

ARTICLE

Received 12 Mar 2015 | Accepted 27 Oct 2015 | Published 21 Jan 2016

DOI: 10.1038/ncomms10023

OPEN

Genetic associations at 53 loci highlight cell types and biological pathways relevant for kidney function

Cristian Pattaro *et al.*[#]

Reduced glomerular filtration rate defines chronic kidney disease and is associated with cardiovascular and all-cause mortality. We conducted a meta-analysis of genome-wide association studies for estimated glomerular filtration rate (eGFR), combining data across 133,413 individuals with replication in up to 42,166 individuals. We identify 24 new and confirm 29 previously identified loci. Of these 53 loci, 19 associate with eGFR among individuals with diabetes. Using bioinformatics, we show that identified genes at eGFR loci are enriched for expression in kidney tissues and in pathways relevant for kidney development and transmembrane transporter activity, kidney structure, and regulation of glucose metabolism. Chromatin state mapping and DNase I hypersensitivity analyses across adult tissues demonstrate preferential mapping of associated variants to regulatory regions in kidney but not extra-renal tissues. These findings suggest that genetic determinants of eGFR are mediated largely through direct effects within the kidney and highlight important cell types and biological pathways.

Correspondence and requests for materials should be addressed to C.P. (email: cristian.pattaro@eurac.edu) or to A.K. (email: anna.koettgen@uniklinik-freiburg.de) or to C.S.F. (email: foxca@nhlbi.nih.gov).

[#]A full list of authors and their affiliations appears at the end of the paper.

Chronic kidney disease (CKD) is a global public health problem^{1–3}, and is associated with an increased risk for cardiovascular disease, all-cause mortality and end-stage renal disease^{4,5}. Few new therapies have been developed to prevent or treat CKD over the past two decades^{1,6}, underscoring the need to identify and understand the underlying mechanisms of CKD.

Prior genome-wide association studies (GWAS) have identified multiple genetic loci associated with CKD and estimated glomerular filtration rate (eGFR), a measure of the kidney's filtration ability that is used to diagnose and stage CKD^{7–15}. Subsequent functional investigations point towards clinically relevant novel mechanisms in CKD that were derived from initial GWAS findings¹⁶, providing proof of principle that locus discovery through large-scale GWAS efforts can translate to new insights into CKD pathogenesis.

To identify additional genetic variants associated with eGFR and guide future experimental studies of CKD-related mechanisms, we have now performed GWAS meta-analyses in up to 133,413 individuals, more than double the sample size of previous studies. Here we describe multiple novel genomic loci associated with kidney function traits and provide extensive locus characterization and bioinformatics analyses, further delineating the physiologic basis of kidney function.

Results

Stage 1 discovery analysis. We analysed associations of eGFR based on serum creatinine (eGFRcrea), cystatin C (eGFRcys, an additional, complementary biomarker of renal function) and CKD (defined as eGFRcrea <60 ml min⁻¹ per 1.73 m²) with ~2.5 million autosomal single-nucleotide polymorphisms (SNPs) in up to 133,413 individuals of European ancestry from 49 predominantly population-based studies (Supplementary Table 1). Results from discovery GWAS meta-analysis are publicly available at <http://fox.nhlbi.nih.gov/CKDGen/>. We performed analyses in each study sample in the overall population and stratified by diabetes status, since genetic susceptibility to CKD may differ in the presence of this strong clinical CKD risk factor. Population stratification did not impact our results as evidenced by low genomic inflation factors in our meta-analyses, which ranged from 1.00 to 1.04 across all our analyses (Supplementary Fig. 1).

In addition to confirming 29 previously identified loci^{7–9} (Fig. 1a; Supplementary Table 2), we identified 48 independent novel loci (Supplementary Table 3) where the index SNP, defined as the variant with the lowest *P* value in the region, had an association *P* value <1.0 × 10⁻⁶. Of these 48 novel SNPs, 21 were genome-wide significant with *P* values <5.0 × 10⁻⁸. Overall, 43 SNPs were identified in association with eGFRcrea (nine in the non-diabetes sample), one with eGFRcys and four with CKD, as reported in Supplementary Table 3. Manhattan plots for CKD, eGFRcys and eGFRcrea in diabetes are shown in Fig. 1b,c and Supplementary Fig. 2, respectively.

Stage-2 replication. Novel loci were tested for replication in up to 42,166 additional European ancestry individuals from 15 studies (Supplementary Table 1). Of the 48 novel candidate SNPs submitted to replication, 24 SNPs demonstrated a genome-wide significant combined stage 1 and 2 *P* value <5.0 × 10⁻⁸ (Table 1). Of these, 23 fulfilled additional replication criteria (*q*-value <0.05 in stage 2). Only rs6795744 at the *WNT7A* locus demonstrated suggestive replication (*P* value <5.0 × 10⁻⁸, *q*-value >0.05). Because serum creatinine is used to estimate eGFRcrea, associated genetic loci may be relevant to creatinine production or metabolism rather than kidney function *per se*. For this reason, we contrasted associations of eGFRcrea versus eGFRcys, the latter estimated from an alternative and

creatinine-independent biomarker of GFR (Supplementary Fig. 3; Supplementary Table 4). The majority of loci (22/24) demonstrated consistent effect directions of their association with both eGFRcrea and eGFRcys.

Association plots of the 24 newly identified genomic regions that contain a replicated or suggestive index SNP appear in Supplementary Fig. 4. The odds ratio for CKD for each of the novel loci ranged from 0.93 to 1.06 (Supplementary Table 4). As evidenced by the relatively small effect sizes, the proportion of phenotypic variance of eGFRcrea explained by all new and known loci was 3.22%: 0.81% for the newly uncovered loci and 2.41% for the already known loci.

Associations stratified by diabetes and hypertension status. The effects of the 53 known and novel loci in individuals with (stage 1 + stage 2 *n* = 16,477) and without (stage 1 + stage 2 *n* = 154,881) diabetes were highly correlated (correlation coefficient: 0.80; 95% confidence interval: 0.67, 0.88; Supplementary Fig. 5) and of similar magnitude (Fig. 2; Supplementary Table 5), suggesting that identification of genetic loci in the overall population may also provide insights into loci with potential importance among individuals with diabetes. The previously identified *UMOD* locus showed genome-wide significant association with eGFRcrea among those with diabetes (Supplementary Fig. 2; rs12917707, *P* value = 2.5 × 10⁻⁸), and six loci (*NFKB1*, *UNCX*, *TSPAN9*, *AP5B1*, *SIPA1L3* and *PTPRO*) had nominally significant associations with eGFRcrea among those with diabetes. Of the previously identified loci, 13 demonstrated nominal associations among those with diabetes, for a total of 19 loci associated with eGFRcrea in diabetes.

Exploratory comparison of the association effect sizes in subjects with and without hypertension based on our previous work⁷ showed that novel and known loci are also similarly associated with eGFRcrea among individuals with and without hypertension (Supplementary Fig. 6).

Tests for SNP associations with related phenotypes. We tested for overlap with traits that are known to be associated with kidney function in the epidemiologic literature by investigating SNP associations with systolic and diastolic blood pressure¹⁷, myocardial infarction¹⁸, left ventricular mass¹⁹, heart failure²⁰, fasting glucose²¹ and urinary albumin excretion (CKDGen Consortium, personal communication). We observed little association of the 24 novel SNPs with other kidney function-related traits, with only 2 out of 165 tests reaching the Bonferroni significance level of 0.0003 (see Methods and Supplementary Table 6).

To investigate whether additional traits are associated with the 24 new eGFR loci, we queried the NHGRI GWAS catalog (www.genome.gov). Overall, nine loci were previously identified in association with other traits at a *P* value of 5.0 × 10⁻⁸ or lower (Supplementary Table 7), including body mass index (*ETV5*) and serum urate (*INHBC*, *A1CF* and *AP5B1*).

Trans-ethnic analyses. To assess the generalizability of our findings across ethnicities, we evaluated the association of the 24 newly identified loci with eGFRcrea in 16,840 participants of 12 African ancestry population studies (Supplementary Table 8) and in up to 42,296 Asians from the AGEN consortium¹¹ (Supplementary Table 9). Seven SNPs achieved nominal direction-consistent significance (*P* < 0.05) in AGEN, and one SNP was significant in the African ancestry meta-analysis (Supplementary Table 9). Random-effect meta-analysis showed that 12 loci (*SDCCAG8*, *LRP2*, *IGFBP5*, *SKIL*, *UNCX*, *KBTBD2*, *A1CF*, *KCNQ1*, *AP5B1*, *PTPRO*, *TP53INP2* and *BCAS1*) had fully consistent effect direction across the three ethnic groups




Figure 1 | Discovery stage genome-wide association analysis. Manhattan plots for eGFRcrea, CKD and eGFRcys. Previously reported loci are highlighted in light blue (grey labels). **(a)** Novel loci uncovered for eGFRcrea in the overall and in the non-diabetes groups are highlighted in blue and green, respectively. **(b)** Results from CKD analysis with highlighted known and novel loci for eGFRcrea. **(c)** Results from eGFRcys with highlighted known and novel loci for eGFRcrea and known eGFRcys loci.

Table 1 | The 24 novel SNPs associated with eGFRcrea in European ancestry individuals.

SNP ID*	Chr.	Position (bp) [†]	Locus name [‡]	Effect/Non effect allele (EAF)	SNP function [§]	Stage 1 (discovery)		Stage 2 (replication)		Combined analysis [¶]		
						Beta	P value	Beta	q-value	Beta	P value [#]	I ² % ^{**}
The eight loci whose smallest P value was observed in the 'no diabetes' group												
rs3850625	1	201,016,296	CACNA1S	A/G (0.12)	Exonic, nonsyn. SNV	0.0080	2.55E-09	0.0071	5.46E-03	0.0083	6.82E-11	0
rs2712184	2	217,682,779	IGFBP5	A/C (0.58)	Intergenic	-0.0049	1.65E-08	-0.0055	2.06E-03	-0.0053	1.33E-10	0
rs9682041	3	170,091,902	SKIL	T/C (0.87)	Intronic	-0.0067	1.36E-07	-0.0046	2.33E-02	-0.0068	2.58E-08	2
rs10513801	3	185,822,353	ETV5	T/G (0.87)	Intronic	0.0070	3.80E-09	0.0046	1.79E-02	0.0072	1.03E-09	0
rs10994860	10	52,645,424	AICF	T/C (0.19)	UTR5	0.0075	1.00E-11	0.0061	5.46E-03	0.0077	1.07E-12	2
rs163160	11	2,789,955	KCNQ1	A/G (0.82)	Intronic	0.0067	9.02E-09	0.0050	9.89E-03	0.0065	2.26E-09	14
rs164748	16	89,708,292	DPEP1	C/G (0.53)	Intergenic	0.0047	9.92E-09	0.0019	4.19E-02	0.0046	1.95E-08	17
rs8091180	18	77,164,243	NFATC1	A/G (0.56)	Intronic	-0.0054	1.43E-08	-0.0052	5.46E-03	-0.0060	1.28E-09	0
The 16 loci whose smallest P value was observed in the 'overall' group												
rs12136063	1	110,014,170	SYPL2	A/G (0.70)	Intronic	0.0049	2.33E-07	0.0028	2.31E-02	0.0045	4.71E-08	0
rs2802729	1	243,501,763	SDCCAG8	A/C (0.43)	Intronic	-0.0050	7.37E-08	-0.0029	2.05E-02	-0.0046	2.20E-08	9
rs4667594	2	170,008,506	LRP2	A/T (0.53)	Intronic	-0.0045	2.37E-07	-0.0043	5.62E-03	-0.0044	3.52E-08	4
rs6795744††	3	13,906,850	WNT7A	A/G (0.15)	Intronic	0.0071	9.60E-09	0.0019	5.15E-02	0.0060	3.33E-08	18
rs228611	4	103,561,709	NFKB1	A/G (0.47)	Intronic	-0.0055	4.66E-10	-0.0060	8.91E-04	-0.0056	3.58E-12	3
rs7759001	6	27,341,409	ZNF204	A/G (0.76)	ncRNA intronic	-0.0053	2.64E-07	-0.0045	9.10E-03	-0.0051	1.75E-08	0
rs10277115	7	1,285,195	UNCX	A/T (0.23)	Intergenic	0.0095	1.05E-10	0.0079	9.03E-04	0.0090	8.72E-14	0
rs3750082	7	32,919,927	KBTD2	A/T (0.33)	Intronic	0.0049	2.52E-07	0.0031	1.96E-02	0.0045	3.22E-08	2
rs6459680	7	156,258,568	RNF32	T/G (0.74)	Intergenic	-0.0065	1.96E-10	-0.0019	4.62E-02	-0.0055	1.07E-09	0
rs4014195	11	65,506,822	AP5B1	C/G (0.64)	Intergenic	0.0061	2.19E-11	0.0034	1.42E-02	0.0055	1.10E-11	0
rs10491967	12	3,368,093	TSPAN9	A/G (0.10)	Intronic	-0.0092	3.03E-10	-0.0106	3.93E-04	-0.0095	5.18E-14	0
rs7956634	12	15,321,194	PTPRO	T/C (0.81)	Intronic	-0.0068	2.46E-09	-0.0069	1.51E-03	-0.0068	7.17E-12	0
rs1106766	12	57,809,456	INHBC	T/C (0.22)	Intergenic	0.0062	4.67E-08	0.0058	8.79E-03	0.0061	2.41E-09	9
rs11666497	19	38,464,262	SIPA1L3	T/C (0.18)	Intronic	-0.0064	8.58E-08	-0.0041	1.53E-02	-0.0058	4.25E-08	24
rs6088580	20	33,285,053	TP53INP2	C/G (0.47)	Intergenic	-0.0055	7.17E-10	-0.0027	2.31E-02	-0.0049	1.79E-09	0
rs17216707	20	52,732,362	BCAS1	T/C (0.79)	Intergenic	-0.0084	5.96E-13	-0.0051	6.69E-03	-0.0077	8.83E-15	1
bp, basepairs; Chr, chromosome; EAF, effect allele frequency; eGFRcrea, eGFR based on serum creatinine; GWAS, genome-wide association studies; SNP, single-nucleotide polymorphism; UTR, untranslated region.												
*SNPs are grouped by the stratum where the smallest P value in the discovery and combined analysis was observed. In the 'no diabetes' group, sample size/number of studies were equal to 118,448/45, 36,433/13 and 154,881/58, in the discovery, replication and combined analyses, respectively. In the 'overall' group, the numbers for the three analyses were equal to 133,413/48, 42,116/14 and 175,579/62, respectively.												
†On the basis of RefSeq genes (build 37).												
‡Conventional locus name based on relevant genes in the region as identified by bioinformatic investigation (Supplementary Table 12) or closest gene. A complete overview of the genes in each locus is given in the regional association plots (Supplementary Fig. 4).												
§SNP function is derived from NCBI RefSeq genes and may not correspond to the named gene.												
Twice genomic-control (GC) corrected P value from discovery GWAS meta-analysis: at the individual study level and after the meta-analysis.												
¶For random-effect estimate, see Supplementary Table 4.												
#P value of the meta-analysis of the twice GC-corrected discovery meta-analysis results and replication studies.												
**Between-study heterogeneity, as assessed by the I ² . Q statistic P value > 0.05 for all SNPs, except rs11666497 (SIPA1L3, P value = 0.04).												
††For this SNP, the conditions for replication were not all met (q-value > 0.05 in the replication stage).												

(Supplementary Fig. 7), suggesting that our findings can likely be generalized beyond the European ancestry group.

To identify additional potentially associated variants and more formally evaluate trans-ethnic heterogeneity of the loci identified through meta-analysis in European ancestry populations, we performed a trans-ethnic meta-analysis²², combining the 12 African ancestry studies with the 48 European Ancestry studies used in the discovery analysis of eGFRcrea. Of the 24 new loci uncovered for eGFRcrea, 15 were also genome-wide significant in the trans-ethnic meta-analysis (defined as log₁₀ Bayes Factor > 6, Supplementary Table 10), indicating that for most of these loci, there is little to no allelic effect heterogeneity across the two ethnic groups. No additional loci were significantly associated with log₁₀ Bayes Factor > 6.

Bioinformatic and functional characterization of new loci. We used several techniques to prioritize and characterize genes underlying the identified associations, uncover connections between associated regions, detect relevant tissues and assign functional annotations to associated variants. These included expression quantitative trait loci (eQTL) analyses, pathway analyses, DNase I hypersensitivity site (DHS) mapping, chromatin mapping, manual curation of genes in each region and zebrafish knockdown.

eQTL analysis. We performed eQTL analysis using publically available eQTL databases (see Methods). These analyses

connected novel SNPs to transcript abundance of *SYPL2*, *SDCCAG8*, *MANBA*, *KBTD2*, *PTPRO* and *SPATA33* (*C16orf55*), thereby supporting these as potential candidate genes in the respective associated regions (Supplementary Table 11).

Pathway analyses. We used a novel method, Data-driven Expression Prioritized Integration for Complex Traits (DEPICT)²³, to prioritize genes at associated loci, to test whether genes at associated loci are highly expressed in specific tissues or cell types and to test whether specific biological pathways and gene sets are enriched for genes in associated loci. On the basis of all SNPs with eGFRcrea association P values < 10⁻⁵ in the discovery meta-analysis, representing 124 independent regions, we identified at least one significantly prioritized gene in 49 regions, including in 9 of the 24 novel genome-wide significant regions (Supplementary Table 12). Five tissue and cell type annotations were enriched for expression of genes from the associated regions, including the kidney and urinary tract, as well as hepatocytes and adrenal glands and cortex (Fig. 3a; Supplementary Table 13). Nineteen reconstituted gene sets showed enrichment of genes mapping into the associated regions at a permutation P value < 10⁻⁵ (Supplementary Table 14; Fig. 4), highlighting processes related to renal development, kidney transmembrane transporter activity, kidney and urogenital system morphology, regulation of glucose metabolism, as well as specific protein complexes important in renal development.




Figure 2 | Association eGFRcrea loci in subjects with and without diabetes. Novel (a) and known (b) loci were considered. Displayed are effects and their 95% confidence intervals on ln(eGFRcrea). Results are sorted by increasing effects in the diabetes group. The majority of loci demonstrated similar effect sizes in the diabetes as compared with non-diabetes strata. SNP-specific information and detailed sample sizes are reported in Supplementary Table 5.

DNase I hypersensitivity and H3K4m3 chromatin mark analyses.

To evaluate whether eGFRcrea-associated SNPs map into gene regulatory regions and to thereby gain insight into their potential function, we evaluated the overlap of independent eGFRcrea-associated SNPs with P values $< 10^{-4}$ (or their proxies) with DHSs using publicly available data from the Epigenomics Roadmap Project and ENCODE for 123 cell types (see Methods). DHSs mark accessible chromatin regions where transcription may occur. Compared with a set of control SNPs (see Methods), eGFRcrea-associated SNPs were significantly more likely to map to DHS in six specific tissues or cell types (Fig. 3b), including adult human renal cortical epithelial cells, adult renal proximal tubular epithelial cells, H7 embryonic stem cells (differentiated 2 days), adult human renal epithelial cells, adult small airway epithelial cells and amniotic epithelial cells. No significant enrichment was observed for adult renal glomerular endothelial cells, the only other kidney tissue evaluated.

Next, we analysed the overlap of the same set of SNPs with H3K4me3 chromatin marks, promoter-specific histone modifications associated with active transcription²⁴, in order to gather more information about cell-type specific regulatory potential of eGFRcrea-associated SNPs. Comparing 33 available adult-derived cell types, we found that eGFRcrea-associated SNPs showed the most significant overlap with H3K4me3 peaks in adult kidney (P value = 0.0029), followed by liver (P value = 0.0117), and rectal mucosa (P value = 0.0445). Taken together, these findings are suggestive of cell-type-specific regulatory roles for eGFR loci, with greatest specificity for the kidney.

Chromatin annotation maps. In addition to assessing individual regulatory marks separately, we annotated the known and replicated novel SNPs, as well as their perfect proxies in a complementary approach. Chromatin annotation maps were generated integrating > 10 epigenetic marks from cells derived from adult human kidney tissue and a variety of non-renal tissues from the ENCODE project (see Methods). The proportion of variants to which a function could be assigned was significantly higher when using chromatin annotation maps from renal tissue compared with using maps that investigated the same epigenetic marks in other non-renal tissues (Fig. 3c), again indicating that eGFRcrea associated SNPs are, or tag, kidney-specific regulatory variants. The difference between kidney and non-renal tissues was particularly evident for marks that define enhancers: the proportion of SNPs mapping to weak and strong enhancer regions in the kidney tissue was higher than in all non-kidney tissues (Fishers' exact test P values from 3.1×10^{-3} to 7.9×10^{-6} , multiple testing threshold $\alpha = 5.6 \times 10^{-3}$).

Functional characterization of new loci. To prioritize genes for functional studies, we applied gene prioritization algorithms including GRAIL²⁵, DEPICT and manual curation of selected genes in each region (Supplementary Table 12). For each region, gene selection criteria were as follows: (1) either GRAIL P value < 0.05 or DEPICT false discovery rate (FDR) < 0.05 ; (2) the effect of a given allele on eGFRcrea and on eGFRcys was direction-consistent and their ratio was between 0.2 and 5




Figure 3 | Bioinformatic analysis of eGFR-associated SNPs. Connection of eGFR-associated SNPs to gene expression and variant function across a variety of tissues, pathways and regulatory marks was considered. **(a)** The DEPICT method shows that implicated eGFR-associated genes are highly expressed in particular tissues, including kidney and urinary tract. Shown are permutation test P values (see Methods). **(b)** Enrichment of eGFRcrea-associated SNPs in DHS according to discovery P value threshold. SNPs from the eGFR discovery genome-wide scan meeting a series of P value thresholds in the range 10^{-4} – 10^{-16} preferentially map to DHSs, when compared with a set of control SNPs, in 6 of 123 cell types. Represented are main effects odds ratios from a logistic mixed effect model. Cell types indicated with coloured lines had nominally significant enrichment (* indicate P values < 0.05) at the P value $< 10^{-16}$ threshold and/or were derived from renal tissues (H7esDiffa2d: H7 embryonic stem cells, differentiated 2 days with BMP4, activin A and bFGF; Hae, amniotic epithelial cells; Hrc, renal cortical epithelial cells; Hre, renal epithelial cells; Hrgec, renal glomerular endothelial cells; Rptec, renal proximal tubule epithelial cells; Saec, small airway epithelial cells). **(c)** ENCODE/Chromatin ChIP-seq mapping: known and replicated novel eGFRcrea-associated SNPs and their perfect proxies were annotated based on genomic location using chromatin annotation maps from different tissues including adult kidney epithelial cells. P values from Fisher's exact tests for 2×2 tables are reported (significance level = 5.6×10^{-3} , see Methods). There is significant enrichment of variants mapping to enhancer regions specifically in kidney but not other non-renal tissues.

(to ensure relative homogeneity of the beta coefficients); (3) nearest gene if the signal was located in a region containing a single gene. Using this approach, *NFKB1*, *DPEP1*, *TSPAN9*, *NFATC1*, *WNT7A*, *PTPRO*, *SYPL2*, *UNCX*, *KBTBD2*, *SKIL* and *AICF* were prioritized as likely genes underlying effects at the new loci (Supplementary Table 12).

We investigated the role of these genes during vertebrate kidney development by examining the functional consequences of gene knockdown in zebrafish embryos utilizing antisense morpholino oligonucleotide (MO) technology. After knockdown, we assessed the expression of established renal markers *pax2a* (global kidney), *nephrin* (podocytes) and *slc20a1a* (proximal tubule) at 48 hours post fertilization by *in situ* hybridization¹². In all cases, morphant embryos did not display significant gene expression defects compared with controls (Supplementary Table 15).

Discussion

We identified 24 new loci in association with eGFR and confirmed 29 previously identified loci. A variety of

complementary analytic, bioinformatic and functional approaches indicate enrichment of implicated gene products in kidney and urinary tract tissues. A greater proportion of the lead SNPs or their perfect proxies map into gene regulatory regions, specifically enhancers, in adult renal tissues compared with non-renal tissues. In addition to the importance in the adult kidney, our results indicate a role for kidney function variants during development.

We extend our previous findings, as well as those from other groups^{7–13} by identifying > 50 genomic loci for kidney function, many of which were not previously known to be connected to kidney function and disease. Using a discovery data set that is nearly double in size to our prior effort⁷, we are now able to robustly link associated SNPs to kidney-specific gene regulatory function. Our work further exemplifies the continued value of increasing the sample size of GWAS meta-analyses to uncover additional loci and gain novel insights into the mechanisms underlying common phenotypes²⁶.

There are several messages from our work. First, many of the genetic variants associated with eGFR appear to affect processes specifically within the kidney. The kidney is a highly vascular and




Figure 4 | Gene set overlap analysis. The 19 reconstituted gene sets with P value $< 10^{-5}$ were considered. Their overlap was estimated by computing the pairwise Pearson correlation coefficient ρ between each pair of gene sets followed by discretization into one of three bins: $0.3 \leq \rho < 0.5$, low overlap; $0.5 \leq \rho < 0.7$, medium overlap; $\rho \geq 0.7$, high overlap. Overlap is shown by edges between gene set nodes and edges representing overlap corresponding to $\rho < 0.3$ are not shown. The network was drawn with Cytoscape⁴⁸.

metabolically active organ that receives 20% of all cardiac output, contains an extensive endothelium-lined capillary network, and is sensitive to ischaemic and toxic injury. As a result, hypertension, cardiovascular diseases and diabetes each affect renal hemodynamics and contribute to kidney injury. However, many of the eGFR-associated SNPs in our GWAS could be assigned gene regulatory function specifically in the kidney and its epithelial cells, but not in human glomerular endothelial cells or the general vasculature. In addition, variants associated with eGFR were not associated with vascular traits, such as blood pressure or myocardial infarction. Taken together, these findings suggest that genetic determinants of eGFR may be mediated largely through direct effects within the kidney.

Second, despite the specificity related to renal processes, we also identified several SNPs that are associated with eGFR in diabetes, and our pathway analyses uncovered gene sets associated with glucose transporter activity and abnormal glucose homeostasis. Uncovering *bona fide* genetic loci for diabetic CKD has been difficult. We have now identified a total of 19 SNPs that demonstrate at least nominal association with eGFR in diabetes. The diabetes population is at particularly high risk of CKD, and identifying kidney injury pathways may help develop new treatments for diabetic CKD.

Finally, even though CKD is primarily a disease of the elderly, our pathway enrichment analyses highlight developmental processes relevant to the kidney and the urogenital tract. Kidney disease has been long thought to have developmental origins, in part related to early programming (Barker hypothesis)²⁷, low birth weight, nephron endowment and early growth and early-life nutrition²⁸. Our pathway enrichment analyses suggest that developmental pathways such as placental morphology, kidney weight and embryo size, as well as protein complexes of importance in renal development may in part contribute to the developmental origins of CKD.

A limitation of our work is that causal variants and precise molecular mechanisms underlying the observed associations were not identified and will require additional experimental follow-up projects. Our attempt to gain insights into potentially causal genes through knockdown in zebrafish did not yield any clear CKD candidate gene, although the absence of a zebrafish

phenotype upon gene knockdown does not mean that the gene cannot be the one underlying the observed association signal in humans. Finally, our conclusions that eGFR_{crea}-associated SNPs regulate the expression of nearby genes specifically in kidney epithelial cells are based on DHSs, H3K4me3 chromatin marks and chromatin annotation maps. Since these analyses rely mostly on variant positions, additional functional investigation such as luciferase assay that assess transcriptional activity more directly are likely to gain additional insights into the variants' mechanism of action.

The kidney specificity for loci we identified may have important translational implications, particularly since our DHS and chromatin annotation analyses suggest that at least a set of gene regulatory mechanisms is important in the adult kidney. Kidney-specific pathways are important for the development of novel therapies to prevent and treat CKD and its progression with minimal risk of toxicity to other organs. Finally, the biologic insights provided by these new loci may help elucidate novel mechanisms and pathways implicated not only in CKD but also of kidney function in the physiological range.

In conclusion, we have confirmed 29 genomic loci and identified 24 new loci in association with kidney function that together highlight target organ-specific regulatory mechanisms related to kidney function.

Methods

Overview. This was a collaborative meta-analysis with a distributive data model. Briefly, an analysis plan was created and circulated to all participating studies. Studies then uploaded study-specific data centrally; files were cleaned, and a specific meta-analysis for each trait was performed. Details regarding each step are provided below. All participants in all discovery and replication studies provided informed consent. Each study had its research protocol approved by the local ethics committee.

Phenotype definitions. Serum creatinine was measured in each discovery and replication study as described in Supplementary Tables 16 and 17, and statistically calibrated to the US nationally representative National Health and Nutrition Examination Study data in all studies to account for between-laboratory variation^{9,29,30}. eGFR_{crea} was estimated using the four-variable Modification of Diet in Renal Disease Study Equation. Cystatin C, an alternative biomarker for kidney function, was measured in a sub-set of participating studies. eGFR_{cys} was estimated as $76.7 \times (\text{serum cystatin C})^{-1.19}$ (ref. 31). eGFR_{crea} and eGFR_{cys} values

$<15 \text{ ml min}^{-1} \text{ per } 1.73 \text{ m}^2$ were set to 15, and those >200 were set to $200 \text{ ml min}^{-1} \text{ per } 1.73 \text{ m}^2$. CKD was defined as $\text{eGFR}_{\text{crea}} < 60 \text{ ml min}^{-1} \text{ per } 1.73 \text{ m}^2$.

Diabetes was defined as fasting glucose $\geq 126 \text{ mg dl}^{-1}$, pharmacologic treatment for diabetes or by self-report. In all studies, diabetes and kidney function were assessed at the same point in time.

Genotypes. Genotyping was conducted in each study as specified in Supplementary Tables 18 and 19. After applying appropriate quality filters, 45 studies used markers of highest quality to impute ~ 2.5 million SNPs, based on European-ancestry haplotype reference samples (HapMap II CEU). Four studies based their imputation on the 1000 Genomes Project data. Imputed genotypes were coded as the estimated number of copies of a specified allele (allelic dosage).

Genome-wide association analysis. By following a centralized analysis plan, each study regressed sex- and age-adjusted residuals of the logarithm of $\text{eGFR}_{\text{crea}}$ or eGFR_{cys} on SNP dosage levels. Logistic regression of CKD status was performed on SNP dosage levels adjusting for sex and age. For all traits, adjustment for appropriate study-specific features, including study site and genetic principal components was included in the regression and family-based studies appropriately accounted for relatedness.

Stage 1 discovery meta-analysis. GWAS of $\text{eGFR}_{\text{crea}}$ were contributed by 48 studies (total sample size, $N = 133,413$); 45 studies contributed GWAS data for the non-diabetes subgroup ($N = 118,448$) and 39 for the diabetes group ($N = 11,522$). GWAS of CKD were comprised by 43 studies, for a total sample size of 117,165, including 12,385 CKD cases. GWAS of eGFR_{cys} were comprised by 16 studies for a total sample size of 32,834. All GWAS files underwent quality control using the GWAtoolbox package³² in R, before including them into the meta-analysis. Genome-wide meta-analysis was performed with the software METAL³³, assuming fixed effects and using inverse-variance weighting. The genomic inflation factor λ was estimated for each study as the ratio between the median of all observed test statistics ($b/\text{s.e.}$)² and the expected median of a χ^2 with 1 degree of freedom, with b and s.e. representing the effect of each SNP on the phenotype and its standard error, respectively³⁴. Genomic-control (GC) correction was applied to P values and s.e. 's in case of $\lambda > 1$ (first GC correction). SNPs with an average minor allele frequency (MAF) of ≥ 0.01 were used for the meta-analysis. To limit the possibility of false positives, after the meta-analysis, a second GC correction on the aggregated results was applied. Between-study heterogeneity was assessed through the I^2 statistic.

After removing SNPs with MAF of <0.05 and which were available in $<50\%$ of the studies, SNPs with a P value of $\leq 10^{-6}$ were selected and clustered into independent loci through LD pruning based on an r^2 of ≤ 0.2 within a window of $\pm 1 \text{ MB}$ to each side of the index SNP. After removing loci containing variants that have been previously replicated at a P value of 5.0×10^{-8} (refs 7,8), the SNP with the lowest P value within each locus was selected for replication ('index SNP'). If a SNP had an association P value of $\leq 10^{-6}$ with more than one trait, the trait where the SNP had the lowest P value was selected as discovery trait/stratum. Altogether, this resulted in 48 SNPs: 34 from $\text{eGFR}_{\text{crea}}$, 9 from $\text{eGFR}_{\text{crea}}$ among those without diabetes, 4 from CKD and 1 from eGFR_{cys} .

Stage 2 replication analysis. *In silico* replication analysis for any of the studied traits was carried out using eight independent studies whose genotyping platforms are provided in Supplementary Table 19. *De novo* genotyping was performed in seven additional studies ($N = 22,850$ individuals) of European ancestry (Supplementary Table 20), including the Bus Santé, ESTHER, KORA-F3 (subset of F3 without GWAS), KORA-F4 (subset of F4 without GWAS), Ogliastro Genetic Park (OGP, without Talana whose GWAS was included in the discovery analysis), SAPHIR and SKIPOGH studies (Supplementary Table 20). Summarizing all *in silico* and *de novo* replication studies (Supplementary Table 1), replication data for $\text{eGFR}_{\text{crea}}$ were contributed by 14 studies (total sample size = 42,166), which also contributed $\text{eGFR}_{\text{crea}}$ results from non-diabetes (13 studies, $N = 36,433$) and diabetes samples (13 studies, $N = 4,955$). Thirteen studies contributed replication data on CKD ($N = 33,972$; 4,245 CKD cases; studies with <50 CKD cases were excluded) and five on eGFR_{cys} ($N = 14,930$).

Association between $\text{eGFR}_{\text{crea}}$, CKD and eGFR_{cys} and each of the 48 SNPs in the replication studies was assessed using the same analysis protocol detailed for the discovery studies above. Quality control of the replication files was performed with the same software as described above.

We performed a combined fixed-effect meta-analysis of the double-GC corrected results from the discovery meta-analysis and the replication studies, based on inverse-variance weighting. The total sample size in the combined analysis of $\text{eGFR}_{\text{crea}}$ was 175,579 subjects (154,881 in the non-diabetes stratum and 16,477 in the diabetes stratum; the sum of these two sample sizes is smaller than the sample size of the overall analysis because some studies did not contribute both strata), 151,137 samples for CKD (16,630 CKD cases) and 47,764 for eGFR_{cys} . Three criteria were used to ensure validity of novel loci declared as significant: (1) P value from the combined meta-analysis $\leq 5.0 \times 10^{-8}$ in accordance with previously published guidelines³⁵; (2) direction-consistent associations of the beta coefficients in stage 1 and stage 2 (one-sided P values were

estimated to test for consistent effect direction with the discovery stage); (3) q -value < 0.05 in the replication stage. Q -values were estimated using the package QVALUE³⁶ in R. The tuning parameter lambda for the estimation of the overall proportion of true null hypotheses, π_0 , was estimated using the bootstrap method³⁷. When the third criterion was not satisfied, the locus was declared 'suggestive'.

Power analysis. With the sample size achieved in the combined analysis of stage 1 and stage 2 data, the power to assess replication at the canonical genome-wide significance level of 5.0×10^{-8} was estimated with the software QUANTO³⁸ version 1.2.4, assuming the same MAF and effect size observed in the discovery sample. Power to replicate associations ranged from 87 to 100% for $\text{eGFR}_{\text{crea}}$ associated SNPs (median = 98%), from 72 to 96% for the CKD-associated SNPs, and was equal to 59% for the eGFR_{cys} -associated SNP (Supplementary Table 3).

Associations stratified by diabetes and hypertension status. For all the 24 novel and 29 known SNPs, the difference between the SNP effect on $\text{eGFR}_{\text{crea}}$ in the diabetes versus the non-diabetes groups was assessed by means of a two-sample t -test for correlated data at a significance level of 0.05. We used the following two-sample t -test for correlated data:

$$t = \frac{(b_{\text{DM}} - b_{\text{nonDM}})}{\left\{ \text{s.e.}(b_{\text{DM}})^2 + \text{s.e.}(b_{\text{nonDM}})^2 - 2 \times \rho(b_{\text{DM}}, b_{\text{nonDM}}) \times \text{s.e.}(b_{\text{DM}}) \times \text{s.e.}(b_{\text{nonDM}}) \right\}^{0.5}},$$

where b_{DM} and b_{nonDM} represent the SNP effects on $\log(\text{eGFR}_{\text{crea}})$ in the two groups, s.e. is the standard error of the estimate and $\rho(\cdot)$ indicates the correlation between effects in the two groups, which was estimated as 0.044 by sampling 100,000 independent SNPs from our DM and nonDM GWAS, after removing known and novel loci associated with $\text{eGFR}_{\text{crea}}$. For a large sample size, as in our case, t follows a standard normal distribution.

A similar analysis was performed to compare results in subjects with and without hypertension, based on results from our previous work⁷. The correlation between the two strata was of 0.01.

Proportion of phenotypic variance explained. The percent of phenotypic

variance explained by novel and known loci was estimated as $\sum_{i=1}^{53} R_i^2$, where

$R_i^2 = b_i^2 \text{var}(\text{SNP}_i) / \text{var}(y)$ is the coefficient of determination for each of the 53 individual SNPs associated with $\text{eGFR}_{\text{crea}}$ uncovered to date (24 novel and 29 known ones), b_i is the estimated effect of the i^{th} SNP on y , y corresponds to the sex- and age-adjusted residuals of the logarithm of $\text{eGFR}_{\text{crea}}$ and $\text{var}(\text{SNP}_i) = 2 \times \text{MAF}_{\text{SNP}_i} \times (1 - \text{MAF}_{\text{SNP}_i})$ ³⁹. $\text{var}(y)$ was estimated in the ARIC study and all loci were assumed to have independent effects on the phenotype.

Test for SNP associations with related traits. We performed evaluations of SNP association with results generated from consortia investigating other traits. Specifically, we evaluated systolic and diastolic blood pressure in ICBP¹⁷, myocardial infarction in CARDIOGRAM¹⁸, left ventricular mass¹⁹, heart failure²⁰, the urinary albumin to creatinine ratio (CKDGen consortium, personal communication) and fasting plasma glucose in MAGIC²¹. In total, we performed 165 tests, corresponding to 7 traits tested for association against each of the 24 novel SNP, with the exception of myocardial infarction for which results from 3 SNPs were not available (Supplementary Table 6). Significance was evaluated at the Bonferroni corrected level of $0.05/165 = 0.0003$.

Lookup of replicated loci in the NHGRI GWAS catalog. All replicated SNPs, as well as SNPs in LD ($r^2 > 0.2$) within $\pm 1 \text{ MB}$ distance were checked for their association with other traits according to the NHGRI GWAS catalog⁴⁰ (accessed April 14, 2014).

SNP assessments in other ethnic groups. We performed cross-ethnicity SNP evaluations in participants of African ancestry from a meta-analysis of African ancestry individuals and from participants of Asian descent from the AGEN consortium¹¹.

African ancestry meta-analysis. We performed fixed-effect meta-analysis of the genome-wide association data from 12 African ancestry studies (Supplementary Table 8) with imputation to HapMap reference panel, based on inverse-variance weighting using METAL. Only SNPs with MAF ≥ 0.01 and imputation quality $r^2 \geq 0.3$ were considered for the meta-analysis. After meta-analysis, we removed SNPs with MAF < 0.05 and which were available in $<50\%$ of the studies. Statistical significance was assessed at the standard threshold of 5.0×10^{-8} . Genomic control correction was applied at both the individual study level before meta-analysis and after the meta-analysis.

Transethnic meta-analysis. We performed a trans-ethnic meta-analysis of GWAS data from cohorts of different ethnic backgrounds using MANTRA (Meta-Analysis

of Trans-ethnic Association studies) software²². We combined the 48 European ancestry studies that contributed eGFRcrea, which were included in stage 1 discovery meta-analysis, and the 12 African ancestry studies mentioned above for a total sample size of 150,253 samples. We limited our analysis to biallelic SNPs with $MAF \geq 0.01$ and imputation quality $r^2 \geq 0.3$. Relatedness between the 60 studies was estimated using default settings from up to 5.9 million SNPs. Only SNPs that were present in more than 25 European ancestry studies and 6 African ancestry studies (total sample size $\geq 120,000$) were considered after meta-analysis. Genome-wide significance was defined as a \log_{10} Bayes' Factor ($\log_{10}BF$) ≥ 6 (ref. 41).

Gene Relationships Across Implicated Loci (GRAIL). To prioritize the gene(s) most likely to give rise to association signals in a given region, the software GRAIL was used²⁵. The index SNP of all previously known kidney function associated regions, as well as the novel SNPs identified here was used as input, using the CEU HapMap (hg18 assembly) and the functional datasource text_2009_03, established before the publication of kidney function-related GWAS. Results from GRAIL were used to prioritize genes for follow-up functional work.

Expression quantitative trait loci analysis. We identified alias rsIDs and proxies ($r^2 > 0.8$) for our index SNPs using SNAP software across 4 HapMap builds. SNP rsIDs and aliases were searched for primary SNPs and LD proxies against a collected database of expression SNP (eSNP) results. The collected eSNP results met criteria for statistical thresholds for association with gene transcript levels in their respective original analyses (for references see Supplementary Table 11). Correlation of selected eSNPs to the best eSNPs per transcript per expression quantitative trait loci (eQTL) data set were assessed by pairwise LD. All results are reported in Supplementary Table 11.

DEPICT analysis. In this work, we first used PLINK⁴² to identify independently associated SNPs using all SNPs with eGFRcrea association P values $< 10^{-5}$ (HapMap release 27 CEU data⁴³; LD r^2 threshold = 0.01; physical kb threshold = 1,000). We then used the DEPICT method²³ to construct associated regions by mapping genes to independently associated SNPs if they overlapped or resided within LD ($r^2 > 0.5$) distance of a given associated SNP. After merging overlapping regions and discarding regions that mapped within the major histocompatibility complex locus (chromosome 6, base pairs 20,000,000–40,000,000), 124 non-overlapping regions remained that covered a total of 320 genes. Finally, we ran the DEPICT software program on the 124 regions to prioritize genes that may represent promising candidates for experimental follow up studies, identify reconstituted gene sets that are enriched in genes from associated regions and therefore may provide insight into general kidney function biology, and identify tissue and cell-type annotations in which genes from associated regions are highly expressed. Specifically, for each tissue, the DEPICT method performs a t -test comparing the tissue-specific expression of eGFRcrea-associated genes and all other genes. Next, for each tissue, empirical enrichment P values are computed by repeatedly sampling random sets of loci (matched to the actual eGFRcrea loci by gene density) from the entire genome to estimate the empirical mean and s.d. of the enrichment statistic's null distribution. To visualize the nineteen reconstituted gene sets with P value $< 1e-5$ (Fig. 4), we estimated their overlap by computing the pairwise Pearson correlation coefficient ρ between each pair of gene sets followed by discretization into one of three bins; $0.3 \leq \rho < 0.5$, low overlap; $0.5 \leq \rho < 0.7$, medium overlap; $\rho \geq 0.7$, high overlap.

DNase I hypersensitivity analysis. The overlap of SNPs associated with eGFRcrea at $P < 10^{-4}$ with DHSs was examined using publicly available data from the Epigenomics Roadmap Project and ENCODE. In all, DHS mappings were available for 123 mostly adult cells and tissues⁴⁴ (downloaded from <http://hgdownload.cse.ucsc.edu/goldenPath/hg19/encodeDCC/wgEncodeUwDnase/>). The analysis here pertains to DHS's defined as "broad" peaks, which were available as experimental replicates (typically duplicates) for the majority of cells and tissues.

SNPs from our stage 1 eGFRcrea GWAS meta-analysis were first clumped in PLINK⁴² in windows of 100 kb and maximum r^2 of 0.1 using LD relationships from the 1,000 Genomes EUR panel (phase I, v3, 3/14/2012 haplotypes) using a series of P value thresholds (10^{-4} , 10^{-6} , 10^{-8} , ... and 10^{-16}). LD proxies of the index SNPs from the clumping procedure were then identified by LD tagging in PLINK with $r^2 = 0.8$ in windows of 100 kb, again using LD relationships in the 1000G EUR panel, restricted to SNPs with $MAF > 1\%$ and also present in the HapMap2 CEU population. A reference set of control SNPs was constructed using the same clumping and tagging procedures applied to NHGRI GWAS catalog SNPs (available at <http://www.genome.gov/gwastudies/>, accessed 13 March 2013) with discovery P values $< 5.0 \times 10^{-8}$ in European populations. In total, there were 1,204 such reference SNPs after LD pruning. A small number of reference SNPs or their proxies overlapping with the eGFRcrea SNPs or their proxies were excluded. For each cell-type and P value threshold, the enrichment of eGFR SNPs (or their LD proxies) mapping to DHSs relative to the GWAS catalog reference SNPs (or their LD proxies) was expressed as an odds ratio from logistic mixed effect models that treated the replicate peak determinations as random effects (lme4 package in R). Significance for enrichment odds ratio was derived from the significance of beta coefficients for the main effects in the mixed models.

Interrogation of human kidney chromatin annotation maps. Different chromatin modification patterns can be used to generate tissue-specific chromatin-state annotation maps. These can serve as a valuable resource to discover regulatory regions and study their cell-type-specific distributions and activities, which may help with the interpretation especially of intergenic variants identified in association studies⁴⁵. We therefore investigated the genomic mapping of the known and replicated novel index SNPs, as well as their perfect LD proxies ($n = 173$, $r^2 = 1$ for proxies) using a variety of resources, including chromatin maps generated from human kidney tissue cells (HKC-E cells). Chromatin immune-precipitation sequencing (ChIP-seq) data from human kidney samples were generated by NIH Roadmap Epigenomics Mapping Consortium⁴⁶. Briefly, proximal tubule cells derived from an adult human kidney were collected and cross-linked with 1% formaldehyde. Subsequently, ChIP-seq was conducted using whole-cell extract from adult kidney tissue as the input (GSM621638) and assessing the following chromatin marks: H3K36me3 (GSM621634), H3K4me1 (GSM670025), H3K4me3 (GSM621648), H3K9ac (GSM772811) and H3K9me3 (GSM621651). The MACS version 1.4.1 (model-based analysis of ChIP-Seq) peak-finding algorithm was used to identify regions of ChIP-Seq enrichment⁴⁷. A FDR threshold of enrichment of 0.01 was used for all data sets. The resulting genomic coordinates in bed format were further used in ChromHMM v1.06 for chromatin annotation⁴⁵. For comparison, the same genomic coordinates were investigated in chromatin annotation maps of renal tissue, as well as across nine different cell lines from the ENCODE Project: umbilical vein endothelial cells (HUVEC), mammary epithelial cells (HMEC), normal epidermal keratinocytes (NHEK), B-lymphoblastoid cells (GM12878), erythrocytic leukemia cells (K562), normal lung fibroblasts (NHLF), skeletal muscle myoblasts (HSM), embryonic stem cells (H1 ES) and hepatocellular carcinoma cells (HepG2). We tested whether the proportion of SNPs pointing to either strong or weak enhancers in the human kidney tissue cells was different from that of the other nine tissues by means of a Fishers' exact test for 2×2 tables, contrasting each of the nine cell lines listed above against the reference kidney cell line, at a Bonferroni-corrected significance level of $0.05/9 = 5.6 \times 10^{-3}$.

Functional characterization of new loci. Replicated gene regions were prioritized for functional studies using the following criteria: (1) GRAIL identification of a gene in each region of P value < 0.05 or DEPICT, FDR < 0.05 ; (2) an eGFRcrea to eGFRcys ratio between 0.2 and 5 with direction consistency between the beta coefficients; (3) nearest gene if the signal was located in a gene-poor region. The list of genes selected for functional work can be found in Supplementary Table 12. This same prioritization scheme was also used to assign locus names. Morpholino knockdowns were performed in zebrafish.

Zebrafish (strain Tübingen, TU) were maintained according to established Harvard Medical School Institutional Animal Care and Use Committee protocols (protocol # 04626). Male and female fish were mated (age 6–12 months) for embryo production. Embryos were injected at the one-cell stage with MOs (GeneTools) designed to block either the ATG start site or an exon–intron splice site of the target gene (Supplementary Table 21). In cases where human loci are duplicated in zebrafish, both orthologues were knocked down simultaneously by combination MO injection. MOs were injected in escalating doses at concentrations up to 250 μ M. Embryos were fixed in 4% paraformaldehyde at 48 h post fertilization for *in situ* hybridization using published methods (<http://zfin.org/ZFIN/Methods/ThiSeProtocol.html>). Gene expression was visualized using established renal markers *pax2a* (global kidney), *nephrin* (podocytes) and *slc20a1a* (proximal tubule). The number of morphant embryos displaying abnormal gene expression was compared with control embryos by means of a Fisher's exact test.

References

- Eckardt, K. U. *et al.* Evolving importance of kidney disease: from subspecialty to global health burden. *Lancet* **382**, 158–169 (2013).
- El Nahas, M. The global challenge of chronic kidney disease. *Kidney Int.* **68**, 2918–2929 (2005).
- Baumeister, S. E. *et al.* Effect of chronic kidney disease and comorbid conditions on health care costs: A 10-year observational study in a general population. *Am. J. Nephrol.* **31**, 222–229 (2010).
- Fox, C. S. *et al.* Associations of kidney disease measures with mortality and end-stage renal disease in individuals with and without diabetes: a meta-analysis. *Lancet* **380**, 1662–1673 (2012).
- Mahmoodi, B. K. *et al.* Associations of kidney disease measures with mortality and end-stage renal disease in individuals with and without hypertension: a meta-analysis. *Lancet* **380**, 1649–1661 (2012).
- Himmelfarb, J. & Tuttle, K. R. New therapies for diabetic kidney disease. *N. Engl. J. Med.* **369**, 2549–2550 (2013).
- Pattaro, C. *et al.* Genome-wide association and functional follow-up reveals new loci for kidney function. *PLoS Genet.* **8**, e1002584 (2012).
- Kottgen, A. *et al.* New loci associated with kidney function and chronic kidney disease. *Nat. Genet.* **42**, 376–384 (2010).
- Kottgen, A. *et al.* Multiple loci associated with indices of renal function and chronic kidney disease. *Nat. Genet.* **41**, 712–717 (2009).
- Chambers, J. C. *et al.* Genetic loci influencing kidney function and chronic kidney disease. *Nat. Genet.* **42**, 373–375 (2010).

11. Okada, Y. *et al.* Meta-analysis identifies multiple loci associated with kidney function-related traits in east Asian populations. *Nat. Genet.* **44**, 904–909 (2012).
12. Liu, C. T. *et al.* Genetic association for renal traits among participants of African ancestry reveals new loci for renal function. *PLoS Genet.* **7**, e1002264 (2011).
13. Boger, C. A. *et al.* Association of eGFR-Related Loci Identified by GWAS with Incident CKD and ESRD. *PLoS Genet.* **7**, e1002292 (2011).
14. Chasman, D. I. *et al.* Integration of genome-wide association studies with biological knowledge identifies six novel genes related to kidney function. *Hum. Mol. Genet.* **21**, 5329–5343 (2012).
15. Pattaro, C. *et al.* A meta-analysis of genome-wide data from five European isolates reveals an association of COL22A1, SYT1, and GABRR2 with serum creatinine level. *BMC Med. Genet.* **11**, 41 (2010).
16. Trudu, M. *et al.* Common noncoding UMOD gene variants induce salt-sensitive hypertension and kidney damage by increasing uromodulin expression. *Nat. Med.* **19**, 1655–1660 (2013).
17. International Consortium for Blood Pressure Genome-Wide Association Studies *et al.* Genetic variants in novel pathways influence blood pressure and cardiovascular disease risk. *Nature* **478**, 103–109 (2011).
18. Schunkert, H. *et al.* Large-scale association analysis identifies 13 new susceptibility loci for coronary artery disease. *Nat. Genet.* **43**, 333–338 (2011).
19. Vasan, R. S. *et al.* Genetic variants associated with cardiac structure and function: a meta-analysis and replication of genome-wide association data. *JAMA* **302**, 168–178 (2009).
20. Smith, N. L. *et al.* Association of genome-wide variation with the risk of incident heart failure in adults of European and African ancestry: a prospective meta-analysis from the cohorts for heart and aging research in genomic epidemiology (CHARGE) consortium. *Circ. Cardiovasc. Genet.* **3**, 256–266 (2010).
21. Dupuis, J. *et al.* New genetic loci implicated in fasting glucose homeostasis and their impact on type 2 diabetes risk. *Nat. Genet.* **42**, 105–116 (2010).
22. Morris, A. P. Transethnic meta-analysis of genomewide association studies. *Genet. Epidemiol.* **35**, 809–822 (2011).
23. Pers, T. H. *et al.* Biological interpretation of genome-wide association studies using predicted gene functions. *Nat. Commun.* **6**, 5890 (2015).
24. Trynka, G. *et al.* Chromatin marks identify critical cell types for fine mapping complex trait variants. *Nat. Genet.* **45**, 124–130 (2013).
25. Raychaudhuri, S. *et al.* Identifying relationships among genomic disease regions: predicting genes at pathogenic SNP associations and rare deletions. *PLoS Genet.* **5**, e1000534 (2009).
26. Panagiotou, O. A., Willer, C. J., Hirschhorn, J. N. & Ioannidis, J. P. The power of meta-analysis in genome-wide association studies. *Annu. Rev. Genomics Hum. Genet.* **14**, 441–465 (2013).
27. Barker, D. J. The fetal and infant origins of adult disease. *BMJ* **301**, 1111 (1990).
28. Boubred, F. *et al.* Developmental origins of chronic renal disease: an integrative hypothesis. *Int. J. Nephrol.* **2013**, 346067 (2013).
29. Fox, C. S. *et al.* Predictors of new-onset kidney disease in a community-based population. *JAMA* **291**, 844–850 (2004).
30. Coresh, J. *et al.* Calibration and random variation of the serum creatinine assay as critical elements of using equations to estimate glomerular filtration rate. *Am. J. Kidney Dis.* **39**, 920–929 (2002).
31. Stevens, L. A. *et al.* Estimating GFR using serum cystatin C alone and in combination with serum creatinine: a pooled analysis of 3,418 individuals with CKD. *Am. J. Kidney Dis.* **51**, 395–406 (2008).
32. Fuchsberger, C., Taliun, D., Pramstaller, P. P. & Pattaro, C. CKDgen consortium. GWAToolbox: an R package for fast quality control and handling of GWAS meta-analysis data. *Bioinformatics* **28**, 444–445 (2012).
33. Willer, C. J., Li, Y. & Abecasis, G. R. METAL: fast and efficient meta-analysis of genomewide association scans. *Bioinformatics* **26**, 2190–2191 (2010).
34. Devlin, B. & Roeder, K. Genomic control for association studies. *Biometrics* **55**, 997–1004 (1999).
35. Skol, A. D., Scott, L. J., Abecasis, G. R. & Boehnke, M. Joint analysis is more efficient than replication-based analysis for two-stage genome-wide association studies. *Nat. Genet.* **38**, 209–213 (2006).
36. Storey, J. D. A direct approach to false discovery rates. *J. R. Stat. Soc. B* **64**, 479–498 (2002).
37. Storey, J. D., Taylor, J. E. & Siegmund, D. Strong control, conservative point estimation, and simultaneous conservative consistency of false discovery rates: a unified approach. *J. R. Stat. Soc. B* **66**, 187–205 (2004).
38. Gauderman, W. J. Sample size requirements for matched case-control studies of gene-environment interaction. *Stat. Med.* **21**, 35–50 (2002).
39. Rosner, B. in *Fundamentals of Biostatistics* (Thomson-Brooks/Cole, Duxbury, 2006).
40. Welter, D. *et al.* The NHGRI GWAS Catalog, a curated resource of SNP-trait associations. *Nucleic Acids Res.* **42**, D1001–D1006 (2014).
41. Wang, X. *et al.* Comparing methods for performing trans-ethnic meta-analysis of genome-wide association studies. *Hum. Mol. Genet.* **22**, 2303–2311 (2013).
42. Purcell, S. *et al.* PLINK: a tool set for whole-genome association and population-based linkage analyses. *Am. J. Hum. Genet.* **81**, 559–575 (2007).
43. International HapMap Consortium. The International HapMap Project. *Nature* **426**, 789–796 (2003).
44. Maurano, M. T. *et al.* Systematic localization of common disease-associated variation in regulatory DNA. *Science* **337**, 1190–1195 (2012).
45. ENCODE Project Consortium *et al.* An integrated encyclopedia of DNA elements in the human genome. *Nature* **489**, 57–74 (2012).
46. Ko, Y. A. *et al.* Cytosine methylation changes in enhancer regions of core pro-fibrotic genes characterize kidney fibrosis development. *Genome Biol.* **14**, R108 (2013).
47. Zhang, Y. *et al.* Model-based analysis of ChIP-Seq (MACS). *Genome Biol.* **9**, R137 (2008).
48. Shannon, P. *et al.* Cytoscape: a software environment for integrated models of biomolecular interaction networks. *Genome Res.* **13**, 2498–2504 (2003).

Acknowledgements

Study-specific acknowledgements and funding sources for participating studies are reported in Supplementary Note. Zebrafish work was supported by NIH R01DK090311 and R24OD017870 to W.G.

Author contributions

Study design: C Helmer, B Stengel, J Chalmers, M Woodward, P Hamet, G Eiriksdottir, LJ Launer, TB Harris, V Gudnason, JR O'Connell, A Köttgen, E Boerwinkle, WHL Kao, P Mitchell, I Guessous, JM Gaspoz, N Bouatia-Naji, P Froguel, A Metspalu, T Esko, BA Oostra, CM van Duijn, V Emilsson, H Brenner, I Borecki, CS Fox, Q Yang, BK Krämer, PS Wild, BI Freedman, J Ding, Y Liu, AB Zonderman, MK Evans, A Adeyemo, CN Rotimi, D Cusi, P Gasparini, M Ciullo, D Toniolo, C Gieger, C Meisinger, CA Böger, HE Wichmann, T Illig, I Rudan, W März, PP Pramstaller, EP Bottinger, BW Penninx, H Snieder, U Gyllenstein, AF Wright, H Campbell, JF Wilson, SH Wild, GJ Navis, BM Buckley, I Ford, JW Jukema, B Paulweber, L Kedenko, F Kronenberg, K Endlich, R Rettig, R Biffar, H Völzke, JK Fernandes, MM Sale, M Pruijm, GB Ehret, A Tönjes, M Stumvoll, JC Denny, RJ Carroll, N Hastie, O Polasek, PM Ridker, J Viikari, M Kähönen, O Raitakari, T Lehtimäki.

Study management: C Helmer, M Metzger, J Tremblay, J Chalmers, M Woodward, P Hamet, G Eiriksdottir, TB Harris, T Aspelund, V Gudnason, A Parsa, AR Shuldiner, BD Mitchell, E Boerwinkle, J Coresh, WHL Kao, R Schmidt, L Ferrucci, E Rohtchina, JJ Wang, J Attia, P Mitchell, I Guessous, JM Gaspoz, M Bochud, DS Siscovick, O Devuyst, P Froguel, T Esko, BA Oostra, CM van Duijn, V Emilsson, AK Dieffenbach, H Brenner, I Borecki, CS Fox, M Rheinberger, ST Turner, S Kloiber, PS Wild, J Ding, Y Liu, SLR Kardia, AB Zonderman, MK Evans, MC Cornelis, A Adeyemo, CN Rotimi, D Cusi, E Salvi, PB Munroe, P Gasparini, M Ciullo, R Sorice, C Sala, D Toniolo, AW Dreisbach, DI Chasman, C Gieger, C Meisinger, M Waldenberger, HE Wichmann, T Illig, W Koenig, I Rudan, I Kolcic, M Boban, T Zemunik, W März, H Kramer, PP Pramstaller, EP Bottinger, O Gottesman, BW Penninx, H Snieder, JH Smit, AF Wright, H Campbell, JF Wilson, SH Wild, W Lieb, GJ Navis, BM Buckley, I Ford, JW Jukema, A Hofman, OH Franco, M Adam, M Imboden, N Probst-Hensch, B Paulweber, L Kedenko, F Kronenberg, S Coassin, M Haun, HK Kroemer, K Endlich, M Nauck, R Rettig, R Biffar, S Stracke, U Völker, H Wallaschofski, H Völzke, KL Keene, MM Sale, B Ponte, D Ackermann, M Pruijm, GB Ehret, A Tönjes, P Kovacs, JC Denny, RJ Carroll, C Hayward, O Polasek, V Vitart, PM Ridker, J Viikari, M Kähönen, O Raitakari, T Lehtimäki.

Subject recruitment: C Helmer, P Hamet, TB Harris, T Aspelund, V Gudnason, AR Shuldiner, BD Mitchell, J Coresh, WHL Kao, M Cavallieri, R Schmidt, JB Whitfield, NG Martin, L Ferrucci, P Mitchell, I Guessous, DS Siscovick, O Devuyst, A Metspalu, BA Oostra, CM van Duijn, BK Krämer, ST Turner, S Kloiber, PS Wild, BI Freedman, MA McEvoy, RJ Scott, AB Zonderman, MK Evans, GC Curhan, A Adeyemo, CN Rotimi, D Cusi, A Lupo, G Gambaro, P d'Adamo, A Robino, S Ulivi, D Ruggiero, M Ciullo, R Sorice, D Toniolo, C Gieger, C Meisinger, CA Böger, HE Wichmann, T Illig, W Koenig, I Rudan, I Kolcic, M Boban, T Zemunik, PP Pramstaller, EP Bottinger, BW Penninx, Å Johansson, I Persico, M Pirastu, JF Wilson, SH Wild, A Franke, G Jacobs, GJ Navis, IM Leach, BM Buckley, I Ford, JW Jukema, N Probst-Hensch, B Paulweber, L Kedenko, F Kronenberg, R Rettig, R Biffar, S Stracke, H Völzke, P Muntner, JK Fernandes, MM Sale, B Ponte, D Ackermann, M Pruijm, GB Ehret, A Tönjes, JC Denny, RJ Carroll, O Polasek, J Viikari, M Kähönen, O Raitakari, T Lehtimäki.

Interpretation of results: C Helmer, JC Lambert, M Metzger, B Stengel, V Chouraki, J Tremblay, J Chalmers, M Woodward, P Hamet, AV Smith, A Parsa, JR O'Connell, A Tin, A Köttgen, M Li, WHL Kao, Y Li, EG Holliday, J Attia, I Guessous, CA Peralta, AC Morrison, JF Felix, C Pattaro, G Li, IH de Boer, O Devuyst, H Lin, A Isaacs, V Emilsson, AD Johnson, CS Fox, M Olden, Q Yang, EJ Atkinson, M de Andrade, ST Turner, T Zeller, J Ding, Y Liu, M Nalls, A Adeyemo, D Shriner, D Cusi, E Salvi, V Mijatovic, D Ruggiero, R Sorice, AW Dreisbach, AY Chu, DI Chasman, CA Böger, IM Heid, M Gorski, B Tayo, C Fuchsberger, H Snieder, IM Nolte, W Igl, K Susztak,

N Verweij, S Trompet, A Dehghan, B Kollerits, F Kronenberg, A Teumer, J Divers, KL Keene, MM Sale, WM Chen, GB Ehret, I Prokopenko, R Mägi, JC Denny, RJ Carroll.

Design, performance and interpretation of zebrafish experiments: M Garnaas, W Goessling.

Drafting manuscript: J Chalmers, A Tin, A Köttgen, WHL Kao, M Bochud, CA Peralta, C Pattaro, IH de Boer, CS Fox, M Garnaas, W Goessling, N Soranzo, CA Böger, IM Heid, M Gorski, S Trompet, A Dehghan, A Teumer, KL Keene, MM Sale.

Critical review of the manuscript: J Tremblay, J Chalmers, M Woodward, P Hamet, TB Harris, V Gudnason, A Parsa, AR Shuldiner, BD Mitchell, A Tin, A Köttgen, E Boerwinkle, J Coresh, M Li, WHL Kao, Y Li, H Schmidt, M Cavalieri, R Schmidt, JB Whitfield, EG Holliday, JJ Wang, J Attia, P Mitchell, I Guessous, JM Gaspoz, M Bochud, CA Peralta, AC Morrison, JF Felix, C Pattaro, DS Siscovick, IH de Boer, M Rao, R Katz, O Devuyt, TH Pers, A Isaacs, H Brenner, M Garnaas, W Goessling, BK Krämer, M Rheinberger, ST Turner, D Czamara, S Kloiber, T Zeller, BI Freedman, JM Stafford, J Ding, Y Liu, MA McEvoy, RJ Scott, SJ Hancock, JA Smith, JD Faul, SLR Kardia, AB Zonderman, M Nalls, MK Evans, FB Hu, GC Curhan, MC Cornelis, A Lupo, G Gambaro, G Malerba, M Ciullo, R Sorice, AW Dreisbach, AY Chu, DI Chasman, C Gieger, H Grallert, C Meisinger, M Waldenberger, CA Böger, HE Wichmann, IM Heid, M Gorski, T Illig, W Koenig, I Kolcic, M Boban, T Zemunik, O März, B Tayo, H Kramer, SE Rosas, C Fuchsberger, D Ruderfer, EP Bottinger, O Gottesman, RJF Loos, Y Lu, H Snieder, H Campbell, A Franke, W Lieb, IM Leach, BM Buckley, I Ford, JW Jukema, S Trompet, A Dehghan, S Sedaghat, GA Thun, M Adam, M Imboden, N Probst-Hensch, B Kollerits, B Paulweber, L Kedenko, F Kronenberg, A Teumer, K Endlich, H Völzke, KL Keene, MM Sale, WM Chen, B Ponte, D Ackermann, M Pruijm, GB Ehret, A Tönjes, I Prokopenko, M Stumvoll, P Kovacs, R Mägi, JC Denny, O Polasek, J Viikari, LP Lyytikäinen, M Kähönen, O Raitakari, T Lehtimäki.

Statistical methods and analysis: C Helmer, JC Lambert, M Metzger, V Chouraki, J Tremblay, P Hamet, AV Smith, T Aspelund, A Parsa, JR O'Connell, A Tin, A Köttgen, M Li, M Foster, WHL Kao, Y Li, H Schmidt, M Struchalin, NG Martin, RPS Middelberg, T Tanaka, E Rohtchina, EG Holliday, I Guessous, M Bochud, JF Felix, C Pattaro, G Li, R Katz, JN Hirschhorn, J Karjalainen, L Franke, TH Pers, L Yengo, N Bouatia-Naji, H Lin, T Nikopensius, T Esko, A Isaacs, A Demirkan, MF Feitosa, M Olden, MH Chen, Q Yang, SJ Hwang, M Garnaas, W Goessling, EJ Atkinson, M de Andrade, D Czamara, S Kloiber, C Müller, JM Stafford, J Ding, K Lohman, Y Liu, JA Smith, JD Faul, M Nalls, MC Cornelis, A Adeyemo, D Shriner, E Salvi, V Mijatovic, A Robino, S Ulivi, R Sorice, G Pistis, M Cocca, AY Chu, DI Chasman, LM Rose, CA Böger, IM Heid, M Gorski, ME Kleber, B Tayo, C Fuchsberger, A Saint-Pierre, D Taliun, D Ruderfer, Y Lu, IM Nolte, PJ van der Most, S Enroth, W Igl, F Murgia, L Portas, K Susztak, YA Ko, N Verweij, S Trompet, A Dehghan, S Sedaghat, GA Thun, M Adam, M Imboden, N Probst-Hensch, B Kollerits, A Teumer, J Divers, WM Chen, GB Ehret, I Prokopenko, R Mägi, CM Shaffer, RJ Carroll, C Hayward, V Vitart, LP Lyytikäinen, V Aalto.

Genotyping: JC Lambert, J Tremblay, P Hamet, E Boerwinkle, WHL Kao, H Schmidt, GW Montgomery, L Ferrucci, M Bochud, BA Oostra, CM van Duijn, K Butterbach, I Borecki, M de Andrade, T Zeller, Y Liu, RJ Scott, SLR Kardia, M Nalls, FB Hu, GC Curhan, A Adeyemo, D Shriner, D Cusi, N Soranzo, P d'Adamo, D Ruggiero, M Ciullo, R Sorice, DI Chasman, H Grallert, T Zemunik, ME Kleber, EP Bottinger, O Gottesman, RJF Loos, AF Wright, JF Wilson, A Franke, D Ellinghaus, JW Jukema, S Trompet, AG Uitterlinden, R Rivadeneira, F Kronenberg, S Coassin, M Haun, F Ernst, G Homuth, HK Kroemer, M Nauck, U Völker, H Wallaschofski, MM Sale, GB Ehret, A Tönjes, M Stumvoll, P Kovacs, CM Shaffer, JC Denny, PM Ridker, T Lehtimäki.

Bioinformatics: JC Lambert, V Chouraki, J Tremblay, P Hamet, AV Smith, T Aspelund, JR O'Connell, E Boerwinkle, Y Li, M Struchalin, GW Montgomery, RPS Middelberg, T Tanaka, C Pattaro, G Li, JN Hirschhorn, J Karjalainen, L Franke, TH Pers, L Yengo, T Esko, AD Johnson, M Olden, M Garnaas, W Goessling, D Czamara, C Müller, JA Smith, SLR Kardia, M Nalls, E Salvi, G Malerba, V Mijatovic, P d'Adamo, S Ulivi, R Sorice, C Sala, G Pistis, M Cocca, DI Chasman, H Grallert, M Waldenberger, CA Böger, IM Heid, M Gorski, ME Kleber, D Taliun, O Gottesman, S Enroth, K Susztak, YA Ko, D Ellinghaus, N Verweij, I Ford, S Trompet, F Rivadeneira, WM Chen, GB Ehret, R Mägi, CM Shaffer, JC Denny, RJ Carroll, C Hayward, LP Lyytikäinen, V Aalto.

Additional information

Supplementary Information accompanies this paper at <http://www.nature.com/naturecommunications>

Competing financial interests: J.T. and P.H. are consultants for Servier. J.C. received research grants and honoraria from Servier. K.S. obtained research support from Boehringer Ingelheim. The remaining authors declared no competing financial interests.

Reprints and permission information is available online at <http://npg.nature.com/reprintsandpermissions/>

How to cite this article: Pattaro, C. *et al.* Genetic associations at 53 loci highlight cell types and biological pathways relevant for kidney function. *Nat. Commun.* 7:10023 doi: 10.1038/ncomms10023 (2016).



This work is licensed under a Creative Commons Attribution 4.0 International License. The images or other third party material in this article are included in the article's Creative Commons license, unless indicated otherwise in the credit line; if the material is not included under the Creative Commons license, users will need to obtain permission from the license holder to reproduce the material. To view a copy of this license, visit <http://creativecommons.org/licenses/by/4.0/>

Cristian Pattaro^{1,*}, Alexander Teumer^{2,3,*}, Mathias Gorski^{4,5,*}, Audrey Y. Chu^{6,*}, Man Li^{7,*}, Vladan Mijatovic^{8,*}, Maija Garnaas⁹, Adrienne Tin⁷, Rossella Sorice¹⁰, Yong Li¹¹, Daniel Taliun¹, Matthias Olden^{4,5}, Meredith Foster¹², Qiong Yang¹³, Ming-Huei Chen^{13,14}, Tune H. Pers^{15,16}, Andrew D. Johnson¹⁷, Yi-An Ko¹⁸, Christian Fuchsberger¹, Bamidele Tayo¹⁹, Michael Nalls²⁰, Mary F. Feitosa²¹, Aaron Isaacs^{22,23}, Abbas Dehghan²⁴, Pio d'Adamo²⁵, Adebawale Adeyemo²⁶, Aida Karina Dieffenbach^{27,28}, Alan B. Zonderman²⁹, Ilja M. Nolte³⁰, Peter J. van der Most³⁰, Alan F. Wright³¹, Alan R. Shuldiner^{32,33}, Alanna C. Morrison³⁴, Albert Hofman²⁴, Albert V. Smith^{35,36}, Albert W. Dreisbach³⁷, Andre Franke³⁸, Andre G. Uitterlinden³⁹, Andres Metspalu^{40,41}, Anke Tonjes⁴², Antonio Lupo⁴³, Antonietta Robino²⁵, Åsa Johansson⁴⁴, Ayse Demirkan²², Barbara Kollerits⁴⁵, Barry I. Freedman⁴⁶, Belen Ponte⁴⁷, Ben A. Oostra⁴⁸, Bernhard Paulweber⁴⁹, Bernhard K. Krämer⁵⁰, Braxton D. Mitchell^{32,33}, Brendan M. Buckley⁵¹, Carmen A. Peralta⁵², Caroline Hayward³¹, Catherine Helmer^{53,54}, Charles N. Rotimi²⁶, Christian M. Shaffer⁵⁵, Christian Müller^{56,57}, Cinzia Sala⁵⁸, Cornelia M. van Duijn²², Aude Saint-Pierre^{1,59}, Daniel Ackermann⁴⁷, Daniel Shriner²⁶, Daniela Ruggiero¹⁰, Daniela Toniolo^{58,60}, Yingchang Lu⁶¹, Daniele Cusi⁶², Darina Czamara⁶³, David Ellinghaus³⁸, David S. Siscovick⁶⁴, Douglas Ruderfer⁶⁵, Christian Gieger⁶⁶, Harald Grallert^{67,68,69}, Elena Rohtchina⁷⁰, Elizabeth J. Atkinson⁷¹, Elizabeth G. Holliday^{72,73}, Eric Boerwinkle³⁴, Erika Salvi⁶², Erwin P. Bottinger⁶¹, Federico Murgia⁷⁴, Fernando Rivadeneira³⁹, Florian Ernst², Florian Kronenberg⁴⁵, Frank B. Hu⁷⁵, Gerjan J. Navis⁷⁶, Gary C. Curhan⁷⁷, George B. Ehret⁷⁸, Georg Homuth², Stefan Coassin⁴⁵, Gian-Andri Thun^{79,80}, Giorgio Pistis⁵⁸,

Giovanni Gambaro⁸¹, Giovanni Malerba⁸, Grant W. Montgomery⁸², Gudny Eiriksdottir³⁵, Gunnar Jacobs⁸³, Guo Li⁶⁴, H-Erich Wichmann^{84,85,86}, Harry Campbell⁸⁷, Helena Schmidt⁸⁸, Henri Wallaschofski^{89,90}, Henry Völzke^{3,90}, Hermann Brenner^{27,28}, Heyo K. Kroemer⁹¹, Holly Kramer¹⁹, Honghuang Lin⁹², I. Mateo Leach⁹³, Ian Ford⁹⁴, Idris Guessous^{95,96,97}, Igor Rudan⁸⁷, Inga Prokopenko⁹⁸, Ingrid Borecki²¹, Iris M. Heid^{4,66}, Ivana Kolcic⁹⁹, Ivana Persico⁷⁴, J. Wouter Jukema^{100,101,102,103}, James F. Wilson⁸⁷, Janine F. Felix²⁴, Jasmin Divers¹⁰⁴, Jean-Charles Lambert¹⁰⁵, Jeanette M. Stafford¹⁰⁴, Jean-Michel Gaspoz⁹⁵, Jennifer A. Smith¹⁰⁶, Jessica D. Faul¹⁰⁷, Jie Jin Wang¹⁰⁸, Jingzhong Ding¹⁰⁹, Joel N. Hirschhorn^{15,16,110}, John Attia^{71,72}, John B. Whitfield⁸², John Chalmers¹¹¹, Jorma Viikari¹¹², Josef Coresh^{7,113}, Joshua C. Denny¹¹⁴, Juha Karjalainen¹¹⁵, Jyotika K. Fernandes¹¹⁶, Karlhans Endlich¹¹⁷, Katja Butterbach²⁷, Keith L. Keene¹¹⁸, Kurt Lohman⁴⁶, Laura Portas⁷⁴, Lenore J. Launer¹¹⁹, Leo-Pekka Lyytikäinen¹²⁰, Loic Yengo^{121,122,123}, Lude Franke¹¹⁵, Luigi Ferrucci¹²⁴, Lynda M. Rose⁶, Lyudmyla Kedenko⁴⁹, Madhumathi Rao¹², Maksim Struchalin^{125,126}, Marcus E. Kleber¹²⁷, Margherita Cavalieri¹²⁸, Margot Haun⁴⁵, Marilyn C. Cornelis⁷⁵, Marina Ciullo¹⁰, Mario Pirastu⁷⁴, Mariza de Andrade⁷¹, Mark A. McEvoy¹²⁹, Mark Woodward^{7,111,112,130}, Martin Adam^{79,80}, Massimiliano Cocca⁵⁸, Matthias Nauck^{89,90}, Medea Imboden^{79,80}, Melanie Waldenberger⁶⁷, Menno Pruijm¹³¹, Marie Metzger¹³², Michael Stumvoll⁴², Michele K. Evans¹³³, Michele M. Sale¹³⁴, Mika Kähönen¹³⁵, Mladen Boban⁹⁹, Murielle Bochud¹³⁶, Myriam Rheinberger⁵, Niek Verweij⁹³, Nabila Bouatia-Naji^{137,138}, Nicholas G. Martin^{82,139}, Nick Hastie³¹, Nicole Probst-Hensch^{79,80}, Nicole Soranzo¹⁴⁰, Olivier Devuyst¹⁴¹, Olli Raitakari¹⁴², Omri Gottesman⁶¹, Oscar H. Franco²⁴, Ozren Polasek⁹⁹, Paolo Gasparini²⁵, Patricia B. Munroe^{143,144}, Paul M. Ridker¹⁴⁵, Paul Mitchell¹⁰⁸, Paul Muntner^{146,147}, Christa Meisinger⁶⁸, Johannes H. Smit¹⁴⁸, ICBP Consortium[†], AGEN Consortium[†], CARDIOGRAM[†], CHARGE-Heart Failure Group[†], ECHOGen Consortium[†], Peter Kovacs¹⁴⁹, Philipp S. Wild¹⁵⁰, Philippe Froguel^{121,122,123}, Rainer Rettig¹⁵¹, Reedik Mägi⁴⁰, Reiner Biffar¹⁵², Reinhold Schmidt¹²⁸, Rita P.S. Middelberg⁸², Robert J. Carroll¹¹⁴, Brenda W. Penninx¹⁴⁸, Rodney J. Scott¹⁵³, Ronit Katz¹⁵⁴, Sanaz Sedaghat²⁴, Sarah H. Wild⁸⁷, Sharon L.R. Kardia¹⁰⁶, Sheila Ulivi¹⁵⁵, Shih-Jen Hwang¹⁷, Stefan Enroth⁴⁴, Stefan Kloiber⁶³, Stella Trompet¹⁰⁰, Benedicte Stengel¹³², Stephen J. Hancock^{72,73}, Stephen T. Turner¹⁵⁶, Sylvia E. Rosas¹⁸, Sylvia Stracke^{105,157}, Tamara B. Harris¹¹⁹, Tanja Zeller^{56,57}, Tatijana Zemunik⁹⁹, Terho Lehtimäki¹²⁰, Thomas Illig⁶⁸, Thor Aspelund^{35,36}, Tiit Nikopensius^{40,41}, Tõnu Esko^{15,40,41}, Toshiko Tanaka¹²⁴, Ulf Gyllenstein⁴⁴, Uwe Völker^{2,90}, Valur Emilsson^{35,158}, Veronique Vitart³¹, Ville Aalto¹⁵⁹, Vilmundur Gudnason^{35,36}, Vincent Chouraki¹⁰⁵, Wei-Min Chen¹³⁴, Wilmar Igl⁴⁴, Winfried März¹⁶⁰, Wolfgang Koenig¹⁶¹, Wolfgang Lieb⁸³, Ruth J.F. Loos^{61,162}, Yongmei Liu⁴⁶, Harold Snieder³⁰, Peter P. Pramstaller^{1,163,164}, Afshin Parsa¹⁶⁵, Jeffrey R. O'Connell³², Katalin Susztak¹⁸, Pavel Hamet¹⁶⁶, Johanne Tremblay¹⁶⁶, Ian H. de Boer¹⁵⁴, Carsten A. Böger^{5,**}, Wolfram Goessling^{9,**}, Daniel I. Chasman^{6,145,**}, Anna Köttgen^{7,11,**}, W.H. Linda Kao^{7,113,**,‡} & Caroline S. Fox^{17,167,**}

¹ Center for Biomedicine, European Academy of Bozen/Bolzano (EURAC), affiliated to the University of Lübeck, Via Galvani 31, Bolzano 39100, Italy.

² Interfaculty Institute for Genetics and Functional Genomics, Ernst-Moritz-Arndt-University Greifswald, Friedrich-Loeffler-Straße 15a, Greifswald 17487, Germany. ³ Institute for Community Medicine, University of Greifswald, Walther-Rathenau-Strasse 48, Greifswald 17487, Germany. ⁴ Department of Genetic Epidemiology, Institute of Epidemiology and Preventive Medicine, University of Regensburg, Franz-Josef-Strauß-Allee 11, Regensburg 93053, Germany.

⁵ Department of Nephrology, University Hospital Regensburg, Franz-Josef-Strauß-Allee 11, Regensburg 93053, Germany. ⁶ Preventive Medicine, Brigham and Women's Hospital, 900 Commonwealth Avenue East, Boston, Massachusetts 02215, USA. ⁷ Department of Epidemiology, Johns Hopkins Bloomberg School of Public Health, 615 North Wolfe Street, Baltimore, Maryland 21205, USA. ⁸ Department of Life and Reproduction Sciences, University of Verona, Strada Le Grazie 8, Verona 37134, Italy. ⁹ Division of Genetics, Department of Medicine, Brigham and Women's Hospital, Harvard Medical School, New Research Building 77 Avenue Louis Pasteur, Room 458, Boston, Massachusetts 02115, USA. ¹⁰ Institute of Genetics and Biophysics "Adriano Buzzati-Traverso"—CNR, Via P. Castellino 111, Napoli 80131, Italy. ¹¹ Department of Internal Medicine IV, University Hospital Freiburg, Berliner Allee 29, Freiburg 79110, Germany.

¹² Division of Nephrology/Tufts Evidence Practice Center, Tufts University School of Medicine, Tufts Medical Center, Boston, Massachusetts 02111, USA.

¹³ Department of Biostatistics, Boston University School of Public Health, 715 Albany Street, Boston, Massachusetts 02118, USA. ¹⁴ Department of Neurology, Boston University School of Medicine, 72 East Concord ST B603, Boston, Massachusetts 02118, USA. ¹⁵ Division of Endocrinology and Center for Basic and Translational Obesity Research, Boston Children's Hospital, 300 Longwood Avenue, Boston, Massachusetts 02115, USA. ¹⁶ Medical and Population Genetics Program, Broad Institute of MIT and Harvard, Cambridge, Massachusetts 02142, USA. ¹⁷ NHLBI's Framingham Heart Study and the Center for Population

Studies, 73 Mt Wayte Avenue, Suite 2, Framingham, Massachusetts 01702, USA. ¹⁸ Renal Electrolyte and Hypertension Division, Perelman School of Medicine, University of Pennsylvania, 415 Curie Boulevard, 405B Clinical Research Building, Philadelphia, Pennsylvania 19104-4539, USA. ¹⁹ Department of Public Health Sciences, Loyola Medical Center, 2160 S First Avenue, Maywood, Illinois 60153, USA. ²⁰ Laboratory of Neurogenetics, Building 35—Porter Building, 1A1015, National Institute on Aging/NIH, Bethesda, Maryland 20892, USA. ²¹ Division of Statistical Genomics, Washington University School of Medicine, 4444 Forest Park Boulevard, Box 8506, St Louis, Missouri 63108, USA. ²² Genetic Epidemiology Unit, Department of Epidemiology, Erasmus University Medical Center, Dr Molewaterplein 50, PO Box 2040, Rotterdam 3000 CA, The Netherlands. ²³ Centre for Medical Systems Biology Leiden, Dr Molewaterplein 50, PO Box 2040, Rotterdam 3000 CA, The Netherlands. ²⁴ Department of Epidemiology, Erasmus University Medical Center, PO Box 2040, Rotterdam 3000 CA, The Netherlands. ²⁵ Institute for Maternal and Child Health—IRCCS “Burlo Garofolo” and University of Trieste, via dell’Istria 65/1, Trieste 34137, Italy. ²⁶ Center for Research on Genomics and Global Health, National Human Genome Research Institute, Building 12A, Room 4047, 12 South Dr, MSC 5635, Bethesda, Maryland 20892-5635, USA. ²⁷ Division of Clinical Epidemiology and Aging Research, German Cancer Research Center (DKFZ), Im Neuenheimer Feld 581, Heidelberg 69120, Germany. ²⁸ German Cancer Consortium (DKTK), Im Neuenheimer Feld 581, Heidelberg 69120, Germany. ²⁹ Laboratory of Personality and Cognition, National Institute on Aging, National Institutes of Health, NIH Biomedical Center, 251 Bayview Boulevard, Suite 100, Baltimore, Maryland 21224, USA. ³⁰ Unit of Genetic Epidemiology and Bioinformatics, Department of Epidemiology, University Medical Center Groningen, PO Box 30001, Groningen 9700 RB, The Netherlands. ³¹ MRC Human Genetics Unit, Institute of Genetics and Molecular Medicine, University of Edinburgh, Crewe Road, Edinburgh EH4 2XU, UK. ³² Department of Medicine, University of Maryland School of Medicine, 685 West Baltimore Street, Baltimore, Maryland 21201, USA. ³³ Geriatric Research and Education Clinical Center, Veterans Administration Medical Center, 10 North Greene Street, Baltimore, Maryland 21201, USA. ³⁴ Human Genetics Center, University of Texas Health Science Center at Houston, 1200 Pressler St Suite 453E, Houston, Texas 77030, USA. ³⁵ Icelandic Heart Association, Research Institute, Holtasmari 1, Kopavogur 201, Iceland. ³⁶ University of Iceland, Sæmundargötu 2, Reykjavík 101, Iceland. ³⁷ Division of Nephrology, University of Mississippi, 2500 North State Street, Jackson, Mississippi 39216, USA. ³⁸ Institute of Clinical Molecular Biology, Christian-Albrechts University of Kiel, Schittenhelmstraße 12, Kiel 24105, Germany. ³⁹ Department of Internal Medicine, Erasmus University Medical Center, PO Box 1738, Rotterdam 3000 DR, The Netherlands. ⁴⁰ Estonian Genome Center of University of Tartu (EGCUT), Riia 23B, Tartu 51010, Estonia. ⁴¹ Institute of Molecular and Cell Biology, University of Tartu and Estonian Biocenter, Riia 23, Tartu 51010, Estonia. ⁴² Department of Medicine, University of Leipzig, Liebigstraße 18, Leipzig 04103, Germany. ⁴³ Division of Nephrology, Department of Medicine, University of Verona, Piazzale Aristide Stefani 1, Verona 37126, Italy. ⁴⁴ Uppsala University, Department of Immunology, Genetics and Pathology, Biomedical Center, SciLifeLab, Uppsala University, Uppsala SE- 75108, Sweden. ⁴⁵ Innsbruck Medical University, Division of Genetic Epidemiology, Schoepfstraße 41, Innsbruck 6020, Austria. ⁴⁶ Internal Medicine Department, Wake Forest School of Medicine, Medical Center Boulevard, Winston-Salem, North Carolina 27157-1053, USA. ⁴⁷ Nephrology Division, Department of Specialties of Internal Medicine, Geneva University Hospital, 4 rue Gabrielle-Perret-Gentil, Geneve 1211, Switzerland. ⁴⁸ Department of Clinical Genetics, Erasmus University Medical Center, Dr Molewaterplein 50, PO Box 2040, Rotterdam 3000 CA, The Netherlands. ⁴⁹ First Department of Internal Medicine, Paracelsus Medical University/Salzbürger Landeskliniken, Müllner Hauptstraße 48, Salzburg 5020, Austria. ⁵⁰ University Medical Centre Mannheim, 5th Department of Medicine, University of Heidelberg, Theodor Kutzer Ufer 1-3, Mannheim 68167, Germany. ⁵¹ Department of Pharmacology and Therapeutics, University College Cork, Clinical Investigations Building, Western Rd, Cork, Ireland. ⁵² Division of Nephrology, University of California, San Francisco Medical School and San Francisco VA Medical Center, 4150 Clement Street, San Francisco, California 94121, USA. ⁵³ INSERM, ISPED, Centre INSERM U897—Epidemiologie-Biostatistique, Bordeaux F-33000, France. ⁵⁴ Université Bordeaux, ISPED, Centre INSERM U897-Epidemiologie-Biostatistique, Bordeaux F-33000, France. ⁵⁵ Vanderbilt University School of Medicine, 2215-B Garland Avenue 1224—MRB4 (Light Hall) Nashville, Tennessee 37232, USA. ⁵⁶ University Heart Center Hamburg, Clinic for general and interventional cardiology, Martinistraße 52, Hamburg 20246, Germany. ⁵⁷ German Center for Cardiovascular Research (DZHK), Partner Site Hamburg/Lübeck/Kiel, Martinistraße 52, Hamburg 20246, Germany. ⁵⁸ Division of Genetics and Cell Biology, San Raffaele Scientific Institute, Via Olgettina 58, Milano 20132, Italy. ⁵⁹ INSERM U1078, Etablissement Français du Sang, 46 rue Félix Le Dantec, CS 51819, Brest Cedex 2 29218, France. ⁶⁰ Institute of Molecular Genetics-CNR, Via Abbiategrosso 207, Pavia 27100, Italy. ⁶¹ The Charles Bronfman Institute for Personalized Medicine, Icahn School of Medicine at Mount Sinai, New York, New York 10029, USA. ⁶² Department of Health Sciences, University of Milano, Via Antonio di Rudini 8, Milano 20142, Italy. ⁶³ Max Planck Institute of Psychiatry, Kraepelinstraße 2-10, Munich 80804, Germany. ⁶⁴ Cardiovascular Health Research Unit, Departments of Epidemiology and Medicine, University of Washington, 1730 Minor Ave, Suite 1360, Seattle, Washington 98101, USA. ⁶⁵ Division of Psychiatric Genomics, Department of Psychiatry, Icahn School of Medicine at Mount Sinai, New York, New York 10029, USA. ⁶⁶ Institute of Genetic Epidemiology, Helmholtz Zentrum München, German Research Center for Environmental Health, Ingolstädter Landstraße 1, 85764 Neuherberg, Germany. ⁶⁷ Research Unit of Molecular Epidemiology, Helmholtz Zentrum München, German Research Center for Environmental Health, Ingolstädter Landstraße 1, 85764 Neuherberg, Germany. ⁶⁸ Institute of Epidemiology II, Helmholtz Zentrum München, German Research Center for Environmental Health, 85764 Neuherberg, Germany. ⁶⁹ German Center for Diabetes Research (DZD), Ingolstädter Landstraße 1, Neuherberg 85764, Germany. ⁷⁰ Westmead Millennium Institute, Centre for Vision Research, University of Sydney, C24 Westmead Hospital, New South Wales 2145, Australia. ⁷¹ Division of Biomedical Statistics and Informatics, Mayo Clinic, 200 First Street SW, Rochester, Minnesota 55905, USA. ⁷² Centre for Clinical Epidemiology and Biostatistics, School of Medicine and Public Health, University of Newcastle, HMRI Building1, Kookaburra Circuit, New Lambton New South Wales 2305, Australia. ⁷³ Clinical Research Design, Information Technology and Statistical Support, Hunter Medical Research Institute, Newcastle, 1 Kookaburra Circuit, New Lambton Heights, New South Wales 2305, Australia. ⁷⁴ Institute of Population Genetics—CNR, Traversa La Crucca 3, 07040 Reg. Balduina, Li Punti, Sassari, Italy. ⁷⁵ Department of Nutrition, Harvard School of Public Health, 665 Huntington Avenue, Building 2, Boston, Massachusetts 02115, USA. ⁷⁶ Department of Internal Medicine, University Medical Center Groningen, University of Groningen, Hanzeplein 1, Groningen 9713 GZ, The Netherlands. ⁷⁷ Brigham and Women’s Hospital and Channing Laboratory, Harvard Medical School, 181 Longwood Avenue, Boston, Massachusetts 02115, USA. ⁷⁸ Cardiology, Department of Specialties of Internal Medicine, Geneva University Hospitals, Rue Gabrielle-Perret-Gentil 4, Geneva 1205, Switzerland. ⁷⁹ Swiss Tropical and Public Health Institute, PO Box 4002, Basel, Switzerland. ⁸⁰ University of Basel, Petersplatz 1, Basel 4003, Switzerland. ⁸¹ Division of Nephrology, Department of Internal Medicine and Medical Specialties, Columbus-Gemelli University Hospital, Catholic University, Via Moscatti 31, Rome 00168, Italy. ⁸² Genetic Epidemiology, Queensland Institute of Medical Research, QIMR, PO Royal Brisbane Hospital, Queensland 4029, Australia. ⁸³ Institute of Epidemiology and Biobank popgen, Christian-Albrechts University, Niemannsweg 11, Kiel 24105, Germany. ⁸⁴ Institute of Epidemiology I, Helmholtz Zentrum München, German Research Center for Environmental Health, Ingolstädter Landstraße 1, Neuherberg 85764, Germany. ⁸⁵ Institute of Medical Informatics, Biometry and Epidemiology, Ludwig-Maximilians-Universität, Ingolstädter Landstrasse 1, 85764 Neuherberg, Germany. ⁸⁶ Klinikum Grosshadern, Ingolstädter Landstraße 1, Neuherberg 85764, Germany. ⁸⁷ Centre for Population Health Sciences, University of Edinburgh, Teviot Place, Edinburgh, EH8 9AG Scotland, UK. ⁸⁸ Austrian Stroke Prevention Study, Institute of Molecular Biology and Biochemistry, Department of Neurology, Medical University Graz, Harrachgasse 21, Graz 8010, Austria. ⁸⁹ Institute of Clinical Chemistry and Laboratory Medicine, University Medicine Greifswald, Ferdinand-Sauerbruch-Straße, Greifswald 17475, Germany. ⁹⁰ German Center for Cardiovascular Research (DZHK), Partner site Greifswald, Ferdinand-Sauerbruch-Straße, Greifswald 17475, Germany. ⁹¹ Institute of Pharmacology, University of Greifswald, Friedrich-Loeffler-Straße 23d, Greifswald 17487, Germany. ⁹² Boston University School of Medicine, 72 East Concord Street, B-616, Boston, Massachusetts 02118, USA. ⁹³ Department of Cardiology, University Medical Center Groningen, University of Groningen, PO Box 30.001, Groningen 9700 RB, The Netherlands. ⁹⁴ Robertson Centre for Biostatistics, University of Glasgow, R1122B Level 11, Robertson Centre, Boyd Orr Building, Glasgow G12 8QQ, UK. ⁹⁵ Division of Primary Care Medicine, Department of Community

Medicine, Primary Care and Emergency Medicine, Geneva University Hospitals, Faculty of Medicine, University of Geneva, Geneva 1211, Switzerland. ⁹⁶Community Prevention Unit, University Institute of Social and Preventive Medicine, Lausanne University Hospital, Route de la Corniche 10, Lausanne 1010, Switzerland. ⁹⁷Department of Epidemiology, Rollins School of Public Health, Emory University, 1518 Clifton Road, NE, Atlanta, Georgia 30322, USA. ⁹⁸Department of Genomics of Common Disease, School of Public Health, Imperial College London, London W12 0NN, UK. ⁹⁹Croatian Centre for Global Health, University of Split Medical School, Šoltanska 2, Split 21000, Croatia. ¹⁰⁰Department of Cardiology, Leiden University Medical Center, PO Box 9600, Leiden 2300 RC, The Netherlands. ¹⁰¹Interuniversity Cardiology Institute of the Netherlands (ICIN), Moreelsepark 1, Utrecht 3511 EP, The Netherlands. ¹⁰²Eindhoven Laboratory for Experimental Vascular Medicine, Albinusdreef 2, Leiden 2333 ZA, The Netherlands. ¹⁰³Durrer Center for Cardiogenetic Research, Meibergdreef 9, Amsterdam 1105 AZ, The Netherlands. ¹⁰⁴Division of Public Health Sciences, Department of Biostatistical Sciences, Wake Forest University Health Sciences, 2326 Medical Center Boulevard, Winston-Salem, North Carolina 27157-1063, USA. ¹⁰⁵INSERM U744, Institut Pasteur de Lille, 1 rue du Pr. Calmette, Lille Cédex 59019, France. ¹⁰⁶Department of Epidemiology, School of Public Health, University of Michigan, 1415 Washington Heights, Ann Arbor, Michigan 48109-2029, USA. ¹⁰⁷Survey Research Center, Institute for Social Research, University of Michigan, 426 Thompson Street, #3456, Ann Arbor, Michigan 48104, USA. ¹⁰⁸Centre for Vision Research, Westmead Millennium Institute, University of Sydney, C24 Westmead Hospital, New South Wales 2145, Australia. ¹⁰⁹Wake Forest School of Medicine, Department of Internal Medicine/Geriatrics, Medical center Boulevard, Winston-Salem, North Carolina 27157, USA. ¹¹⁰Department of Genetics, Harvard Medical School, 77 Avenue Louis Pasteur, NRB 0330, Boston, Massachusetts 02115, USA. ¹¹¹University of Sydney, The George Institute for Global Health, Level 10, King George V Building, 83-117 Missenden Road, Camperdown, New South Wales 2050, Australia. ¹¹²Department of Medicine, University of Turku, Turku University Hospital, PO Box 52, Turku 20521, Finland. ¹¹³Welch Center for Prevention, Epidemiology and Clinical Research, 2024 East Monument St, Suite 2-600, Baltimore, Maryland 21287, USA. ¹¹⁴Vanderbilt University School of Medicine, 448 Eskind Biomedical Library, 2209 Garland Avenue, Nashville, Tennessee 37212, USA. ¹¹⁵Department of Genetics, University of Groningen, University Medical Centre Groningen, PO Box 72, Groningen 9700 AB, The Netherlands. ¹¹⁶Division of Endocrinology, Medical University of South Carolina, 171 Ashley Avenue, Charleston, South Carolina 29425, USA. ¹¹⁷Institute of Anatomy and Cell Biology, University of Greifswald, Friedrich-Loeffler-Straße 23c, Greifswald 17487, Germany. ¹¹⁸Center for Health Disparities, Department of Biology, East Carolina University, 1001 East 10th Street, N209 Howell Science Complex Mailstop 551, Greenville, North Carolina 27858, USA. ¹¹⁹Intramural Research Program, Laboratory of Epidemiology, Demography, and Biometry, National Institute on Aging, Gateway Building, 3C309, 7201 Winkscon Avenue, Bethesda, Maryland 20892-9205, USA. ¹²⁰Department of Clinical Chemistry, Fimlab Laboratories, University of Tampere, School of Medicine, Tampere 33520, Finland. ¹²¹CNRS UMR 8199, 1 Rue du Professeur Calmette, Lille 59000, France. ¹²²Lille Pasteur Institute, 1 Rue du Professeur Calmette, Lille 59000, France. ¹²³Lille II University, 42 Rue Paul Duez, Lille 59000, France. ¹²⁴Clinical Research Branch, National Institute on Aging, 251 Bayview Blvd, Baltimore, Maryland 21250, USA. ¹²⁵Department of Epidemiology and Biostatistics, Erasmus University Medical Center, Dr Molewaterplein, Rotterdam 50-603015 GE, The Netherlands. ¹²⁶Department of Forensic Molecular Biology, Erasmus University Medical Center, Dr Molewaterplein, Rotterdam 50-603015 GE, The Netherlands. ¹²⁷Medical Clinic V, Medical Faculty Mannheim, Heidelberg University, Theodor-Kutzer-Ufer 1-3, Mannheim 68167, Germany. ¹²⁸Austrian Stroke Prevention Study, Department of Neurology, Division of Special Neurology, Medical University Graz, Auenbruggerplatz 22, Graz 8036, Austria. ¹²⁹Centre for Clinical Epidemiology and Biostatistics, University of Newcastle, Hunter Medical Research Institute, John Hunter Hospital, Locked Bag 1, HRMC, New South Wales 2310, Australia. ¹³⁰The George Institute for Global Health, Nuffield Department of Population Health, University of Oxford, Old Road Campus, Roosevelt Drive, Oxford OX3 7LF, UK. ¹³¹Service of Nephrology, Lausanne University Hospital, Rue du Bugnon 17, Lausanne 1005, Switzerland. ¹³²Inserm UMRS 1018, CESP Team 10, Université Paris-Sud, 16 avenue Paul Vaillant Couturier, Villejuif 94807, France. ¹³³Health Disparities Research Section, Clinical Research Branch, National Institute on Aging, National Institutes of Health, NIH Biomedical Center, 251 Bayview Boulevard, Suite 100, Baltimore, Maryland 21224, USA. ¹³⁴Center for Public Health Genomics, Department of Medicine (Cardiovascular Medicine), University of Virginia, PO Box 800717, Charlottesville, Virginia 22908, USA. ¹³⁵Department of Clinical Physiology, Tampere University Hospital, University of Tampere, School of Medicine, Tampere 33521, Finland. ¹³⁶University Institute of Social and Preventive Medicine, Centre Hospitalier Universitaire Vaudois, University of Lausanne, Route de la Corniche 2, Epalinges CH-1066, Switzerland. ¹³⁷INSERM UMR970, Paris Cardiovascular Research Center (PARCC), 56 rue Leblanc, Paris F-75015, France. ¹³⁸Paris Descartes University, Faculty of medicine, Paris Cité Sorbonne, 12 Rue de l'école de Médecine, Paris F-75006, France. ¹³⁹Brigham and Women's Hospital, Harvard Medical School, 77 Avenue Louis Pasteur, Boston, Massachusetts 02115, USA. ¹⁴⁰Wellcome Trust Sanger Institute, Hinxton CB10 1HH, UK. ¹⁴¹University of Zurich, Institute of Physiology, Mechanisms of Inherited Kidney Disorders Group, Winterthurerstrasse 190, Zürich 8057, Switzerland. ¹⁴²Research Centre of Applied and Preventive Cardiovascular Medicine, University of Turku, Turku University Hospital, Department of Clinical Physiology, PO Box 52, Turku 20521, Finland. ¹⁴³Department Clinical Pharmacology, William Harvey Research Institute, Queen Mary University of London, London EC1M 6BQ, UK. ¹⁴⁴NIHR Barts Cardiovascular Biomedical Research Unit, Queen Mary University of London, London EC1M 6BQ, UK. ¹⁴⁵Harvard Medical School, 900 Commonwealth Avenue East, Boston, Massachusetts 02115, USA. ¹⁴⁶University of Alabama at Birmingham, Department of Medicine, 1530 3rd Avenue, South Birmingham, Alabama 35294-0022, USA. ¹⁴⁷University of Alabama at Birmingham, Department of Epidemiology, 1530 3rd Avenue, South Birmingham, Alabama 35294-0022, USA. ¹⁴⁸Department of Psychiatry and EMGO + Institute, VU University Medical Center, A.J. Ernststraat 1187, Amsterdam 1081 HL, The Netherlands. ¹⁴⁹IFB AdiposityDiseases, University of Leipzig, Liebigstraße 21, Leipzig 04103, Germany. ¹⁵⁰Medical University Center Mainz, Langenbeckstraße 1, Mainz 55131, Germany. ¹⁵¹Institute of Physiology, University of Greifswald, Greifswald 17487, Germany. ¹⁵²Clinic for Prosthodontic Dentistry, Gerostomatology and Material Science, University of Greifswald, Rotgerberstraße 8, Greifswald 17475, Germany. ¹⁵³School of Biomedical Sciences and Pharmacy, University of Newcastle, Hunter Medical Research Institute, John Hunter Hospital, Locked Bag 1, HRMC, New South Wales 2310, Australia. ¹⁵⁴Kidney Research Institute, University of Washington, Box 359606, 325 9th Avenue, Seattle, Washington 98104, USA. ¹⁵⁵Institute for Maternal and Child Health—IRCCS "Burlo Garofolo", Via dell'Istria 65, Trieste 34137, Italy. ¹⁵⁶Department of Internal Medicine, Division of Nephrology and Hypertension, Mayo Clinic, 200 First Street SW, Rochester, Minnesota 55905, USA. ¹⁵⁷Clinic for Internal Medicine A, University of Greifswald, Friedrich-Loeffler-Straße 23a, Greifswald 17475, Germany. ¹⁵⁸Faculty of Pharmaceutical Sciences, University of Iceland, Sæmundargata 2, Reykjavík 101, Iceland. ¹⁵⁹Department of Clinical Physiology, Turku University Hospital, Research Centre of Applied and Preventive Cardiovascular Medicine, University of Turku, PO Box 52, Turku 20521, Finland. ¹⁶⁰Synlab Academy, Synlab Services GmbH, Oberer Eselsberg 45, Ulm 89081, Germany. ¹⁶¹Department of Internal Medicine II—Cardiology, University of Ulm Medical Centre, Albert-Einstein-Allee 23, Ulm 89081, Germany. ¹⁶²The Mindich Child Health and Development Institute, Ichan School of Medicine at Mount Sinai, New York, New York 10029, USA. ¹⁶³Department of Neurology, General Central Hospital, Via Lorenz Bohler 5, Bolzano 39100, Italy. ¹⁶⁴Department of Neurology, University of Lübeck, Ratzeburger Allee 160, Lübeck 23538, Germany. ¹⁶⁵University of Maryland Medical School, Division of Nephrology, 685 W. Baltimore Street, MSTF 314, Baltimore, Maryland 21201, USA. ¹⁶⁶CRCHUM, University of Montreal, CHUM Research Center, Technopôle Angus, 900 Saint-Denis, Montreal, Québec, Canada H2X 0A9. ¹⁶⁷Division of Endocrinology, Brigham and Women's Hospital, Harvard Medical School, 221 Longwood Avenue, Boston, Massachusetts 02115, USA. * These authors contributed equally to this work. ** These authors jointly supervised this work. †A full list of consortium members appears at the end of the paper.

‡Deceased.

ICBP Consortium

Goncalo R. Abecasis¹⁶⁸, Linda S. Adair¹⁶⁹, Myriam Alexander¹⁷⁰, David Altshuler^{171,172}, Najaf Amin²⁴, Dan E. Arking¹⁷³, Pankaj Arora¹⁷⁴, Yurii Aulchenko²⁴, Stephan J.L. Bakker⁷⁶, Stefania Bandinelli¹⁷⁵, Ines Barroso¹⁴⁰, Jacques S. Beckmann¹⁷⁶, John P. Beilby¹⁷⁷, Richard N. Bergman¹⁷⁸, Sven Bergmann¹⁷⁶, Joshua C. Bis¹⁷⁹, Michael Boehnke¹⁶⁸, Lori L. Bonnycastle¹⁸⁰, Stefan R. Bornstein¹⁸¹, Michiel L. Bots¹⁸², Jennifer L. Bragg-Gresham¹⁶⁸, Stefan-Martin Brand¹⁸³, Eva Brand¹⁸⁴, Peter S. Braund¹⁸⁵, Morris J. Brown¹⁸⁶, Paul R. Burton¹⁸⁷, Juan P. Casas¹⁸⁸, Mark J. Caulfield¹⁸⁹, Aravinda Chakravarti¹⁷³, John C. Chambers¹⁹⁰, Giriraj R. Chandak¹⁹¹, Yen-Pei C. Chang¹⁹², Fadi J. Charchar¹⁹³, Nish Chaturvedi¹⁹⁴, Yoon Shin Cho¹⁹⁵, Robert Clarke¹⁹⁶, Francis S. Collins¹⁸⁰, Rory Collins¹⁹⁶, John M. Connell¹⁹⁷, Jackie A. Cooper¹⁹⁸, Matthew N. Cooper¹⁹⁹, Richard S. Cooper²⁰⁰, Anna Maria Corsi²⁰¹, Marcus Dörr²⁰², Santosh Dahgam²⁰³, John Danesh¹⁷⁰, George Davey Smith²⁰⁴, Ian N.M. Day²⁰⁴, Panos Deloukas¹⁴⁰, Matthew Denniff¹⁸⁵, Anna F. Dominiczak²⁰⁵, Yanbin Dong²⁰⁶, Ayo Doumatey²⁶, Paul Elliott¹⁹⁰, Roberto Elosua²⁰⁷, Jeanette Erdmann²⁰⁸, Susana Eyheramendy²⁰⁹, Martin Farrall²¹⁰, Cristiano Fava²¹¹, Terrence Forrester²¹², F. Gerald R. Fowkes⁸⁷, Ervin R. Fox²¹³, Timothy M. Frayling²¹⁴, Pilar Galan²¹⁵, Santhi K. Ganesh²¹⁶, Melissa Garcia²¹⁷, Tom R. Gaunt²⁰⁴, Nicole L. Glazer¹⁷⁹, Min Jin Go¹⁹⁵, Anuj Goel²¹⁰, Jürgen Grässler¹⁸¹, Diederick E. Grobbee¹⁸², Leif Groop²¹⁸, Simonetta Guarrera²¹⁹, Xiuqing Guo²²⁰, David Hadley²²¹, Anders Hamsten²²², Bok-Ghee Han¹⁹⁵, Rebecca Hardy²²³, Anna-Liisa Hartikainen²²⁴, Simon Heath²²⁵, Susan R. Heckbert²²⁶, Bo Hedblad²¹¹, Serge Hercberg²¹⁵, Dena Hernandez²⁰, Andrew A. Hicks¹, Gina Hilton¹⁷³, Aroon D. Hingorani²²⁷, Judith A. Hoffman Bolton⁷, Jemma C. Hopewell¹⁹⁶, Philip Howard²²⁸, Steve E. Humphries¹⁹⁸, Steven C. Hunt²²⁹, Kristian Hveem²³⁰, M. Arfan Ikram²⁴, Muhammad Islam^{231,232}, Naoharu Iwai^{233,234}, Marjo-Riitta Jarvelin¹⁹⁰, Anne U. Jackson¹⁶⁸, Tazeen H. Jafar^{231,232}, Charles S. Janipalli¹⁹¹, Toby Johnson¹⁸⁹, Sekar Kathiresan²³⁵, Kay-Tee Khaw¹⁷⁰, Hyung-Lae Kim¹⁹⁵, Sanjay Kinra²³⁶, Yoshikuni Kita²³⁷, Mika Kivimäki²²⁷, Jaspal S. Kooner²³⁸, M.J. Kranthi Kumar¹⁹¹, Diana Kuh²²³, Smita R. Kulkarni²³⁹, Meena Kumari²⁴⁰, Johanna Kuusisto²⁴¹, Tatiana Kuznetsova²⁴², Markku Laakso²⁴¹, Maris Laan²⁴³, Jaana Laitinen²⁴⁴, Edward G. Lakatta²⁴⁵, Carl D. Langefeld²⁴⁶, Martin G. Larson²⁴⁷, Mark Lathrop²²⁵, Debbie A. Lawlor²⁰⁴, Robert W. Lawrence¹⁹⁹, Jong-Young Lee¹⁹⁵, Nanette R. Lee²⁴⁸, Daniel Levy²⁴⁷, Yali Li²⁴⁹, Will T. Longstreth²⁵⁰, Jian'an Luan²⁵¹, Gavin Lucas²⁰⁷, Barbara Ludwig¹⁸¹, Massimo Mangino²⁵², K. Radha Mani¹⁹¹, Michael G. Marmot²²⁷, Francesco U.S. Mattace-Raso²⁴, Giuseppe Matullo²⁵³, Wendy L. McArdle²⁵⁴, Colin A. McKenzie²¹², Thomas Meitinger²⁵⁵, Olle Melander²¹¹, Pierre Meneton²⁵⁶, James F. Meschia²⁵⁷, Tetsuro Miki^{258,259}, Yuri Milaneschi¹²⁴, Karen L. Mohlke²⁶⁰, Vincent Moser²⁶¹, Mario A. Morken¹⁸⁰, Richard W. Morris²⁶², Thomas H. Mosley²⁶³, Samer Najjar²⁶⁴, Narisu Narisu¹⁸⁰, Christopher Newton-Cheh¹⁷⁴, Khanh-Dung Hoang Nguyen¹⁷³, Peter Nilsson²¹¹, Fredrik Nyberg²⁰³, Christopher J. O'Donnell²⁴⁷, Toshio Ogihara²⁶⁵, Takayoshi Ohkubo²⁶⁶, Tomonori Okamura^{233,234}, Rick Twee-Hee Ong²⁶⁷, Halit Ongen²¹⁰, N. Charlotte Onland-Moret¹⁸², Paul F. O'Reilly¹⁹⁰, Elin Org²⁴³, Marco Orru²⁶⁸, Walter Palmas²⁶⁹, Jutta Palmen¹⁹⁸, Lyle J. Palmer¹⁹⁹, Nicholette D. Palmer²⁴⁶, Alex N. Parker²⁷⁰, John F. Peden²¹⁰, Leena Peltonen¹⁴⁰, Markus Perola²⁷¹, Vasyl Pihur¹⁷³, Carl G.P. Platou²³⁰, Andrew Plump²⁷², Dorairajan Prabhakaran²⁷³, Bruce M. Psaty¹⁷⁹, Leslie J. Raffel²²⁰, Dabeeru C. Rao²⁷⁴, Asif Rasheed²⁷⁵, Fulvio Ricceri²⁵³, Kenneth M. Rice²⁷⁶, Annika Rosengren²⁷⁷, Jerome I. Rotter²²⁰, Megan E. Rudock²⁷⁸, Siim Söber²⁴³, Tunde Salako²⁷⁹, Danish Saleheen²⁷⁵, Veikko Salomaa²⁷¹, Niles J. Samani¹⁸⁵, Steven M. Schwartz²²⁶, Peter E.H. Schwarz²⁸⁰, Laura J. Scott¹⁶⁸, James Scott²³⁸, Angelo Scuteri²⁴⁵, Joban S. Sehmi²³⁸, Mark Seielstad²⁸¹, Sudha Seshadri¹⁴, Pankaj Sharma²⁸², Sue Shaw-Hawkins¹⁸⁹, Gang Shi²⁷⁴, Nick R.G. Shrine¹⁸⁷, Eric J.G. Sijbrands²⁴, Xueling Sim²⁸³, Andrew Singleton²⁰, Marketa Sjögren²¹¹,

Nicholas L. Smith²²⁶, Maria Soler Artigas¹⁸⁷, Tim D. Spector²⁵², Jan A. Staessen²⁴⁴, Alena Stancakova²⁴¹, Nanette I. Steinle¹⁹², David P. Strachan²²¹, Heather M. Stringham¹⁶⁸, Yan V. Sun¹⁰⁶, Amy J. Swift¹⁸⁰, Yasuharu Tabara^{258,259}, E-Shyong Tai²⁸⁴, Philippa J. Talmud¹⁹⁸, Andrew Taylor²⁴⁰, Janos Terzic²⁸⁵, Dag S. Thelle²⁸⁶, Martin D. Tobin¹⁸⁹, Maciej Tomaszewski¹⁸⁵, Vikal Tripathy²⁷³, Jaakko Tuomilehto²⁸⁷, Ioanna Tzoulaki¹⁹⁰, Manuela Uda²⁶⁸, Hirotugu Ueshima²⁸⁸, Cuno S.P.M. Uiterwaal¹⁸⁴, Satoshi Umemura²⁸⁹, Pim van der Harst⁹³, Yvonne T. van der Schouw¹⁸², Wiek H. van Gilst⁹³, Erkki Vartiainen²⁷¹, Ramachandran S. Vasan²⁴⁷, Gudrun Veldre²⁴³, Germaine C. Verwoert²⁴, Margus Viigimaa²⁹⁰, D.G. Vinay¹⁹¹, Paolo Vineis²⁹¹, Benjamin F. Voight²³⁵, Peter Vollenweider²⁹², Lynne E. Wagenknecht²⁴⁶, Louise V. Wain¹⁸⁷, Xiaoling Wang²⁰⁶, Thomas J. Wang²⁴⁷, Nicholas J. Wareham²⁵¹, Hugh Watkins²¹⁰, Alan B. Weder²¹⁶, Peter H. Whincup²²¹, Kerri L. Wiggins¹⁷⁹, Jacqueline C.M. Witteman²⁴, Andrew Wong²²³, Ying Wu²⁵⁹, Chittaranjan S. Yajnik²³⁹, Jie Yao²²⁰, J.H. Young²⁹³, Diana Zelenika²²⁵, Guangju Zhai²⁵², Weihua Zhang¹⁹⁰, Feng Zhang²⁵², Jing Hua Zhao²⁵¹, Haidong Zhu²⁰⁶, Xiaofeng Zhu²⁴⁹, Paavo Zitting²⁹⁴, Ewa Zukowska-Szczechowska²⁹⁵

¹⁶⁸Center for Statistical Genetics, Department of Biostatistics, University of Michigan, School of Public Health, Ann Arbor, Michigan 48103, USA.

¹⁶⁹Department of Nutrition, University of North Carolina, Chapel Hill, North Carolina 27599, USA. ¹⁷⁰Department of Public Health and Primary Care, University of Cambridge, Cambridge, CB1 8RN, UK. ¹⁷¹Department of Medicine, Harvard Medical School, Boston, Massachusetts 02115, USA. ¹⁷²Department of Genetics, Harvard Medical School, Boston, Massachusetts 02115, USA. ¹⁷³Center for Complex Disease Genomics, McKusick-Nathans Institute of Genetic Medicine, Johns Hopkins University School of Medicine, Baltimore, Maryland 21205, USA. ¹⁷⁴Center for Human Genetic Research, Cardiovascular Research Center, Massachusetts General Hospital, Boston, Massachusetts 02114, USA. ¹⁷⁵Geriatric Rehabilitation Unit, Azienda Sanitaria Firenze (ASF), 50125 Florence, Italy. ¹⁷⁶Département de Génétique Médicale, Université de Lausanne, Lausanne 1015, Switzerland. ¹⁷⁷Pathology and Laboratory Medicine, University of Western Australia, 6009 Crawley, Western Australia, Australia. ¹⁷⁸Department of Physiology and Biophysics, Keck School of Medicine, University of Southern California, Los Angeles, California 90033, USA. ¹⁷⁹Cardiovascular Health Research Unit, Department of Medicine, University of Washington, Seattle 98195, Washington, USA. ¹⁸⁰National Human Genome Research Institute, National Institutes of Health, Bethesda, Maryland 20892, USA. ¹⁸¹Department of Medicine III, Medical Faculty Carl Gustav Carus at the Technical University of Dresden, Dresden 01307, Germany. ¹⁸²Julius Center for Health Sciences and Primary Care, University Medical Center Utrecht, Heidelberglaan 100, Utrecht 3508 GA, The Netherlands. ¹⁸³Leibniz-Institute for Arteriosclerosis Research, Department of Molecular Genetics of Cardiovascular Disease, University of Münster, Münster, Germany. ¹⁸⁴University Hospital Münster, Internal Medicine D, Münster, Germany. ¹⁸⁵Department of Cardiovascular Sciences, University of Leicester, Glenfield Hospital, Leicester LE3 9QP, UK. ¹⁸⁶Clinical Pharmacology Unit, University of Cambridge, Addenbrookes Hospital, Hills Road, Cambridge CB2 2QQ, UK. ¹⁸⁷Department of Health Sciences, University of Leicester, University Rd, Leicester LE1 7RH, UK. ¹⁸⁸Faculty of Epidemiology and Population Health, London School of Hygiene and Tropical Medicine, London WC1E 7HT, UK. ¹⁸⁹Clinical Pharmacology and The Genome Centre, William Harvey Research Institute, Barts and The London School of Medicine and Dentistry, Queen Mary University of London, London EC1M 6BQ, UK. ¹⁹⁰Department of Epidemiology and Biostatistics, School of Public Health, Imperial College London, Norfolk Place, London W2 1PG, UK. ¹⁹¹Centre for Cellular and Molecular Biology (CCMB), Council of Scientific and Industrial Research (CSIR), Uppal Road, Hyderabad 500 007, India. ¹⁹²University of Maryland, School of Medicine, Baltimore, Maryland 21201, USA. ¹⁹³School of Science and Engineering, University of Ballarat, Ballarat 3353, Australia. ¹⁹⁴International Centre for Circulatory Health, National Heart & Lung Institute, Imperial College, London, UK. ¹⁹⁵Center for Genome Science, National Institute of Health, Seoul, Korea. ¹⁹⁶Clinical Trial Service Unit and Epidemiological Studies Unit, University of Oxford, Oxford OX3 7LF, UK. ¹⁹⁷University of Dundee, Ninewells Hospital & Medical School, Dundee DD1 9SY, UK. ¹⁹⁸Centre for Cardiovascular Genetics, University College London, London WC1E 6JF, UK. ¹⁹⁹Centre for Genetic Epidemiology and Biostatistics, University of Western Australia, Crawley, Western Australia, Australia. ²⁰⁰Department of Preventive Medicine and Epidemiology, Loyola University Medical School, Maywood, Illinois, USA. ²⁰¹Tuscany Regional Health Agency, Florence, Italy. ²⁰²Department of Internal Medicine B, Ernst-Moritz-Arndt-University Greifswald, Greifswald 17487, Germany. ²⁰³Occupational and Environmental Medicine, Department of Public Health and Community Medicine, Institute of Medicine, Sahlgrenska Academy, University of Gothenburg, Gothenburg 40530, Sweden. ²⁰⁴MRC Centre for Causal Analyses in Translational Epidemiology, School of Social & Community Medicine, University of Bristol, Bristol BS8 2BN, UK. ²⁰⁵BHF Glasgow Cardiovascular Research Centre, University of Glasgow, 126 University Place, Glasgow G12 8TA, UK. ²⁰⁶Georgia Prevention Institute, Department of Pediatrics, Medical College of Georgia, 30912 Augusta, Georgia, USA. ²⁰⁷Cardiovascular Epidemiology and Genetics, Institut Municipal d'Investigació Mèdica, Barcelona Biomedical Research Park, 88 Doctor Aiguader, Barcelona 08003, Spain. ²⁰⁸Medizinische Klinik II, Universität zu Lübeck, 23562 Lübeck, Germany. ²⁰⁹Department of Statistics, Pontificia Universidad Católica de Chile, Vicuña Mackenna, Santiago 4860, Chile. ²¹⁰Department of Cardiovascular Medicine, The Wellcome Trust Centre for Human Genetics, University of Oxford, Oxford OX3 7BN, UK. ²¹¹Department of Clinical Sciences, Lund University, SE-205 02 Malmö, Sweden. ²¹²Tropical Medicine Research Institute, University of the West Indies, Mona, Kingston, Jamaica. ²¹³Department of Medicine, University of Mississippi Medical Center, 2500 North State St, Jackson, Mississippi 39216, USA. ²¹⁴Genetics of Complex Traits, Peninsula Medical School, University of Exeter, EX1 2LU Exeter, UK. ²¹⁵U557 Institut National de la Santé et de la Recherche Médicale, U1125 Institut National de la Recherche Agronomique, Université Paris 13, F-93017 Bobigny, France. ²¹⁶Department of Internal Medicine, Division of Cardiovascular Medicine, University of Michigan Medical Center, Ann Arbor 48108, Michigan, USA. ²¹⁷Laboratory of Epidemiology, Demography, Biometry, National Institute on Aging, National Institutes of Health, Bethesda, Maryland 20892, USA. ²¹⁸Department of Clinical Sciences, Diabetes and Endocrinology Research Unit, University Hospital, SE-205 02 Malmö, Sweden. ²¹⁹Human Genetics Foundation (HUGEF), Via Lagrange 35, Torino 10123, Italy. ²²⁰Medical Genetics Institute, Cedars-Sinai Medical Center, Los Angeles, 90048 California, USA. ²²¹Division of Community Health Sciences, St George's University of London, London SW17 0RE, UK. ²²²Atherosclerosis Research Unit, Department of Medicine, Karolinska Institute, SE-171 77 Stockholm, Sweden. ²²³MRC Unit for Lifelong Health & Ageing, London WC1B 5JU, UK. ²²⁴Institute of Clinical Medicine/Obstetrics and Gynecology, University of Oulu, 90014 Oulu, Finland. ²²⁵Centre National de Génotypage, Commissariat de L'Energie Atomique, Institut de Génétique, 91000 Evry, France. ²²⁶Department of Epidemiology, University of Washington, Seattle, Washington 98195, USA. ²²⁷Epidemiology Public Health, UCL, London WC1E 6BT, UK. ²²⁸William Harvey Research Institute, Barts and The London School of Medicine and Dentistry, Queen Mary University of London, London EC1M 6BQ, UK. ²²⁹Cardiovascular Genetics, University of Utah, School of Medicine, Salt Lake City, 84132 Utah, USA. ²³⁰HUNT Research Centre, Department of Public Health and General Practice, Norwegian University of Science and Technology, Levanger 7600, Norway. ²³¹Department of Community

Health Sciences, Aga Khan University, 74800 Karachi, Pakistan. ²³²Department of Medicine, Aga Khan University, 74800 Karachi, Pakistan. ²³³Department of Genomic Medicine, National Cerebral and Cardiovascular Research Center, Suita 565-8565, Japan. ²³⁴Department of Preventive Cardiology, National Cerebral and Cardiovascular Research Center, Suita 565-8565, Japan. ²³⁵Medical Population Genetics, Broad Institute of Harvard and MIT, 5 Cambridge Center, Cambridge, Massachusetts 02142, USA. ²³⁶Division of Non-communicable disease Epidemiology, The London School of Hygiene and Tropical Medicine London, Keppel Street, London WC1E 7HT, UK. ²³⁷Department of Health Science, Shiga University of Medical Science, Otsu 520-2192, Japan. ²³⁸National Heart and Lung Institute, Imperial College London, London W12 0HS, UK. ²³⁹Diabetes Unit, KEM Hospital and Research Centre, Rasta Peth, Pune, 411011 Maharashtra, India. ²⁴⁰Genetic Epidemiology Group, Epidemiology and Public Health, UCL, London WC1E 6BT, UK. ²⁴¹Department of Medicine, University of Kuopio, Kuopio University Hospital, Kuopio 70210, Finland. ²⁴²Studies Coordinating Centre, Division of Hypertension and Cardiac Rehabilitation, Department of Cardiovascular Diseases, University of Leuven, Campus Sint Rafaël, Kapucijnenvoer 35, Block D, Box 7001, Leuven 3000, Belgium. ²⁴³Institute of Molecular and Cell Biology, University of Tartu, Riia 23, Tartu 51010, Estonia. ²⁴⁴Finnish Institute of Occupational Health, Finnish Institute of Occupational Health, Aapistie 1, Oulu 90220, Finland. ²⁴⁵Gerontology Research Center, National Institute on Aging, Baltimore, Maryland 21224, USA. ²⁴⁶Wake Forest University Health Sciences, Winston-Salem, North Carolina 27157, USA. ²⁴⁷National Heart, Lung and Blood Institute Framingham Heart Study, Framingham, 01702-5827 Massachusetts, USA. ²⁴⁸Office of Population Studies Foundation, University of San Carlos, Talamban, Cebu 6000, Philippines. ²⁴⁹Department of Epidemiology and Biostatistics, Case Western Reserve University, 2103 Cornell Road, Cleveland, Ohio 44106, USA. ²⁵⁰Department of Medicine and Neurology, University of Washington, Seattle, Washington 98195, USA. ²⁵¹MRC Epidemiology Unit, Institute of Metabolic Science, Cambridge CB2 0QQ, UK. ²⁵²Department of Twin Research and Genetic Epidemiology, King's College London, SE1 7EH London, UK. ²⁵³Department of Genetics, Biology and Biochemistry, University of Torino, Via Santena 19, Torino 10126, Italy. ²⁵⁴ALSPAC Laboratory, University of Bristol, Bristol BS8 2BN, UK. ²⁵⁵Institute of Human Genetics, Helmholtz Zentrum Munich, German Research Centre for Environmental Health, Neuherberg 85764, Germany. ²⁵⁶U872 Institut National de la Santé et de la Recherche Médicale, Centre de Recherche des Cordeliers, 75006 Paris, France. ²⁵⁷Mayo Clinic, Jacksonville, 32224 Florida, USA. ²⁵⁸Department of Basic Medical Research and Education, Ehime University Graduate School of Medicine, Toon 791-0295, Japan. ²⁵⁹Department of Geriatric Medicine, Ehime University Graduate School of Medicine, Toon 791-0295, Japan. ²⁶⁰Department of Genetics, University of North Carolina, Chapel Hill, North Carolina 27599, USA. ²⁶¹Division of Genetics, GlaxoSmithKline, Philadelphia, Pennsylvania 19101, USA. ²⁶²Department of Primary Care and Population Health, UCL, London NW3 2PF, UK. ²⁶³Department of Medicine (Geriatrics), University of Mississippi Medical Center, Jackson, 39216 Mississippi, USA. ²⁶⁴Laboratory of Cardiovascular Science, Intramural Research Program, National Institute on Aging, NIH, Baltimore, 21224-6825 Maryland, USA. ²⁶⁵Department of Geriatric Medicine, Osaka University Graduate School of Medicine, Suita 565-0871, Japan. ²⁶⁶Tohoku University, Graduate School of Pharmaceutical Sciences and Medicine, Sendai 980-8578, Japan. ²⁶⁷Genome Institute of Singapore, Agency for Science, Technology and Research, Singapore 138672, Singapore. ²⁶⁸Istituto di Neurogenetica e Neurofarmacologia, Consiglio Nazionale delle Ricerche, Cittadella Universitaria di Monserrato, 09042 Monserrato, Cagliari, Italy. ²⁶⁹Columbia University, New York, 10027 New York, USA. ²⁷⁰Amgen, 1 Kendall Square, Building 100, Cambridge, Massachusetts 02139, USA. ²⁷¹National Institute for Health and Welfare, Helsinki 00271, Finland. ²⁷²Merck Research Laboratory, 126 East Lincoln Avenue, Rahway, New Jersey 07065, USA. ²⁷³South Asia Network for Chronic Disease, Public Health Foundation of India, C-1/52, SDA, New Delhi 100016, India. ²⁷⁴Division of Biostatistics, Washington University School of Medicine, Saint Louis, Missouri 63110, USA. ²⁷⁵Center for Non-Communicable Diseases, 74800 Karachi, Pakistan. ²⁷⁶Department of Biostatistics, University of Washington, Seattle, 98105 Washington, USA. ²⁷⁷Department of Emergency and Cardiovascular Medicine, Institute of Medicine, Sahlgrenska Academy, University of Gothenburg, Gothenburg 41685, Sweden. ²⁷⁸Epidemiology & Prevention, Division of Public Health Sciences, Wake Forest University, School of Medicine, Winston-Salem, North Carolina 27157, USA. ²⁷⁹University of Ibadan, PMB 5017 Ibadan, Nigeria. ²⁸⁰Prevention and Care of Diabetes, Department of Medicine III, Medical Faculty Carl Gustav Carus at the Technical University of Dresden, Dresden 01307, Germany. ²⁸¹Department of Laboratory Medicine, Institute of Human Genetics, University of California, San Francisco, 513 Parnassus Avenue, San Francisco, California 94143, USA. ²⁸²Imperial College Cerebrovascular Unit (ICCRU), Imperial College, London W6 8RF, UK. ²⁸³Centre for Molecular Epidemiology, Yong Loo Lin School of Medicine, National University of Singapore, Singapore 117597, Singapore. ²⁸⁴Department of Medicine, Yong Loo Lin School of Medicine, National University of Singapore, Singapore 119074, Singapore. ²⁸⁵Faculty of Medicine, University of Split, 21000 Split, Croatia. ²⁸⁶Department of Biostatistics, Institute of Basic Medical Sciences, University of Oslo, Oslo 0317, Norway. ²⁸⁷Diabetes Prevention Unit, National Institute for Health and Welfare, Helsinki 00271, Finland. ²⁸⁸Lifestyle-related Disease Prevention Center, Shiga University of Medical Science, Otsu 520-2192, Japan. ²⁸⁹Department of Medical Science and Cardiorespiratory Medicine, Yokohama City University School of Medicine, Yokohama 236-0004, Japan. ²⁹⁰Tallinn University of Technology, Institute of Biomedical Engineering, Ehitajate tee 5, Tallinn 19086, Estonia. ²⁹¹Department of Epidemiology and Public Health, Imperial College, Norfolk Place, London W2 1PG, UK. ²⁹²Department of Internal Medicine, Centre Hospitalier Universitaire Vaudois, Lausanne 1011, Switzerland. ²⁹³Department of Medicine, Johns Hopkins University, Baltimore, 21205 Maryland, USA. ²⁹⁴Lapland Central Hospital, Department of Physiatry, Box 8041, Rovaniemi 96101, Finland. ²⁹⁵Department of Internal Medicine, Diabetology, and Nephrology, Medical University of Silesia, Zabrze 41-800, Poland.

AGEN Consortium

Yukinori Okada^{296,297}, Jer-Yuarn Wu^{298,299}, Dongfeng Gu³⁰⁰, Fumihiko Takeuchi³⁰¹, Atsushi Takahashi²⁹⁶, Shiro Maeda³⁰², Tatsuhiko Tsunoda³⁰³, Peng Chen³⁰⁴, Su-Chi Lim^{305,306}, Tien-Yin Wong^{307,308,309}, Jianjun Liu²⁶⁷, Terri L. Young³¹⁰, Tin Aung^{307,308}, Yik-Ying Teo^{267,283,306,311,312}, Young Jin Kim³¹³, Daehee Kang³¹⁴, Chien-Hsiun Chen^{298,299}, Fuu-Jen Tsai²⁹⁹, Li-Ching Chang²⁹⁸, S.-J. Cathy Fann²⁹⁸, Hao Mei³¹⁵, James E. Hixson³⁴, Shufeng Chen³⁰⁰, Tomohiro Katsuya^{316,317}, Masato Isono³⁰¹, Eva Albrecht⁶⁶, Kazuhiko Yamamoto³¹⁸, Michiaki Kubo³¹⁹, Yusuke Nakamura³²⁰, Naoyuki Kamatani³²¹, Norihiro Kato³⁰¹, Jiang He³¹⁵, Yuan-Tsong Chen²⁹⁸, Toshihiro Tanaka^{297,322}

²⁹⁶Laboratory for Statistical Analysis, RIKEN Center for Integrative Medical Sciences, Yokohama, Kanagawa 230-0045, Japan. ²⁹⁷Department of Human Genetics and Disease Diversity, Graduate School of Medical and Dental Sciences, Tokyo Medical and Dental University, Tokyo 113-8510, Japan. ²⁹⁸Institute of Biomedical Sciences, Academia Sinica, Taipei 115, Taiwan. ²⁹⁹School of Chinese Medicine, China Medical University, Taichung 404, Taiwan. ³⁰⁰Cardiovascular Institute and Fu Wai Hospital, Chinese Academy of Medical Sciences, Peking Union Medical College, Beijing 100037, China. ³⁰¹Department of Gene Diagnostics and Therapeutics, Research Institute, National Center for Global Health and Medicine, Tokyo 162-8655, Japan. ³⁰²Laboratory for Endocrinology and Metabolism, RIKEN Center for Integrative Medical Sciences, Yokohama, Kanagawa 230-0045, Japan. ³⁰³Laboratory for Medical Informatics, RIKEN

Center for Integrative Medical Sciences, Yokohama, Kanagawa 230-0045, Japan. ³⁰⁴Saw Swee Hock School of Public Health, National University of Singapore, Singapore 119077, Singapore. ³⁰⁵Department of Medicine, Khoo Teck Puat Hospital, Singapore 768828, Singapore. ³⁰⁶Department of Epidemiology and Public Health, Yong Loo Lin School of Medicine, National University of Singapore, Singapore 117597, Singapore. ³⁰⁷Singapore Eye Research Institute, Singapore National Eye Centre, Singapore 168751, Singapore. ³⁰⁸Department of Ophthalmology, Yong Loo Lin School of Medicine, National University of Singapore, Singapore 117597, Singapore. ³⁰⁹Centre for Eye Research Australia, University of Melbourne, East Melbourne, 3002 Victoria, Australia. ³¹⁰Center for Human Genetics, Duke University Medical Center, Durham, 27710 North Carolina, USA. ³¹¹Department of Statistics and Applied Probability, National University of Singapore, Singapore 117546, Singapore. ³¹²NUS Graduate School for Integrative Science and Engineering, National University of Singapore, Singapore 119077, Singapore. ³¹³Center for Genome Science, National Institute of Health, Osong Health Technology Administration Complex, 187 Chungcheongbuk-do, Korea. ³¹⁴Department of Preventive Medicine, Seoul National University College of Medicine, Seoul 08826, Korea. ³¹⁵Department of Epidemiology, Tulane University School of Public Health and Tropical Medicine, New Orleans, 70112 Louisiana, USA. ³¹⁶Department of Clinical Gene Therapy, Osaka University Graduate School of Medicine, Suita, Osaka 565-0871, Japan. ³¹⁷Department of Geriatric Medicine and Nephrology, Osaka University Graduate School of Medicine, Suita, Osaka 565-0871, Japan. ³¹⁸Department of Allergy and Rheumatology, Graduate School of Medicine, University of Tokyo, Tokyo 113-0033, Japan. ³¹⁹Laboratory for Genotyping Development, RIKEN Center for Integrative Medical Sciences, Yokohama, Kanagawa 230-0045, Japan. ³²⁰Laboratory of Molecular Medicine, Human Genome Center, Institute of Medical Science, University of Tokyo, Tokyo 108-8639, Japan. ³²¹Laboratory for International Alliance, RIKEN Center for Integrative Medical Sciences, Yokohama, Kanagawa 230-0045, Japan. ³²²Laboratory for Cardiovascular Diseases, RIKEN Center for Integrative Medical Sciences, Yokohama, Kanagawa 230-0045, Japan.

CARDIOGRAM

Muredach P. Reilly³²³, Heribert Schunkert^{57,324,325}, Themistocles L. Assimes³²⁶, Alistair Hall³²⁷, Christian Hengstenberg³²⁸, Inke R. König³²⁹, Reijo Laaksonen³³⁰, Ruth McPherson³³¹, John R. Thompson¹⁸⁷, Unnur Thorsteinsdottir^{332,333}, Andreas Ziegler³²⁹, Devin Absher³³⁴, Li Chen³³⁵, L. Adrienne Cupples^{13,247}, Eran Halperin^{336,337}, Mingyao Li³³⁸, Kiran Musunuru^{140,339,340}, Michael Preuss^{324,329}, Arne Schillert³²⁹, Gudmar Thorleifsson³³², George A. Wells³³⁵, Hilma Holm³³², Robert Roberts³³¹, Alexandre F.R. Stewart³³¹, Stephen Fortmann³²⁶, Alan Go³⁴¹, Mark Hlatky³²⁶, Carlos Iribarren³⁴¹, Joshua Knowles³²⁶, Richard Myers³³⁴, Thomas Quertermous³²⁶, Steven Sidney³⁴¹, Neil Risch³⁴², Hua Tang³⁴³, Stefan Blankenberg³⁴⁴, Renate Schnabel³⁴⁴, Christoph Sinning³⁴⁴, Karl J. Lackner³⁴⁵, Laurence Tiret³⁴⁶, Viviane Nicaud³⁴⁶, Francois Cambien³⁴⁶, Christoph Bickel³⁴⁴, Hans J. Rupprecht³⁴⁴, Claire Perret³⁴⁶, Carole Proust³⁴⁶, Thomas F. Münzel³⁴⁴, Maja Barbalic³⁴, Ida Yii-Der Chen²²⁰, Serkalem Demissie-Banjaw^{246,347}, Aaron Folsom³⁴⁸, Thomas Lumley²⁷⁵, Kristin Marcianti³⁴⁹, Kent D. Taylor²²⁰, Kelly Volcik³⁵⁰, Solveig Gretarsdottir³³², Jeffrey R. Gulcher³³², Augustine Kong³³², Kari Stefansson^{332,333}, Gudmundur Thorgeirsson^{333,351}, Karl Andersen^{333,351}, Marcus Fischer³²⁸, Anika Grosshennig^{324,329}, Patrick Linsel-Nitschke³²⁴, Klaus Stark³²⁸, Stefan Schreiber³⁸, Zouhair Aherrahrou^{57,324}, Petra Bruse^{57,324}, Angela Doering³⁵², Norman Klopp³⁵², Patrick Diemert³²⁴, Christina Loley^{324,329}, Anja Medack^{57,324}, Janja Nahrstedt^{324,329}, Annette Peters⁶⁸, Arnika K. Wagner³²⁴, Christina Willenborg^{57,324}, Bernhard O. Böhm³⁵³, Harald Dobnig³⁵⁴, Tanja B. Grammer³⁵⁵, Michael M. Hoffmann³⁵⁶, Andreas Meinitzer³⁵⁷, Bernhard R. Winkelmann³⁵⁸, Stefan Pilz³⁵⁴, Wilfried Renner³⁵⁷, Hubert Scharnagl³⁵⁷, Tatjana Stojakovic³⁵⁷, Andreas Tomaschitz³⁵⁴, Karl Winkler³⁵⁶, Candace Guiducci¹⁶, Noel Burt¹⁶, Stacey B. Gabriel¹⁶, Sonny Dandona³³¹, Olga Jarinova³³¹, Liming Qu³³⁸, Robert Wilensky³²³, William Matthai³²³, Hakon H. Hakonarson³⁵⁹, Joe Devaney³⁶⁰, Mary Susan Burnett³⁶⁰, Augusto D. Pichard³⁶⁰, Kenneth M. Kent³⁶⁰, Lowell Satler³⁶⁰, Joseph M. Lindsay³⁶⁰, Ron Waksman³⁶⁰, Christopher W. Knouff³⁶¹, Dawn M. Waterworth³⁶¹, Max C. Walker³⁶¹, Stephen E. Epstein³⁶⁰, Daniel J. Rader^{323,362}, Christopher P. Nelson¹⁸⁵, Benjamin J. Wright³⁶³, Anthony J. Balmforth³⁶⁴, Stephen G. Ball³⁶⁵

³²³The Cardiovascular Institute, University of Pennsylvania, Philadelphia, Pennsylvania 19104, USA. ³²⁴Institut für integrative und experimentelle Genomik, Universität zu Lübeck, Lübeck 23562, Germany. ³²⁵Deutsches Herzzentrum München, Technische Universität München, München 80636, Germany. ³²⁶Department of Medicine, Stanford University School of Medicine, Stanford, 94305-5101 California, USA. ³²⁷Division of Cardiovascular and Neuronal Remodelling, Multidisciplinary Cardiovascular Research Centre, Leeds Institute of Genetics, Health and Therapeutics, University of Leeds, Leeds LS2 9JT, UK. ³²⁸Klinik und Poliklinik für Innere Medizin II, Universität Regensburg, 93053 Regensburg, Germany. ³²⁹Institut für Medizinische Biometrie und Statistik, Universität zu Lübeck, 23562 Lübeck, Germany. ³³⁰Science Center, Tampere University Hospital, Tampere 33521, Finland. ³³¹The John & Jennifer Ruddy Canadian Cardiovascular Genetics Centre, University of Ottawa Heart Institute, Ottawa, Ontario, Canada K1Y 4W7. ³³²deCODE Genetics, Reykjavik 101, Iceland. ³³³Faculty of Medicine, University of Iceland, Reykjavik 101, Iceland. ³³⁴Hudson Alpha Institute, Huntsville, 35806 Alabama, USA. ³³⁵Cardiovascular Research Methods Centre, University of Ottawa Heart Institute, 40 Ruskin Street, Ottawa, Ontario, Canada K1Y 4W7. ³³⁶The Blavatnik School of Computer Science, Department of Molecular Microbiology and Biotechnology, Tel-Aviv University, Tel-Aviv 6997801, Israel. ³³⁷The International Computer Science Institute, Berkeley, 94704 California, USA. ³³⁸Biostatistics and Epidemiology, University of Pennsylvania, Philadelphia, 19104 Pennsylvania, USA.

³³⁹Cardiovascular Research Center and Cardiology Division, Massachusetts General Hospital, Boston, 02114 Massachusetts, USA. ³⁴⁰Center for Human Genetic Research, Massachusetts General Hospital, Boston, 02114 Massachusetts, USA. ³⁴¹Division of Research, Kaiser Permanente, Oakland, 94611 California, USA. ³⁴²Institute for Human Genetics, University of California, San Francisco, San Francisco, 94143 California, USA. ³⁴³Department of Cardiovascular Medicine, Cleveland Clinic 7255 Old Oak Blvd, Cleveland, Ohio 44130, USA. ³⁴⁴Medizinische Klinik und Poliklinik, Johannes-Gutenberg Universität Mainz, Universitätsmedizin, 55122 Mainz, Germany. ³⁴⁵Institut für Klinische Chemie und Laboratoriumsmedizin, Johannes-Gutenberg Universität Mainz, Universitätsmedizin, 55122 Mainz, Germany. ³⁴⁶INSERM UMRS 937, Pierre and Marie Curie University (UPMC, Paris 6) and Medical School, 75005 Paris, France. ³⁴⁷Boston University, School of Public Health, Boston, 02118 Massachusetts, USA. ³⁴⁸University of Minnesota School of Public Health, Division of Epidemiology and Community Health, School of Public Health (A.R.F.), Minneapolis, 55454 Minnesota, USA. ³⁴⁹University of Washington, Department of Internal Medicine, Seattle, 98195-6420 Washington, USA. ³⁵⁰University of Texas, School of Public Health, Houston, 77030 Texas, USA. ³⁵¹Department of Medicine, Landspítali University Hospital, Reykjavík 101, Iceland. ³⁵²Institute of Epidemiology, Helmholtz Zentrum München, German Research Center for Environmental Health, 85764 Neuherberg, Germany. ³⁵³Division of Endocrinology and Diabetes, Graduate School of Molecular Endocrinology and Diabetes, University of Ulm, 89069 Ulm, Germany. ³⁵⁴Division of Endocrinology, Department of Medicine, Medical University of Graz, 8010 Graz, Austria. ³⁵⁵Synlab Center of Laboratory Diagnostics Heidelberg, 69037 Heidelberg, Germany. ³⁵⁶Division of Clinical Chemistry, Department of Medicine, Albert Ludwigs University, 79085 Freiburg, Germany. ³⁵⁷Clinical Institute of Medical and Chemical Laboratory Diagnostics, Medical University Graz, 8010 Graz, Austria. ³⁵⁸Cardiology Group Frankfurt-Sachsenhausen, 60594 Frankfurt, Germany. ³⁵⁹The Center for Applied Genomics, Children's Hospital of Philadelphia, 19104 Philadelphia, Pennsylvania, USA. ³⁶⁰Cardiovascular Research Institute, Medstar Health Research Institute, Washington Hospital Center, Washington, DC 20010, USA. ³⁶¹Genetics Division and Drug Discovery, GlaxoSmithKline, King of Prussia, Pennsylvania 19406, USA. ³⁶²The Institute for Translational Medicine and Therapeutics, School of Medicine, University of Pennsylvania, Philadelphia, 19104-5158 Pennsylvania, USA. ³⁶³Department of Cardiovascular Surgery, University of Leicester, Leicester LE1 7RH, UK. ³⁶⁴Division of Cardiovascular and Diabetes Research, Multidisciplinary Cardiovascular Research Centre, Leeds Institute of Genetics, Health and Therapeutics, University of Leeds, Leeds LS2 9JT, UK. ³⁶⁵LIGHT Research Institute, Faculty of Medicine and Health, University of Leeds, Leeds LS2 9JT, UK.

CHARGE-Heart Failure Group

Laura R. Loehr^{366,367,368}, Wayne D. Rosamond³⁶⁷, Emelia Benjamin²⁴⁷, Talin Haritunians²²⁰, David Couper³⁶⁹, Joanne Murabito²⁴⁷, Ying A. Wang¹³, Bruno H. Stricker²⁴, Patricia P. Chang³⁶⁶, James T. Willerson^{370,371}


³⁶⁶Department of Medicine, University of North Carolina at Chapel Hill, North Carolina 27516, USA. ³⁶⁷Department of Epidemiology, University of North Carolina at Chapel Hill, North Carolina 27599-7435, USA. ³⁶⁸National Institute of Environmental Health Sciences, Research Triangle Park, North Carolina 27709, USA. ³⁶⁹Department of Biostatistics, University of North Carolina at Chapel Hill, North Carolina 27514, USA. ³⁷⁰University of Texas, Houston Health Science Center, Houston, Texas 77030, USA. ³⁷¹Texas Heart Institute, Houston, Texas 77225-0345, USA.

ECHOGen Consortium


Stephan B. Felix²⁰², Norbert Watzinger³⁷², Jayashri Aragam²⁴⁷, Robert Zweiker³⁷², Lars Lind³⁷³, Richard J. Rodeheffer³⁷⁴, Karin Halina Greiser³⁷⁵, Jaap W. Deckers³⁷⁶, Jan Stritzke³⁷⁷, Erik Ingelsson³⁷⁸, Iftikhar Kullo³⁷⁴, Johannes Haerting³⁷⁵, Thorsten Reffellmann²⁰², Margaret M. Redfield³⁷⁴, Karl Werdan³⁷⁹, Gary F. Mitchell²⁴⁷, Donna K. Arnett³⁸⁰, John S. Gottdiener³⁸¹, Maria Blettner³⁸², Nele Friedrich³⁸³

³⁷²Department of Internal Medicine, Division of Cardiology, Medical University Graz, Graz 8036, Austria. ³⁷³Department of Medical Sciences, Uppsala University, Uppsala 75185, Sweden. ³⁷⁴Division of Cardiovascular Diseases, Mayo Clinic, Rochester, Minnesota 55905, USA. ³⁷⁵Institute of Medical Epidemiology, Biostatistics and Informatics, Martin Luther University of Halle-Wittenberg, Halle-Wittenberg, Halle (Saale) 06097, Germany. ³⁷⁶Department of Cardiology, Erasmus University Medical Center, Rotterdam 3000 CA, The Netherlands. ³⁷⁷Medical Clinic 2, University of Lübeck, Lübeck 23538, Germany. ³⁷⁸Department of Medical Epidemiology and Biostatistics, Karolinska Institute, Stockholm 17177, Sweden. ³⁷⁹Martin Luther University, Halle-Wittenberg, Halle (Saale) 06097, Germany. ³⁸⁰Department of Epidemiology, University of Alabama at Birmingham, Birmingham, Alabama 35294-0022, USA. ³⁸¹Division of Cardiology, University of Maryland Hospital, Baltimore, Maryland 21201, USA. ³⁸²Institute of Medical Biometry, Epidemiology, and Informatics, Johannes Gutenberg University, Mainz 55101, Germany. ³⁸³Institute for Community Medicine, Ernst-Moritz-Arndt-Universität, Greifswald 17475, Germany.


Supplementary Figure 1. Q-Q plots of eGFRcrea (A), eGFRcys (B), CKD (C), and eGFRcrea in diabetes (D). $\lambda_{1000} = 1.00$ for all four analyses.^{1,2}




Supplementary Figure 2. Manhattan Plots of eGFRcrea in diabetes.



















Supplementary Figure 3. All replicated loci comparing the effect sizes for eGFR_{crea} vs eGFR_{cys}; known loci are in blue, new loci are in orange. 95% confidence intervals appear for the new loci only.




Supplementary Figure 4. Regional association plots for all 24 novel loci.









Supplementary Figure 5. Effects of known and novel loci on eGFR_{crea} in individuals with (n=16,477) and without (n=154,881) diabetes. The Pearson's correlation coefficient between the effects in the two groups was 0.80 across all loci (95% confidence interval: 0.67, 0.88). The corresponding correlation coefficient when considering all ~2.5 million genetic variants was of 0.044 (see "Test for differential effects on eGFR_{crea} between diabetes and hypertension strata", in the Methods), highlighting that most of replicated loci are associated with eGFR_{crea} regardless of the presence or absence of diabetes.



Supplementary Figure 6. Effect of the 53 known (light blue) and new (orange) loci on log(eGFR_{crea}) in the absence versus presence of hypertension. Data are derived from our previous analysis³ and based on n = 29,592 and 44,319 hypertensive and non-hypertensive subjects, respectively.



Supplementary Figure 7. Forest plots from trans-ethnic random-effect meta-analysis for all 24 novel loci. Indicated are: index SNP and effect allele; ancestry (Eu = European, As = Asian; Af = African) along with effect allele frequencies in parenthesis. Dot sizes are proportional to study sample sizes. Bars indicate 95% confidence intervals. The random-effect estimate is reported as a rhomb and as a number (b) followed by the p-value (p).



Supplementary Table 1. Sample characteristics of discovery and replication studies.

Study ¹	Sample size			Women % (N)	Age Mean (SD)	eGFRcrea Mean(SD) ml / min / 1.73m ²	eGFRcys Mean(SD) ml / min / 1.73m ²	CKD % (N)	HTN % (N)	DM % (N)
	eGFRcrea	CKD	eGFRcys							
Discovery studies										
3C	6431	6431	1243	60.8 (3911)	74(5)	73.1(16.9)	90.5(22.7)	20.4 (1311)	79.0 (6431)	9.7 (623)
Advance	2287	2287	NA	32.8 (755)	67(7)	85.1(29.2)	NA	14.7 (337)	47.6 (1096)	100.0 (2287)
AGES	3219	3219	NA	58.0 (1867)	76(5)	73.0(20.0)	NA	24.2 (781)	80.6 (2595)	11.5 (368)
Amish	1211	NA	783	48.9 (592)	49(17)	93.7(19.7)	114.9(18.0)	3.1 (37)	18.9 (229)	1.7 (20)
ARIC	8982	8982	7145	53.1 (4767)	62(6)	81.4(17.5)	84.2(19.7)	8.7 (782)	40.7 (3643)	14.2 (1276)
ASPS	848	848	NA	56.8 (482)	65(8)	96.5(39.9)	NA	8.1 (69)	72.5 (615)	9.2 (78)
AUSTWIN	9592	3320	NA	60.9 (5846)	46(13)	98.3(26.2)	NA	10.5 (350)	NA	NA
BLSA	723	723	NA	46.1 (333)	70(15)	80.3(23.1)	NA	17.4 (126)	21.9 (147)	7.7 (55)
BMES	2437	2437	NA	56.8 (1385)	69(9)	78.7(20.2)	NA	13.2 (322)	76.4 (1861)	10.9 (265)
CHS	2820	2353	2475	61.3 (1729)	72(5)	77.3(20.8)	81.0(17.9)	9.5 (224)	51.4 (1441)	11.0 (307)
CROATIA-KORCULA	888	888	NA	64.0 (568)	56(14)	87.3(20.6)	NA	7.5 (67)	54.2 (474)	13.1 (116)
CROATIA-SPLIT	478	478	NA	59.8 (286)	49(15)	104.8(23.8)	NA	5.0 (24)	39.4 (186)	5.0 (24)
CROATIA-VIS	768	768	NA	58.6 (450)	57(15)	88.2(22.1)	NA	6.9 (53)	52.2 (396)	12.0 (91)
DESIR	715	NA	NA	75.2 (550)	50(8)	92.0(16.9)	NA	0.7 (5)	NA	0.0 (0)
EGCUT 370K	863	NA	NA	51.2 (442)	37(16)	101.5(20.3)	NA	1.7 (15)	18.7 (161)	2.2 (19)
EGCUT Omni	261	261	NA	73.4 (193)	81(9)	71.2(24.1)	NA	34.2 (90)	82.5 (217)	21.8 (57)
ERF	2561	2561	NA	55.3 (1416)	49(14)	93.3(21.4)	NA	3.7 (96)	52.9 (1355)	6.1 (155)
FamHS	3838	3838	521	52.4 (2012)	52(14)	91.6(20.1)	86.3(33.5)	4.2 (161)	25.5 (979)	7.7 (291)
FHS	7782	4142	3002	54.3 (4229)	51(14)	92.1(21.7)	83.8(17.8)	10.7 (445)	57.5 (2390)	9.7 (405)
GENOA	1163	1163	NA	56.3 (655)	59(10)	87.7(24.0)	NA	10.7 (125)	73.3 (852)	15.3 (178)
HABC	1663	1663	1663	47.1 (784)	74(3)	71.2(14.8)	77.0(19.9)	25.0 (415)	63.7 (1060)	13.0 (216)
HCS	1235	1235	NA	49.6 (630)	66(7)	79.9(18.1)	NA	11.3 (144)	45 (544)	10.5 (127)
HPFS	818	818	NA	0.0 (0)	65(8)	85.2(22.7)	NA	9.5 (78)	59 (479)	100.0 (818)
HYPERGENES HTN cases	1591	1591	NA	33.0 (525)	48(10)	94.5(23.1)	NA	3.7 (59)	100.0 (1591)	0.0 (0)
HYPERGENES HTN ctrls	1662	1662	NA	39.6 (659)	60(10)	87.8(19.0)	NA	4.9 (81)	0.0 (0)	0.0 (0)
INCIPE	940	940	875	52.7 (495)	61(11)	82.6(18.5)	88.5(22.1)	8.6 (81)	69.6 (654)	10.6 (100)

Supplementary Table 1 (continued).

Study ¹	Sample size			Women % (N)	Age Mean (SD)	eGFRcrea Mean(SD) ml / min / 1.73m ²	eGFRcys Mean(SD) ml / min / 1.73m ²	CKD % (N)	HTN % (N)	DM % (N)
	eGFRcrea	CKD	eGFRcys							
Discovery studies										
INGI-CARLANTINO	447	NA	NA	60.8 (272)	50(16)	93.9(22.4)	NA	NA	34.9 (156)	9.4 (42)
INGI-CILENTO	821	821	NA	54.9 (451)	54(18)	88.7(21.8)	NA	8.0 (66)	38.4 (315)	10.5 (86)
INGI-FVG	874	874	NA	59.4 (519)	52(16)	90.6(21.8)	NA	6.0 (52)	48.8 (427)	6.7 (59)
INGI-VAL BORBERA	1636	1636	NA	55.8 (913)	55(18)	89.2(23.3)	NA	8.5 (139)	43.8 (717)	6.5 (107)
JUPITER	8780	8780	NA	32.2 (2827)	66(8)	80.1(18.1)	NA	11.5 (1008)	55.8 (4901)	0.4 (37)
KORA-F3	1641	1641	1642	50.5 (831)	62(10)	83.9(21.0)	111.8(26.3)	10.8 (177)	41.1 (674)	11.1 (179)
KORA-F4	1814	1814	1811	51.3 (930)	61(9)	85.1(20.2)	109.7(26.2)	7.0 (127)	20.9 (379)	9.2 (167)
MESA	2521	2521	2521	52.0 (1311)	63(10)	82.4(18.3)	90.0(21.7)	9.7 (245)	38.6 (974)	6.0 (151)
MICROS	1201	1201	1198	56.5 (678)	46(16)	94.6(20.9)	107.4(23.8)	3.8 (46)	37.7 (437)	4.3 (49)
NESDA	1856	NA	1856	67.5 (1253)	42(12)	98.4(20.7)	112.4(24.9)	1.1 (20)	30.3 (563)	5.1 (95)
NHS	786	786	NA	100.0 (786)	59(6)	86.2(22.1)	NA	10.7 (84)	70 (554)	100.0 (786)
NSPHS	565	565	NA	53.1 (300)	52(18)	91.0(22.1)	NA	5.7 (32)	43.4 (242)	7.8 (44)
OGP-TALANA	862	862	NA	57.3 (494)	51(19)	91.2(23.6)	NA	7.5 (65)	37.3 (322)	5.1 (44)
ORCADES	704	704	NA	53.6 (377)	54(15)	89.4(20.7)	NA	6.8 (48)	41.8 (287)	4.0 (28)
POPGEN	1163	1163	NA	44.4 (516)	55(14)	88.1(18.8)	NA	5.1 (59)	46.8 (541)	3.8 (44)
PREVEND	NA	NA	1885	48.9 (923)	50(12)	NA	105.4(25.3)	NA	33.5 (631)	3.7 (70)
PROSPER- PHASE	5236	5236	NA	51.7 (2718)	75(3)	72.0(21.4)	NA	29.6 (1549)	62.1 (3251)	10.4 (544)
RS-I	4390	4390	NA	61.4 (2696)	70(9)	77.1(17.2)	NA	13.7 (600)	34.1 (1497)	10.7 (470)
RS-II	1863	1863	NA	54.5 (1015)	65(8)	81.3(17.2)	NA	9.1 (169)	28.4 (530)	11.1 (207)
SAPALDIA	1444	1444	NA	51.0 (737)	52(11)	90.3(17.3)	NA	3.1 (44)	27.4 (389)	2.8 (40)
SHIP	3228	3228	3228	51.7 (1670)	54(15)	90.4(23.6)	97.1(25.3)	7.7 (248)	51.1 (1649)	11.2 (362)
SHIP-TREND	986	986	986	56.2 (554)	50(14)	92.4(22.1)	122.1(22.1)	4.3 (42)	39.6 (390)	1.8 (18)
SORBS	856	856	NA	58.5 (501)	49(16)	92.2(19.0)	NA	4.1 (35)	53.2 (455)	9.3 (80)
WGHS	21940	23186	NA	100.0 (21,940)	55(7)	90.0(22.5)	NA	6.1 (1329)	24.5 (5374)	2.5 (554)
YFS	2023	NA	NA	54.7 (1107)	38(5)	100.3(15.8)	NA	0.2 (5)	20.0 (404)	1.9 (39)

Supplementary Table 1 (continued).

Study ¹	Sample size			Women % (N)	Age Mean (SD)	eGFRcrea Mean(SD) ml / min / 1.73m ²	eGFRcys Mean(SD) ml / min / 1.73m ²	CKD % (N)	HTN % (N)	DM % (N)
	eGFRcrea	CKD	eGFRcys							
Replication studies										
Bus Santé	4408	4408	NA	49.4 (2178)	58(11)	85.6(15.6)	NA	4.3 (186)	28.3 (1249)	7.4 (327)
EGCUT_replic	1519	1519	1037	46.2 (703)	58(17)	97.4(29.3)	85.5(16.9)	8.8 (134)	59.5 (905)	6.3 (97)
ESTHER	3604	3604	NA	55.6 (2004)	62(7)	90.3(34.4)	NA	15.7 (565)	57.5 (2073)	15.9 (572)
GENDIAN	450	450	450	47.1 (212)	65(10)	69.1(20.2)	85.3(27.1)	32.3 (145)	53.0 (237)	100.0 (450)
GHS 1	2995	2995	NA	48.5 (1452)	56(11)	87.3(16.5)	NA	3.7 (112)	52.6 (1575)	7.2 (215)
GHS 2	1179	1179	NA	50.0 (590)	55(11)	86.7(16.2)	NA	4.5 (53)	48.4 (570)	7.6 (90)
GSK	1721	1721	NA	66.6 (1147)	51(13)	92.3(22.6)	NA	5.5 (95)	43.7 (752)	4.6 (80)
HRS	NA	NA	7700	58.9 (4537)	68(10)	NA	102.5(20.3)	2.9 (221)	66.1 (5087)	20.5 (1557)
KORA-F3 non-GWAS	1494	NA	1493	52.5 (787)	52(13)	92.6(21.3)	123.5(29.1)	2.6 (39)	29.4 (437)	5.1 (76)
KORA-F4 non-GWAS	1199	1200	1196	52.4 (629)	49(15)	92.6(22.4)	118.4(27.5)	5.8 (70)	13.3 (159)	4.0 (48)
IPM_EA_Affy	440	NA	NA	30.2 (133)	62(13)	94.8(36.8)	NA	19.1 (84)	60.5 (266)	26.0 (99)
IPM_EA_Illu	1307	1307	NA	48.6 (635)	68(9)	86.1(27.8)	NA	14.8 (194)	55.1 (720)	15.0 (185)
LURIC	3056	3056	3054	30.0 (917)	63(11)	86.0(21.7)	84.7(22.6)	10.0 (305)	72.7 (2223)	32.6 (996)
OGP	9554	5884	NA	56.9 (5440)	50(17)	98.6(35.0)	NA	8.1 (776)	36.0 (3443)	6.2 (596)
SAPHIR	1721	NA	NA	37.1 (639)	51(6)	91.7(16.1)	NA	6.9 (19)	55.7 (959)	3.3 (56)
SKIPOGH	870	NA	NA	52.3 (455)	47(18)	94.2(24.6)	NA	5.7 (50)	22.9 (198)	4.5 (39)
Vanderbilt Omni1	3221	3221	NA	47.3 (1525)	54(19)	80.0(36.9)	NA	27.7 (891)	70.5 (2271)	18.0 (581)
Vanderbilt Omni5	1129	1129	NA	46.9 (529)	50(21)	89.0(44.4)	NA	21.7 (245)	58.2 (657)	33.3 (376)
Vanderbilt 660W	2299	2299	NA	56.5 (1298)	56(17)	78.6(23.9)	NA	20.6 (474)	57.2 (1316)	17.9 (411)

¹ Extended study names are given in the Acknowledgements section.

Supplementary Table 2. Association of previously identified loci. For each locus, we report the best SNP from the current discovery GWAS (current best SNP). The previously reported best SNP is reported for comparison.

ID of current best SNP in the locus	Chr	Position (Build 37)	Locus name*	Ref. / Non-Ref. All. (Ref All Freq)	eGFRcrea		CKD		Previously reported SNP in the locus		
					beta(SE) [†]	P-value [†]	Odds Ratio (95%CI) [†]	P-value [†]	ID of previously reported SNP ^{REF}	LD# with current best SNP r ² /D'	P-value in current GWAS [†]
rs1800615	1	15,832,281	CASP9	T/C(0.30)	-0.0058(0.0009)	1.90E-09	1.04 (1.00,1.07)	3.40E-02	rs12124078 ³	1.00/1.00	3.09E-09
rs267734	1	150,951,477	LASS2	T/C(0.79)	-0.0079(0.0011)	4.01E-13	1.06 (1.02,1.10)	3.52E-03	same ⁴		
rs807601	2	15,793,014	DDX1	T/G(0.34)	0.0064(0.0009)	6.60E-12	0.99 (0.96,1.02)	5.28E-01	rs6431731 ³	0.04/0.60	2.95E-07
rs1260326	2	27,730,940	GCKR	T/C(0.42)	0.0068(0.0009)	3.38E-14	0.98 (0.95,1.01)	1.36E-01	same ⁴		
rs6546838	2	73,679,280	ALMS1	A/G(0.76)	-0.0093(0.0010)	7.72E-20	1.05 (1.02,1.09)	4.63E-03	rs13538 ⁴	0.95/1.00	3.15E-17
rs7422339	2	211,540,507	CPS1	A/C(0.32)	-0.0106(0.0010)	2.18E-23	1.11 (1.07,1.15)	7.54E-09	same ⁵		
rs2861422	3	141,724,644	TFDP2	T/C(0.27)	0.0074(0.0010)	9.12E-14	0.96 (0.92,0.99)	7.59E-03	rs347685 ⁴	0.96/1.00	1.87E-13
rs17319721	4	77,368,847	SHROOM3	A/G(0.43)	-0.0114(0.0009)	1.32E-37	1.07 (1.04,1.10)	7.71E-06	same ⁵		
rs11959928	5	39,397,132	DAB2	A/T(0.44)	-0.0083(0.0009)	1.66E-20	1.06 (1.02,1.09)	4.20E-04	same ⁴		
rs6420094	5	176,817,636	SLC34A1	A/G(0.66)	0.0096(0.0010)	4.92E-22	0.91 (0.88,0.94)	3.68E-09	same ⁴		
rs9472135	6	43,809,802	VEGFA	T/C(0.71)	-0.0080(0.0010)	3.34E-15	1.07 (1.04,1.11)	3.37E-05	rs881858 ⁴	0.88/0.96	7.51E-15
rs316009	6	160,675,764	SLC22A2	T/C(0.10)	0.0131(0.0014)	4.38E-19	0.96 (0.91,1.01)	9.71E-02	rs2279463 ⁴	0.01/1.00	2.94E-16
rs848490	7	77,555,005	TMEM60	C/G(0.73)	0.0073(0.0010)	7.80E-13	0.93 (0.90,0.97)	1.44E-04	rs6465825 ⁴	0.41/1.00	4.91E-12
rs7805747	7	151,407,801	PRKAG2	A/G(0.25)	-0.0130(0.0011)	7.96E-29	1.16 (1.11,1.20)	2.06E-14	same ⁴		
rs3758086	8	23,714,992	STC1	A/G(0.42)	-0.0071(0.0009)	1.71E-15	1.06 (1.02,1.09)	4.13E-04	rs10109414 ⁵	1.00/1.00	2.31E-15
rs4744712	9	71,434,707	PIP5K1B	A/C(0.40)	-0.0071(0.0009)	4.29E-15	1.06 (1.03,1.09)	7.59E-05	same ⁴		
rs1044261	10	1,065,710	WDR37	T/C(0.08)	-0.0113(0.0016)	1.21E-11	1.15 (1.09,1.22)	3.62E-07	rs10794720 ⁴	0.42/0.69	3.23E-09
rs963837	11	30,749,090	MPPED2	T/C(0.54)	-0.0078(0.0009)	5.69E-18	1.09 (1.05,1.12)	9.03E-08	rs3925584 ³	0.93/0.97	7.58E-18
rs10774021	12	349,298	SLC6A13	T/C(0.65)	-0.0063(0.0009)	4.77E-12	1.04 (1.01,1.07)	1.17E-02	same ⁴		
rs716877	13	72,347,448	DACH1	C/G(0.40)	0.0049(0.0009)	6.22E-08	0.97 (0.94,1.00)	3.15E-02	rs626277 ⁴	1.00/1.00	7.98E-08
rs476633	15	41,392,134	INO80	C/G(0.57)	0.0051(0.0009)	8.90E-09	0.94 (0.91,0.97)	2.84E-05	rs2928148 ³	0.84/1.00	1.28E-07
rs2467853	15	45,698,793	GATM	T/G(0.62)	0.0126(0.0009)	1.05E-42	0.90 (0.87,0.93)	1.88E-11	rs2453533 ⁵	1.00/1.00	4.26E-42
rs491567	15	53,946,593	WDR72	A/C(0.78)	-0.0084(0.0010)	2.86E-15	1.08 (1.04,1.12)	7.48E-05	same ⁴		
rs1394125	15	76,158,983	UBE2Q2	A/G(0.35)	-0.0073(0.0010)	5.47E-14	1.07 (1.04,1.11)	3.02E-05	same ⁴		
rs13329952	16	20,366,507	UMOD	T/C(0.81)	-0.0158(0.0011)	9.47E-43	1.24 (1.19,1.29)	1.98E-25	rs12917707 ⁵	0.95/1.00	1.16E-41
rs2453580	17	19,438,321	SLC47A1	T/C(0.59)	0.0064(0.0009)	2.93E-11	0.94 (0.91,0.97)	2.60E-04	same ³		
rs9916302	17	37,499,949	CDK12 / FBXL20	T/C(0.74)	-0.008(0.0010)	4.78E-15	1.03 (0.99,1.07)	9.43E-02	rs7208487 ³ rs11078903 ⁶	0.49/1.00 0.87/1.00	1.53E-12 5.09E-13
rs11657044	17	59,450,105	BCAS3	T/C(0.19)	-0.0115(0.0012)	7.89E-22	1.06 (1.02,1.10)	6.56E-03	rs9895661 ⁴	0.94/1.00	2.76E-21
rs12460876	19	33,356,891	SLC7A9	T/C(0.60)	-0.0066(0.0009)	1.86E-13	1.05 (1.02,1.08)	2.19E-03	same ⁴		

*Based on previously reported index gene to facilitate comparison.

[†]Beta is the effect on log(eGFRcrea in ml/min/1.73 m²); all reported standard errors, confidence intervals, and P-values, are based on the twice-GC corrected results from discovery GWAS meta-analysis.

[#]LD lookup is based on HapMap 22 CEU obtained using SNAP⁷ version 2.2.

Supplementary Table 3. All SNPs tested for replication.

SNPID Locus name*	Chr	Position (bp) (Build 37)	Eff./ Non Eff All (EAF)#	STAGE 1 (discovery)†			STAGE 2 (replication)‡				Combined analysis					
				N	Beta (SE)	P-value	N	Beta (SE)	1-sided P-value	q-value	N	Beta (SE)	P-value	I² (%)	Power	Median Rs _q
eGFRcrea in the non-diabetes group																
rs3850625 CACNA1S	1	201,016,296	A/G(0.12)	116,689	0.0086 (0.0014)	2.55E-09	36,418	0.0071 (0.0028)	5.10E-03	5.46E-03	153,107	0.0083 (0.0013)	6.82E-11	0	0.997	1.00
rs2712184 IGFBP5	2	217,682,779	A/C(0.58)	118,440	-0.0052 (0.0009)	1.65E-08	35,414	-0.0055 (0.0018)	1.20E-03	2.06E-03	153,854	-0.0053 (0.0008)	1.33E-10	0	0.983	1.00
rs9682041 SKIL	3	170,091,902	T/C(0.87)	118,454	-0.0072 (0.0013)	1.36E-07	32,457	-0.0046 (0.0031)	6.81E-02	2.33E-02	150,911	-0.0068 (0.0012)	2.58E-08	2	0.948	1.00
rs10513801 ETV5	3	185,822,353	T/G(0.87)	118,374	0.0079 (0.0013)	3.80E-09	36,400	0.0046 (0.0026)	3.96E-02	1.79E-02	154,774	0.0072 (0.0012)	1.03E-09	0	0.992	1.00
rs10994860 A1CF	10	52,645,424	T/C(0.18)	118,358	0.0082 (0.0012)	1.00E-11	36,286	0.0061 (0.0024)	5.10E-03	5.46E-03	154,644	0.0077 (0.0011)	1.07E-12	2	0.999	0.93
rs163160 KCNQ1	11	2,789,955	A/G(0.82)	118,373	0.0069 (0.0012)	9.02E-09	36,311	0.0050 (0.0023)	1.62E-02	9.89E-03	154,684	0.0065 (0.0011)	2.26E-09	14	0.992	0.98
rs164748 DPEP1	16	89,708,292	C/G(0.53)	118,373	0.0053 (0.0009)	9.92E-09	36,124	0.0019 (0.0018)	1.42E-01	4.19E-02	154,497	0.0046 (0.0008)	1.95E-08	17	0.991	0.99
rs8091180 NFATC1	18	77,164,243	A/G(0.56)	117,447	-0.0062 (0.0011)	1.43E-08	36,268	-0.0052 (0.0020)	5.10E-03	5.46E-03	153,715	-0.0060 (0.0010)	1.28E-09	0	0.999	0.81
rs437065 ADAMTS5	21	28,527,399	C/G(0.14)	118,358	0.0066 (0.0013)	8.64E-07	35,562	-0.0026 (0.0026)	1.63E-01	4.62E-02	153,920	0.0047 (0.0012)	6.78E-05	0	0.904	0.99
eGFRcrea in the overall sample																
rs12136063 SYPL2	1	110,014,170	A/G(0.70)	133,723	0.0049 (0.0009)	2.33E-07	41,703	0.0028 (0.0019)	6.48E-02	2.31E-02	175,426	0.0045 (0.0008)	4.71E-08	0	0.922	1.00
rs2802729 SDCCAG8	1	243,501,763	A/C(0.44)	133,608	-0.0050 (0.0009)	7.37E-08	41,200	-0.0029 (0.0018)	5.02E-02	2.05E-02	174,808	-0.0046 (0.0008)	2.20E-08	9	0.983	0.88
rs2888875 THADA	2	43,788,092	A/G(0.69)	129,114	0.0048 (0.0009)	7.75E-07	41,571	0.0012 (0.0018)	2.60E-01	5.72E-02	170,685	0.0041 (0.0008)	6.92E-07	0	0.894	0.99
rs17050272 GLI2	2	121,306,440	A/G(0.43)	132,764	-0.0048 (0.0009)	6.63E-07	41,681	-0.0024 (0.0018)	8.88E-02	2.93E-02	174,445	-0.0043 (0.0008)	1.36E-07	19	0.964	0.90
rs3820716 ACVR2A	2	148,680,260	A/G(0.53)	133,751	-0.0052 (0.0009)	2.73E-09	41,642	0.0006 (0.0017)	6.36E-01	1.22E-01	175,393	-0.0039 (0.0008)	1.36E-06	17	0.994	1.00
rs4667594 LRP2	2	170,008,506	A/T(0.53)	133,715	-0.0045 (0.0009)	2.37E-07	41,622	-0.0043 (0.0017)	5.90E-03	5.62E-03	175,337	-0.0044 (0.0008)	3.52E-08	4	0.922	1.00
rs6795744 WNT7A	3	13,906,850	A/G(0.15)	133,718	0.0071 (0.0012)	9.60E-09	41,772	0.0019 (0.0024)	2.10E-01	5.15E-02	175,490	0.0060 (0.0011)	3.33E-08	18	0.990	0.98

Supplementary Table 3 (continued).

SNPID Locus name*	Chr	Position (bp) (Build 37)	Eff./ Non Eff All (EAF) [#]	STAGE 1 (discovery) [†]			STAGE 2 (replication) [‡]				Combined analysis					
				N	Beta (SE)	P-value	N	Beta (SE)	1-sided P-value	q-value	N	Beta (SE)	P-value	I ² (%)	Power	Median Rsq
eGFRcrea in the overall sample																
rs6795744 WNT7A	3	13,906,850	A/G(0.15)	133,718	0.0071 (0.0012)	9.60E-09	41,772	0.0019 (0.0024)	2.10E-01	5.15E-02	175,490	0.0060 (0.0011)	3.33E-08	18	0.990	0.98
rs16852193 EGFEM1P	3	168,074,912	T/C(0.09)	133,691	0.0076 (0.0015)	8.72E-07	41,664	0.0005 (0.0030)	4.34E-01	8.86E-02	175,355	0.0061 (0.0014)	6.64E-06	0	0.885	0.95
rs228611 NFKB1	4	103,561,709	A/G(0.48)	133,788	-0.0055 (0.0009)	4.66E-10	41,657	-0.0060 (0.0017)	2.08E-04	8.91E-04	175,445	-0.0056 (0.0008)	3.58E-12	4	0.998	1.00
rs7735249 ARL15	5	53,310,139	C/G(0.92)	132,513	-0.0109 (0.0018)	2.09E-09	41,662	-0.0012 (0.0028)	3.28E-01	6.86E-02	174,175	-0.0079 (0.0015)	2.08E-07	8	1.000	0.98
rs11960179 PIK3R1	5	67,820,217	A/G(0.12)	133,714	-0.0072 (0.0014)	2.47E-07	23,693	-0.0051 (0.0033)	6.16E-02	2.31E-02	157,407	-0.0069 (0.0013)	1.57E-07	9	0.908	0.94
rs836788 DHFR	5	79,912,044	T/C(0.35)	133,810	-0.0045 (0.0009)	8.56E-07	41,731	-0.0038 (0.0018)	1.73E-02	9.89E-03	175,541	-0.0043 (0.0008)	9.89E-08	11	0.868	1.00
rs7759001 ZNF204	6	27,341,409	A/G(0.76)	133,723	-0.0053 (0.0010)	2.64E-07	41,760	-0.0045 (0.0020)	1.38E-02	9.10E-03	175,483	-0.0051 (0.0009)	1.75E-08	0	0.930	1.00
rs10277115 UNCX	7	1,285,195	A/T(0.24)	115,895	0.0095 (0.0014)	1.05E-10	40,626	0.0079 (0.0023)	3.16E-04	9.03E-04	156,521	0.0090 (0.0012)	8.72E-14	0	1.000	0.53
rs2290263 MIR148A	7	25,887,278	A/G(0.73)	132,930	0.0053 (0.0010)	2.35E-07	41,756	0.0016 (0.0019)	2.04E-01	5.15E-02	174,686	0.0045 (0.0009)	6.70E-07	5	0.959	1.00
rs3750082 KBTBD2	7	32,919,927	A/T(0.34)	127,284	0.0049 (0.0009)	2.52E-07	41,210	0.0031 (0.0018)	4.58E-02	1.96E-02	168,494	0.0045 (0.0008)	3.22E-08	2	0.934	1.00
rs1004402 GRB10	7	50,755,208	A/C(0.12)	127,292	0.0068 (0.0013)	4.79E-07	41,801	-0.0026 (0.0027)	8.31E-01	1.52E-01	169,093	0.0050 (0.0012)	2.93E-05	11	0.882	1.00
rs868055 UBE2H	7	129,435,191	T/C(0.10)	133,608	0.0082 (0.0016)	3.48E-07	41,843	-0.0008 (0.0029)	6.14E-01	1.22E-01	175,451	0.0060 (0.0014)	2.17E-05	22	0.981	0.83
rs6459680 RNF32	7	156,258,568	T/G(0.74)	133,692	-0.0065 (0.0010)	1.96E-10	41,655	-0.0019 (0.0019)	1.67E-01	4.62E-02	175,347	-0.0055 (0.0009)	1.07E-09	0	1.000	0.98
rs913423 CYP26A1	10	94,845,036	A/G(0.54)	132,861	-0.0052 (0.0009)	5.10E-09	41,638	-0.0014 (0.0017)	2.07E-01	5.15E-02	174,499	-0.0043 (0.0008)	7.65E-08	10	0.993	1.00
rs4014195 AP5B1	11	65,506,822	C/G(0.64)	133,723	0.0061 (0.0009)	2.19E-11	41,677	0.0034 (0.0018)	2.65E-02	1.42E-02	175,400	0.0055 (0.0008)	1.10E-11	0	1.000	1.00
rs10491967 TSPAN9	12	3,368,093	A/G(0.11)	133,800	-0.0092 (0.0014)	3.03E-10	41,870	-0.0106 (0.0027)	4.59E-05	3.93E-04	175,670	-0.0095 (0.0013)	5.18E-14	0	1.000	0.99
rs7956634 PTPRO	12	15,321,194	T/C(0.81)	133,722	-0.0068 (0.0011)	2.46E-09	41,726	-0.0069 (0.0022)	7.04E-04	1.51E-03	175,448	-0.0068 (0.0010)	7.17E-12	0	0.997	1.00

Supplementary Table 3 (continued).

SNPID Locus name*	Chr	Position (bp) (Build 37)	Eff./ Non Eff All (EAF) [#]	STAGE 1 (discovery) [†]			STAGE 2 (replication) [‡]				Combined analysis					
				N	Beta (SE)	P-value	N	Beta (SE)	1-sided P-value	q-value	N	Beta (SE)	P-value	I ² (%)	Power	Median Rsq
eGFRcrea in the overall sample																
rs1106766 <i>INHBC</i>	12	57,809,456	T/C(0.22)	133,660	0.0062 (0.0011)	4.67E-08	21,005	0.0058 (0.0026)	1.23E-02	8.79E-03	154,665	0.0061 (0.0010)	2.41E-09	11	0.973	0.84
rs11180732 <i>PHLDA1</i>	12	76,283,354	T/G(0.31)	133,656	0.0049 (0.0009)	4.65E-07	41,586	0.0008 (0.0018)	3.26E-01	6.86E-02	175,242	0.0041 (0.0008)	6.23E-07	0	0.930	0.95
rs2071047 <i>BMP4</i>	14	54,418,411	A/G(0.41)	133,647	0.0044 (0.0009)	9.15E-07	41,772	0.0016 (0.0017)	1.85E-01	4.95E-02	175,419	0.0038 (0.0008)	3.20E-06	13	0.879	1.00
rs8056893 <i>SLC7A6</i>	16	68,304,392	A/C(0.72)	133,711	0.0051 (0.0010)	2.03E-07	41,695	0.0035 (0.0019)	3.21E-02	1.53E-02	175,406	0.0047 (0.0009)	1.28E-07	0	0.940	0.98
rs9807656 <i>SETBP1</i>	18	42,346,956	T/C(0.91)	133,688	-0.0081 (0.0016)	6.06E-07	40,684	0.0020 (0.0029)	7.61E-01	1.42E-01	174,372	-0.0056 (0.0014)	7.26E-05	22	0.947	0.99
rs9945268 <i>RNF152</i>	18	59,340,526	A/T(0.70)	127,290	-0.0048 (0.0010)	7.05E-07	41,612	-0.0020 (0.0018)	1.32E-01	4.05E-02	168,902	-0.0042 (0.0009)	3.18E-06	0	0.876	1.00
rs11666497 <i>SIPA1L3</i>	19	38,464,262	T/C(0.18)	127,271	-0.0064 (0.0012)	8.58E-08	41,640	-0.0041 (0.0022)	3.06E-02	1.53E-02	168,911	-0.0058 (0.0011)	4.25E-08	24	0.973	0.97
rs12975033 <i>IZUMO1</i>	19	49,249,443	A/T(0.55)	133,724	-0.0052 (0.0009)	4.75E-09	38,469	0.0006 (0.0017)	6.43E-01	1.22E-01	172,193	-0.0039 (0.0008)	1.31E-06	26	0.992	1.00
rs6088580 <i>TP53INP2</i>	20	33,285,053	C/G(0.47)	132,773	-0.0055 (0.0009)	7.17E-10	34,592	-0.0027 (0.0017)	6.28E-02	2.31E-02	167,365	-0.0049 (0.0008)	1.79E-09	0	0.997	0.99
rs2235808 <i>JPH2</i>	20	42,815,795	C/G(0.85)	129,605	0.0071 (0.0013)	1.50E-07	25,036	0.0020 (0.0025)	2.17E-01	5.17E-02	154,641	0.0060 (0.0012)	3.21E-07	0	0.967	0.87
rs17216707 <i>BCAS1</i>	20	52,732,362	T/C(0.79)	133,656	-0.0084 (0.0011)	5.96E-13	39,971	-0.0051 (0.0021)	7.80E-03	6.69E-03	173,627	-0.0077 (0.0010)	8.83E-15	1	1.000	0.85
CKD in the overall sample																
rs1032843 <i>FIGN</i>	2	164,184,329	T/C(0.07)	118,131	0.1508 (0.0299)	4.48E-07	32,338	0.0332 (0.0486)	2.47E-01	5.72E-02	150,469	0.1174 (0.0259)	5.70E-06	2	0.915	0.95
rs11039182 <i>MADD</i>	11	47,346,723	T/C(0.70)	118,130	-0.0818 (0.0167)	9.07E-07	35,062	-0.0325 (0.0271)	1.15E-01	3.66E-02	153,192	-0.0678 (0.0144)	2.64E-06	0	0.722	0.99
rs593790 <i>OR4C45</i>	11	48,391,116	A/T(0.82)	118,137	-0.1045 (0.0209)	5.96E-07	35,015	-0.0213 (0.0320)	2.53E-01	5.72E-02	153,152	-0.0788 (0.0178)	9.77E-06	0	0.956	0.99
rs2832559 <i>GRIK1</i>	21	31,417,144	T/C(0.91)	117,017	-0.1435 (0.0292)	9.56E-07	30,577	0.0668 (0.0540)	8.92E-01	1.61E-01	147,594	-0.0941 (0.0262)	3.31E-04	0	0.889	0.97

Supplementary Table 3 (continued).

SNPID Locus name*	Chr	Position (bp) (Build 37)	Eff./ Non Eff All (EAF) [#]	STAGE 1 (discovery) [†]			STAGE 2 (replication) [‡]				Combined analysis					
				N	Beta (SE)	P-value	N	Beta (SE)	1-sided P-value	q-value	N	Beta (SE)	P-value	I ² (%)	Power	Median Rs _q
eGFRcys in the overall sample																
rs12428035 <i>DZIP1</i>	13	96,300,872	T/C(0.12)	33,145	-0.0144 (0.0029)	7.28E-07	14,919	-0.0089 (0.0039)	1.13E-02	8.79E-03	48,064	-0.0125 (0.0023)	6.33E-08	0	0.588	1.00

*Based on Table 1 for replicated loci and on the closest gene name for the non-replicated loci, according to the UCSC Genome Browser on Human Feb. 2009 (GRCh37/hg19) Assembly.

[#]Effect Allele Frequency as estimated in the combined sample.

[†]All P-values and SEs reported from the STAGE 1 analysis were twice-GC corrected, to account for possible genomic inflation.

[‡]In Stage 2 analysis, 1-sided P-values were computed to assess direction consistent effects with Stage 1; q-values were estimated based on the 1-sided P-values.

Supplementary Table 4. Association of replicated novel loci with eGFRcrea, CKD, and eGFRcys, in the overall combined sample.

SNP ID	Chr	Position (bp) (Build 37)	Locus name*	Eff./ Non Eff. All. (EAF)	eGFRcrea			CKD		eGFRcys	
					beta (SE)	P-value	beta (SE) from random-effect meta-analysis	OR (95%CI)	P-value	beta (SE)	P-value
rs12136063	1	110,014,170	<i>SYPL2</i>	A/G(0.70)	0.0045(0.0008)	4.71E-08	0.0045(0.0008)	0.98(0.96,1.01)	2.37E-01	0.0025(0.0016)	1.25E-01
rs3850625	1	201,016,296	<i>CACNA1S</i>	A/G(0.12)	0.0078(0.0013)	5.53E-10	0.0079(0.0012)	0.95(0.91,0.99)	2.31E-02	0.0016(0.0023)	4.95E-01
rs2802729	1	243,501,763	<i>SDCCAG8</i>	A/C(0.43)	-0.0046(0.0008)	2.20E-08	-0.0047(0.0009)	1.05(1.02,1.08)	4.10E-04	-0.0036(0.0016)	2.49E-02
rs4667594	2	170,008,506	<i>LRP2</i>	A/T(0.53)	-0.0044(0.0008)	3.52E-08	-0.0045(0.0008)	1.05(1.02,1.07)	5.43E-04	0.0001(0.0015)	9.63E-01
rs2712184	2	217,682,779	<i>IGFBP5</i>	A/C(0.58)	-0.0048(0.0008)	3.02E-09	-0.0047(0.0009)	1.02(0.99,1.05)	1.54E-01	0.0001(0.0015)	9.39E-01
rs6795744	3	13,906,850	<i>WNT7A</i>	A/G(0.15)	0.0060(0.0011)	3.33E-08	0.0057(0.0013)	0.96(0.93,1.00)	3.98E-02	0.0049(0.0021)	1.84E-02
rs9682041	3	170,091,902	<i>SKIL</i>	T/C(0.87)	-0.0062(0.0012)	2.95E-07	-0.0062(0.0012)	1.02(0.98,1.06)	2.53E-01	-0.0039(0.0022)	7.99E-02
rs10513801	3	185,822,353	<i>ETV5</i>	T/G(0.87)	0.0070(0.0012)	2.47E-09	0.0069(0.0012)	0.93(0.89,0.96)	5.08E-05	0.0019(0.0023)	3.92E-01
rs228611	4	103,561,709	<i>NFKB1</i>	A/G(0.47)	-0.0056(0.0008)	3.58E-12	-0.0056(0.0008)	1.03(1.00,1.05)	3.36E-02	-0.0023(0.0015)	1.23E-01
rs7759001	6	27,341,409	<i>ZNF204</i>	A/G(0.76)	-0.0051(0.0009)	1.75E-08	-0.0052(0.0009)	1.03(1.00,1.06)	5.66E-02	-0.0023(0.0017)	1.71E-01
rs10277115	7	1,285,195	<i>UNCX</i>	A/T(0.23)	0.0090(0.0012)	8.72E-14	0.0090(0.0012)	0.96(0.92,1.00)	4.09E-02	0.0045(0.0025)	7.62E-02
rs3750082	7	32,919,927	<i>KBTD2</i>	A/T(0.33)	0.0045(0.0008)	3.22E-08	0.0045(0.0009)	0.97(0.95,1.00)	5.76E-02	0.0014(0.0016)	3.95E-01
rs6459680	7	156,258,568	<i>RNF32</i>	T/G(0.74)	-0.0055(0.0009)	1.07E-09	-0.0055(0.0009)	1.04(1.01,1.07)	2.08E-02	-0.0043(0.0017)	1.06E-02
rs10994860	10	52,645,424	<i>A1CF</i>	T/C(0.19)	0.0071(0.0010)	1.66E-12	0.0069(0.0011)	0.99(0.95,1.02)	5.14E-01	0.0015(0.0020)	4.41E-01
rs163160	11	2,789,955	<i>KCNQ1</i>	A/G(0.82)	0.0064(0.0010)	1.72E-10	0.0063(0.0011)	0.95(0.92,0.99)	5.30E-03	0.0073(0.0020)	2.35E-04
rs4014195	11	65,506,822	<i>AP5B1</i>	C/G(0.64)	0.0055(0.0008)	1.10E-11	0.0056(0.0004)	0.93(0.91,0.96)	2.35E-07	0.0053(0.0015)	5.58E-04
rs10491967	12	3,368,093	<i>TSPAN9</i>	A/G(0.10)	-0.0095(0.0013)	5.18E-14	-0.0095(0.0013)	1.06(1.02,1.11)	6.54E-03	-0.0024(0.0024)	3.24E-01
rs7956634	12	15,321,194	<i>PTPRO</i>	T/C(0.81)	-0.0068(0.0010)	7.17E-12	-0.0066(0.0011)	1.03(1.00,1.06)	8.94E-02	-0.0059(0.0019)	2.17E-03
rs1106766	12	57,809,456	<i>INHBC</i>	T/C(0.22)	0.0061(0.0010)	2.41E-09	0.0060(0.0011)	0.98(0.95,1.02)	3.07E-01	0.0048(0.0019)	1.03E-02
rs164748	16	89,708,292	<i>DPEP1</i>	C/G(0.53)	0.0042(0.0008)	1.63E-07	0.0040(0.0010)	0.99(0.96,1.01)	3.27E-01	0.0046(0.0015)	1.80E-03
rs8091180	18	77,164,243	<i>NFATC1</i>	A/G(0.56)	-0.0053(0.0009)	5.57E-09	-0.0053(0.0009)	1.03(1.00,1.06)	3.57E-02	-0.0050(0.0020)	1.19E-02
rs11666497	19	38,464,262	<i>SIPA1L3</i>	T/C(0.18)	-0.0058(0.0011)	4.25E-08	-0.0061(0.0013)	1.00(0.97,1.04)	9.70E-01	-0.0027(0.0019)	1.70E-01
rs6088580	20	33,285,053	<i>TP53INP2</i>	C/G(0.47)	-0.0049(0.0008)	1.79E-09	-0.0049(0.0008)	1.04(1.01,1.07)	5.25E-03	0.0000(0.0015)	9.92E-01
rs17216707	20	52,732,362	<i>BCAS1</i>	T/C(0.79)	-0.0077(0.0010)	8.83E-15	-0.0076(0.0011)	1.05(1.01,1.09)	8.18E-03	-0.0083(0.0020)	3.25E-05

* Conventional locus name based on relevant genes in the region as identified by bioinformatic investigation (Suppl. Tab. 12) or closest gene. A complete overview of the genes in each locus is given in the regional association plots (Suppl. Fig. 4)

Supplementary Table 5. SNP associations for novel and known loci stratified by diabetes status.

SNP ID (effect allele)	Locus name	No Diabetes			Diabetes			P-value for difference*
		N	beta (SE)	P-value	N	beta (SE)	P-value	
Novel loci: sample size based on the combined (discovery and replication) sample								
rs3850625 (A)	CACNA1S	153,107	0.0083 (0.0013)	6.82E-11	16,275	0.0038 (0.0048)	4.34E-01	0.27
rs2712184 (A)	IGFBP5	153,854	-0.0053 (0.0004)	1.33E-10	16,463	-0.0034 (0.0031)	2.70E-01	0.22
rs9682041 (T)	SKIL	150,911	-0.0068 (0.0012)	2.58E-08	16,161	0.0006 (0.0045)	8.86E-01	0.20
rs10513801 (T)	ETV5	154,774	0.0072 (0.0012)	1.03E-09	16,470	0.0022 (0.0045)	6.19E-01	0.28
rs10994860 (T)	A1CF	154,644	0.0077 (0.0011)	1.07E-12	16,451	0.0031 (0.0041)	4.57E-01	0.27
rs163160 (A)	KCNQ1	154,684	0.0065 (0.0011)	2.26E-09	16,457	0.0055 (0.0040)	1.68E-01	0.81
rs164748 (C)	DPEP1	154,497	0.0046 (0.0004)	1.95E-08	16,416	0.0026 (0.0031)	4.07E-01	0.52
rs8091180 (A)	NFATC1	153,715	-0.0060 (0.0010)	1.28E-09	13,764	0.0030 (0.0043)	4.85E-01	0.04
rs12136063 (A)	SYPL2	152,998	0.0043 (0.0009)	2.25E-06	16,253	0.0013 (0.0034)	7.02E-01	0.38
rs2802729 (A)	SDCCAG8	152,501	-0.0042 (0.0009)	2.65E-06	16,156	-0.0050 (0.0033)	1.24E-01	0.81
rs4667594 (A)	LRP2	154,570	-0.0042 (0.0008)	2.09E-07	16,449	-0.0034 (0.0031)	2.67E-01	0.80
rs6795744 (A)	WNT7A	154,716	0.0053 (0.0011)	1.23E-06	16,462	0.0070 (0.0043)	1.02E-01	0.70
rs228611 (A)	NFKB1	154,683	-0.0056 (0.0008)	7.39E-12	16,454	-0.0077 (0.0030)	1.07E-02	0.49
rs7759001 (A)	ZNF204	154,713	-0.0053 (0.0009)	1.02E-08	16,458	-0.0000 (0.0036)	9.92E-01	0.15
rs10277115 (A)	UNCX	141,932	0.0085 (0.0013)	4.97E-11	13,565	0.0156 (0.0053)	3.30E-03	0.19
rs3750082 (A)	KBTD2	148,429	0.0042 (0.0009)	3.82E-06	15,754	0.0050 (0.0033)	1.32E-01	0.81
rs6459680 (T)	RNF32	154,586	-0.0057 (0.0009)	4.64E-10	16,454	-0.0043 (0.0035)	2.25E-01	0.70
rs4014195 (C)	AP5B1	154,634	0.0050 (0.0008)	1.02E-09	16,452	0.0116 (0.0032)	2.52E-04	0.04
rs10491967 (A)	TSPAN9	154,881	-0.0089 (0.0013)	3.18E-11	16,477	-0.0204 (0.0050)	3.80E-05	0.02
rs7956634 (T)	PTPRO	154,677	-0.0065 (0.0010)	1.11E-10	16,460	-0.0076 (0.0038)	4.82E-02	0.78
rs1106766 (T)	INHBC	138,058	0.0063 (0.0010)	1.75E-09	14,990	0.0016 (0.0040)	6.89E-01	0.25
rs11666497 (T)	SIPA1L3	148,770	-0.0058 (0.0011)	6.04E-08	15,831	-0.0096 (0.0041)	2.01E-02	0.37
rs6088580 (C)	TP53INP2	150,326	-0.0050 (0.0008)	1.51E-09	15,387	-0.0010 (0.0031)	7.48E-01	0.21
rs17216707 (T)	BCAS1	153,362	-0.0075 (0.0011)	2.25E-12	15,977	-0.0063 (0.0039)	1.06E-01	0.76
Known loci: sample size based on the discovery sample								
rs12917707 (T)	UMOD	118,365	0.0152 (0.0012)	4.68E-36	11,522	0.0266 (0.0048)	2.48E-08	0.02
rs1260326 (T)	GCKR	118,415	0.0065 (0.0009)	1.86E-12	11,525	0.0112 (0.0037)	2.21E-03	0.21
rs6465825 (T)	SLC22A2	118,448	0.0061 (0.0009)	4.80E-11	11,529	0.0089 (0.0036)	1.44E-02	0.45
rs2279463 (A)	VEGFA	118,368	0.0119 (0.0014)	6.98E-18	11,521	0.0062 (0.0053)	2.43E-01	0.30
rs12124078 (A)	CASP9	116,718	0.0058 (0.0010)	4.54E-09	11,315	0.0039 (0.0039)	3.18E-01	0.63
rs848490 (C)	TMEM60	112,566	0.0074 (0.0010)	1.35E-12	10,897	0.0033 (0.0041)	4.30E-01	0.33
rs2928148 (A)	INO80	118,368	0.0050 (0.0009)	2.82E-08	11,521	0.0026 (0.0035)	4.73E-01	0.50
rs9895661 (T)	BCAS3	118,188	0.0121 (0.0012)	2.80E-22	11,520	-0.0006 (0.0048)	9.00E-01	0.01
rs347685 (A)	TFDP2	118,374	-0.0077 (0.0010)	1.70E-14	11,522	-0.0021 (0.0040)	5.93E-01	0.17
rs7208487 (T)	CDK12	118,453	-0.0091 (0.0012)	8.60E-14	11,526	-0.0024 (0.0048)	6.20E-01	0.17

Supplementary Table 5 (continued).

SNP ID (effect allele)	Locus name	No Diabetes			Diabetes			P-value for difference*
		N	beta (SE)	P-value	N	beta (SE)	P-value	
Known loci: sample size based on the discovery sample								
rs10774021 (T)	SLC6A13	118,448	-0.0067 (0.0009)	1.48E-12	11,526	-0.0031 (0.0038)	4.13E-01	0.35
rs2453580 (T)	SLC47A1	117,386	0.0073 (0.0010)	2.56E-13	11,474	-0.0035 (0.0040)	3.93E-01	0.01
rs4744712 (A)	PIP5K1B	118,374	-0.0072 (0.0009)	7.45E-15	11,518	-0.0053 (0.0037)	1.48E-01	0.62
rs12460876 (T)	SLC7A9	118,361	-0.0065 (0.0009)	2.86E-12	11,521	-0.0062 (0.0037)	9.16E-02	0.94
rs881858 (A)	SLC34A1	118,324	-0.0081 (0.0011)	4.03E-14	11,521	-0.0063 (0.0042)	1.35E-01	0.68
rs10109414 (T)	STC1	118,342	-0.0075 (0.0009)	3.53E-16	11,521	-0.0066 (0.0036)	6.75E-02	0.81
rs6431731 (T)	DDX1	109,308	-0.0124 (0.0025)	4.70E-07	10,419	-0.0070 (0.0100)	4.89E-01	0.60
rs626277 (A)	DACH1	118,450	-0.0049 (0.0009)	1.44E-07	11,528	-0.0072 (0.0036)	4.53E-02	0.53
rs17319721 (A)	SHROOM3	118,359	-0.0113 (0.0009)	5.08E-35	11,522	-0.0075 (0.0036)	3.64E-02	0.30
rs13538 (A)	ALMS1	115,154	-0.0092 (0.0011)	2.79E-16	11,135	-0.0077 (0.0045)	8.45E-02	0.74
rs11959928 (A)	DAB2	118,359	-0.0082 (0.0009)	7.14E-19	11,521	-0.0078 (0.0036)	3.33E-02	0.91
rs2453533 (A)	GATM	118,374	-0.0125 (0.0009)	4.83E-41	11,520	-0.0079 (0.0036)	3.16E-02	0.21
rs7805747 (A)	PRKAG2	116,602	-0.0136 (0.0012)	1.53E-29	11,378	-0.0085 (0.0048)	7.37E-02	0.30
rs1394125 (A)	UBE2Q2	118,404	-0.0072 (0.0010)	5.83E-13	11,525	-0.0089 (0.0041)	2.92E-02	0.69
rs3925584 (T)	MPPED2	118,345	-0.0074 (0.0009)	5.35E-16	11,521	-0.0100 (0.0035)	5.00E-03	0.47
rs267734 (T)	LASS2	116,718	-0.0079 (0.0011)	2.81E-12	11,315	-0.0110 (0.0045)	1.39E-02	0.50
rs7422339 (A)	CPS1	115,499	-0.0106 (0.0011)	3.25E-22	11,422	-0.0137 (0.0043)	1.61E-03	0.48
rs10794720 (T)	WDR37	118,433	-0.0093 (0.0017)	4.95E-08	11,528	-0.0145 (0.0067)	3.06E-02	0.45
rs491567 (A)	WDR72	118,374	-0.0078 (0.0011)	1.45E-12	11,521	-0.0167 (0.0043)	1.25E-04	0.04

* P-value of a two-sample *t* test for correlated data (see Methods: “Associations Stratified by Diabetes and Hypertension Status”).

Supplementary Table 6. Association between replicated SNPs and additional kidney related phenotypes.*

			Myocardial Infarction		Left Ventricular Mass		Incident Heart Failure		DBP		SBP		UACR		Fasting glucose	
SNPID	Locus name	All. #	OR (95%CI)	P-value	beta (SE)	P-value	beta (SE)	P-value	beta (SE)	P-value	beta (SE)	P-value	beta(SE E)	P-value	beta (SE)	P-value
rs3850625	<i>CACNA1S</i>	A/G	1.03 (0.99,1.08)	0.15	0.390 (0.680)	0.57	-0.034 (0.044)	0.44	0.003 (0.094)	0.97	-0.068 (0.147)	0.64	0.017 (0.011)	0.11	0.012 (0.005)	0.013
rs2712184	<i>IGFBP5</i>	A/C	1.00 (0.97,1.03)	0.81	-0.183 (0.449)	0.68	-0.051 (0.034)	0.14	-0.087 (0.062)	0.16	-0.016 (0.098)	0.87	-0.001 (0.006)	0.88	-0.001 (0.003)	0.77
rs9682041	<i>SKIL</i>	T/C	0.99 (0.95,1.04)	0.79	-0.036 (0.639)	0.96	-0.003 (0.043)	0.95	-0.026 (0.093)	0.78	-0.092 (0.147)	0.53	-0.015 (0.009)	0.11	-0.006 (0.005)	0.18
rs10513801	<i>ETV5</i>	T/G	0.97 (0.94,1.01)	0.18	0.255 (0.649)	0.69	0.019 (0.044)	0.67	-0.021 (0.091)	0.82	0.004 (0.145)	0.98	0.027 (0.011)	0.014	0.002 (0.005)	0.71
rs10994860	<i>A1CF</i>	T/C	0.99 (0.96,1.03)	0.56	0.294 (0.595)	0.62	-0.081 (0.041)	0.05	-0.064 (0.081)	0.43	-0.162 (0.127)	0.20	-0.006 (0.009)	0.55	0.004 (0.004)	0.28
rs163160	<i>KCNQ1</i>	A/G	0.98 (0.95,1.02)	0.39	0.360 (0.577)	0.53	-0.063 (0.038)	0.10	-0.040 (0.082)	0.62	0.032 (0.129)	0.80	0.011 (0.010)	0.26	0.001 (0.004)	0.86
rs164748	<i>DPEP1</i>	C/G	0.99 (0.97,1.02)	0.65	0.668 (0.439)	0.13	0.062 (0.029)	0.04	0.173 (0.063)	0.01	0.361 (0.099)	0.00026 ^s	0.021 (0.006)	0.0004	-0.003 (0.003)	0.30
rs8091180	<i>NFATC1</i>	A/G	NA	NA	0.307 (0.816)	0.71	0.056 (0.044)	0.21	0.122 (0.103)	0.23	0.181 (0.162)	0.27	-0.018 (0.015)	0.21	0.006 (0.005)	0.19
rs12136063	<i>SYPL2</i>	A/G	1.03 (1.00,1.06)	0.08	-0.192 (0.478)	0.69	0.023 (0.032)	0.47	0.032 (0.066)	0.62	-0.033 (0.104)	0.75	0.010 (0.008)	0.19	-0.007 (0.003)	0.05
rs2802729	<i>SDCCAG8</i>	A/C	NA	NA	-0.127 (0.470)	0.79	-0.005 (0.032)	0.88	0.147 (0.067)	0.03	0.239 (0.105)	0.02	0.008 (0.007)	0.23	0.002 (0.003)	0.53
rs4667594	<i>LRP2</i>	A/T	0.99 (0.97,1.02)	0.59	0.940 (0.440)	0.03	0.019 (0.029)	0.51	0.098 (0.062)	0.11	0.152 (0.097)	0.12	-0.008 (0.006)	0.18	0.000 (0.003)	0.89
rs6795744	<i>WNT7A</i>	A/G	0.98 (0.94,1.02)	0.27	0.298 (0.643)	0.64	-0.071 (0.045)	0.12	-0.022 (0.088)	0.80	-0.110 (0.139)	0.43	-0.005 (0.009)	0.57	0.005 (0.004)	0.30
rs228611	<i>NFKB1</i>	A/G	0.98 (0.95,1.01)	0.18	0.165 (0.440)	0.71	0.008 (0.028)	0.78	0.090 (0.061)	0.14	0.003 (0.097)	0.98	0.008 (0.006)	0.20	0.011 (0.003)	0.00026 ^s
rs7759001	<i>ZNF204</i>	A/G	0.99 (0.96,1.02)	0.51	0.172 (0.524)	0.74	-0.031 (0.034)	0.36	-0.022 (0.071)	0.75	-0.085 (0.112)	0.45	0.002 (0.007)	0.83	0.006 (0.004)	0.10
rs10277115	<i>UNCX</i>	A/T	NA	NA	-1.058 (0.777)	0.17	0.012 (0.052)	0.82	-0.099 (0.141)	0.48	-0.232 (0.219)	0.29	0.013 (0.012)	0.27	-0.009 (0.006)	0.15
rs3750082	<i>KBTBD2</i>	A/T	0.99 (0.96,1.02)	0.48	0.641 (0.470)	0.17	-0.005 (0.031)	0.86	-0.024 (0.066)	0.72	0.094 (0.104)	0.37	0.019 (0.006)	0.0036	-0.001 (0.003)	0.88
rs6459680	<i>RNF32</i>	T/G	0.98 (0.95,1.01)	0.14	-0.234 (0.514)	0.65	-0.010 (0.034)	0.76	-0.155 (0.071)	0.03	-0.184 (0.112)	0.10	-0.001 (0.008)	0.90	0.002 (0.004)	0.60
rs4014195	<i>AP5B1</i>	C/G	0.98 (0.95,1.01)	0.20	-0.825 (0.458)	0.07	-0.052 (0.035)	0.14	-0.079 (0.064)	0.22	-0.176 (0.101)	0.08	0.007 (0.006)	0.29	0.001 (0.003)	0.70

Supplementary Table 6 (continued).

			Myocardial Infarction		Left Ventricular Mass		Incident Heart Failure		DBP		SBP		UACR		Fasting glucose	
SNPID	Locus name	All. #	OR (95%CI)	P-value	beta (SE)	P-value	beta (SE)	P-value	beta (SE)	P-value	beta (SE)	P-value	beta(SE E)	P-value	beta (SE)	P-value
rs10491967	<i>TSPAN9</i>	A/G	0.95 (0.91,1.00)	0.04	0.087 (0.696)	0.90	-0.033 (0.047)	0.48	0.003 (0.101)	0.98	-0.205 (0.160)	0.20	0.002 (0.011)	0.86	-0.003 (0.005)	0.60
rs7956634	<i>PTPRO</i>	T/C	1.01 (0.98,1.05)	0.49	0.528 (0.564)	0.35	0.010 (0.036)	0.79	0.047 (0.083)	0.57	0.113 (0.130)	0.39	-0.017 (0.008)	0.03	0.006 (0.004)	0.17
rs1106766	<i>INHBC</i>	T/C	0.99 (0.96,1.02)	0.56	-0.322 (0.584)	0.58	0.027 (0.037)	0.46	0.069 (0.079)	0.39	0.093 (0.125)	0.46	0.011 (0.009)	0.22	-0.001 (0.004)	0.82
rs11666497	<i>SIPA1L3</i>	T/C	1.02 (0.98,1.05)	0.42	0.624 (0.590)	0.29	0.058 (0.041)	0.15	-0.056 (0.081)	0.49	-0.034 (0.127)	0.79	0.000 (0.008)	1.00	0.002 (0.004)	0.72
rs6088580	<i>TP53INP2</i>	C/G	1.02 (0.99,1.05)	0.14	-0.864 (0.456)	0.06	0.014 (0.028)	0.63	-0.126 (0.061)	0.04	-0.066 (0.097)	0.49	0.010 (0.006)	0.10	0.010 (0.003)	0.0016
rs17216707	<i>BCAS1</i>	T/C	1.02 (0.98,1.05)	0.40	0.238 (0.605)	0.69	0.038 (0.041)	0.35	0.136 (0.084)	0.11	0.195 (0.132)	0.14	-0.000 (0.009)	0.98	0.004 (0.004)	0.36

* **Myocardial infarction:** results from CARDIoGRAM meta-analysis⁸ of up to 14 studies, for a total sample size comprised between 72,649 and 83,231, except for SNPs rs4014195 (N=61,259) and rs2712184 (N=61,275). SNPs not reported did not pass CARDIoGRAM internal quality control checks and were not assessed. **Left Ventricular Mass:** per-allele effect on left ventricular mass in grams, from a meta-analysis of five community-based studies within the EchoGen consortium, totaling 12,612 European ancestry individuals.⁹ **Incident heart failure:** CHARGE consortium meta-analysis on incident heart failure:¹⁰ four studies on European ancestry individuals totaling a sample size of 20,926, except for SNPs rs2712184 and rs4014195 (N=13,282). **DBP** and **SBP:** diastolic and systolic blood pressure results from the ICBP consortium:¹¹ meta-analysis of up to 43 studies, for a total sample size of 53,302 to 69,671 samples, except for SNPs rs10277115 (27 studies; N=17,644) and rs8091180 (21 studies; N=23,456). **UACR:** urinary albumin-to-creatinine ratio, results from the CKDGen consortium (personal communication): meta-analysis of up to 29 studies, for a total sample size of 39,130 to 54,450. **Fasting glucose:** per-allele effect on fasting glucose (mmol/L) not adjusted for BMI in 29 studies on up 58,074 non-diabetic participants of European ancestry.¹²

All.: first is the effect and second is the non-effect allele, respectively.

§ P-values that are significant at a Bonferroni corrected level of 0.0003, corresponding to 0.05 over 165 tests.

Supplementary Table 7. NHGRI GWAS catalog query of novel loci associated with different traits at a genome-wide significant level.

Locus name	Index SNP	Published SNP	Chr	Position (Build 37)	Region	LD (r^2)*	Disease or Quantitative Trait ^{REF}	P-value
<i>SDCCAG8</i>	rs2802729	rs6703335	1	243,608,967	1q43	0.44	Schizophrenia ¹³	5.00E-08
<i>ETV5</i>	rs10513801	rs1516725	3	185,824,004	3q27.2	1.00	Body mass index ¹⁴	4.00E-08
		rs1516725	3	185,824,004	3q27.2	1.00	Obesity ¹⁴	3.00E-09
		rs7647305	3	185,834,290	3q27.2	0.39	Body mass index ¹⁵	7.00E-11
		rs7647305	3	185,834,290	3q27.2	0.39	Weight ¹⁵	4.00E-09
		rs9816226	3	185,834,499	3q27.2	0.49	Obesity ¹⁴	2.00E-13
		rs9816226	3	185,834,499	3q27.2	0.49	Obesity ¹⁴	2.00E-14
		rs9816226	3	185,834,499	3q27.2	0.49	Body mass index ¹⁶	2.00E-18
<i>NFKB1</i>	rs228611	rs3774959	4	103,511,114	4q24	0.29	Ulcerative colitis ¹⁷	4.00E-12
		rs7665090	4	103,551,603	4q24	0.87	Primary biliary cirrhosis ¹⁸	4.00E-12
<i>UNCX</i>	rs10277115	rs10275044	7	1,273,845	7p22.3	0.73	Blood Urea Nitrogen ¹⁹	4.00E-09
		rs10277115	7	1,285,195	7p22.3	1.00	eGRFcrea ¹⁹	1.00E-10
		rs10277115	7	1,285,195	7p22.3	1.00	Serum creatinine ¹⁹	5.00E-11
<i>A1CF</i>	rs10994860	rs10821905	10	52,646,093	10q11.23	1.00	Urate levels ²⁰	7.00E-17
<i>AP5B1</i>	rs4014195	rs479844	11	65,551,957	11q13.1	0.37	Atopic dermatitis ²¹	1.00E-13
		rs642803	11	65,560,620	11q13.1	0.40	Urate levels ²⁰	3.00E-13
<i>INHBC</i>	rs1106766	rs11613352	12	57,792,580	12q13.3	1.00	HDL cholesterol ²²	2.00E-08
		rs11613352	12	57,792,580	12q13.3	1.00	Triglycerides ²²	4.00E-10
		rs1106766	12	57,809,456	12q13.3	1.00	Urate levels ²³	2.00E-11
		rs3741414	12	57,844,049	12q13.3	0.97	Urate levels ²⁰	2.00E-25
<i>DPEP1</i>	rs164748	rs154657	16	89,708,096	16q24.3	1.00	Homocysteine levels ²⁴	2.00E-43
		rs258322	16	89,755,903	16q24.3	0.30	Melanoma ²⁵	3.00E-27
		rs258322	16	89,755,903	16q24.3	0.30	Melanoma ²⁶	3.00E-27
		rs258322	16	89,755,903	16q24.3	0.30	Black vs. red hair color ²⁷	2.00E-23
		rs12921383	16	89,859,753	16q24.3	0.25	Homocysteine levels ²⁴	8.00E-11
		rs1805007	16	89,986,117	16q24.3	0.25	Hair color ²⁸	3.00E-09
		rs1805007	16	89,986,117	16q24.3	0.25	Non-melanoma skin cancer ²⁸	3.00E-10
		rs1805007	16	89,986,117	16q24.3	0.25	Sunburns ²⁸	2.00E-19
		rs1805007	16	89,986,117	16q24.3	0.25	Tanning ²⁸	1.00E-65
		rs1805007	16	89,986,117	16q24.3	0.25	Basal cell carcinoma ²⁹	4.00E-17
		rs1805007	16	89,986,117	16q24.3	0.25	Blond vs. brown hair color ³⁰	2.00E-13
		rs1805007	16	89,986,117	16q24.3	0.25	Freckles ³⁰	1.00E-96
		rs1805007	16	89,986,117	16q24.3	0.25	Red vs non-red hair color ³⁰	2.00E-142
		rs1805007	16	89,986,117	16q24.3	0.25	Skin sensitivity to sun ³⁰	2.00E-55
<i>TP53INP2</i>	rs6088580	rs2284378	20	32,588,095	20q11.22	0.25	Breast cancer ³¹	1.00E-08
		rs4911414	20	32,729,444	20q11.22	0.25	Tanning ²⁸	4.00E-09
		rs4911414	20	32,729,444	20q11.22	0.25	Burning and freckling ³²	6.00E-37
		rs4911414	20	32,729,444	20q11.22	0.25	Freckles ³²	8.00E-29
		rs4911414	20	32,729,444	20q11.22	0.25	Red vs. non-red hair color ³²	3.00E-09
		rs4911414	20	32,729,444	20q11.22	0.25	Skin sensitivity to sun ³²	2.00E-24
		rs910873	20	33,171,772	20q11.22	0.21	Melanoma ³³	1.00E-15
		rs8114671	20	33,789,142	20q11.22	0.47	Height ¹⁴	1.00E-15
		rs6088765	20	33,799,280	20q11.22	0.34	Ulcerative colitis ¹⁷	2.00E-08

* LD between published and index SNPs.

Supplementary Table 8. Study sample characteristics, African ancestry meta-analysis.

Study	Sample Size	European Ancestry % - change to median with 25/75th percentiles	Women %	Mean age (years)	Mean eGFR_{crea} (ml/min/1.73 m²)	CKD %
ARIC	2786	15.3 (10.7, 22.1)	63.1	53.3	100	3.7
CARDIA	821	16.7 (12.2, 23.2)	61.1	39.4	111	0.9
CHS	728	20.6 (12.4, 32.7)	62.8	72.9	81	18.4
JHS	2135	15.7 (11.8, 21.1)	60.8	50	101	4.2
MESA	1640	18.8 (11.5, 29.7)	54.8	62.2	87	8.6
GENOA	1217	12.6 (7.2, 18.9)	71.7	63.2	88	13.0
HANDLS	989	16.1 (11.2, 22.0)	55.0	48.4	121	5.3
Health ABC	1139	22.4 (12.2, 32.6)	57.2	73.4	76	17.1
HUFS	1013	19.7 (14.3, 27.0)	58.8	48.3	104	4.9
IPM	712	12.5 (7.1, 19.5)	60.0	59.7	76	36.4
SIGNET-Sea Islands	1275	7.0 (4.3, 11.7)	77.1	53.7	106	9.6
SIGNET-REGARDS	2385	14.8 (9.0, 22.9)	63.7	63	100	10.0

Abbreviations: eGFR_{crea}: estimated glomerular filtration rate by serum creatinine, CKD: chronic kidney disease.

Supplementary Table 9. Evaluation of replicated SNPs among individuals of Asian (AGEN) and the African Ancestry Renal meta-analysis.

Locus name	SNPID	Eff. All.	Non Eff. All.	AGEN Consortium ¹⁹				African Ancestry Renal Meta-Analysis			
				N	Eff. All. Freq.	Beta (SE)	P-value	N	Ref. all freq.	Beta (SE)	P-value
<i>SYPL2</i>	rs12136063	A	G	36,057	0.90	-0.0004 (0.0021)	0.85	16,349	0.32	0.0049 (0.0036)	0.16
<i>CACNA1S</i>	rs3850625	A	G	41,963	0.09	-0.0012 (0.0019)	0.53	11,194	0.02	0.0127 (0.0135)	0.35
<i>SDCCAG8</i>	rs2802729	A	C	42,296	0.29	-0.0007 (0.0009)	0.42	15,461	0.47	-0.0007 (0.0036)	0.85
<i>LRP2</i>	rs4667594	A	T	40,415	0.13	-0.0015 (0.0013)	0.24	15,417	0.61	-0.0016 (0.0034)	0.63
<i>IGFBP5</i>	rs2712184	A	C	40,415	0.51	-0.0025 (0.0008)	0.0024	16,210	0.48	-0.0023 (0.0034)	0.50
<i>WNT7A</i>	rs6795744	A	G	42,296	0.11	-0.0001 (0.0016)	0.94	16,420	0.21	0.0032 (0.0043)	0.46
<i>SKIL</i>	rs9682041	T	C	42,296	0.91	-0.0047 (0.0018)	0.0074	15,460	0.75	-0.0005 (0.0041)	0.91
<i>ETV5</i>	rs10513801	T	G	37,250	0.93	0.0006 (0.0026)	0.82	14,243	0.95	-0.0006 (0.0100)	0.95
<i>NFKB1</i>	rs228611	A	G	42,296	0.52	-0.0001 (0.0008)	0.88	16,450	0.24	0.0042 (0.0040)	0.29
<i>ZNF204</i>	rs7759001	A	G	40,415	0.62	-0.0005 (0.0008)	0.56	16,435	0.87	0.0044 (0.0050)	0.38
<i>UNCX</i>	rs10277115	A	T	40,415	0.65	0.0066 (0.0011)	7.30E-10	11,801	0.70	0.0059 (0.0060)	0.32
<i>KBTBD2</i>	rs3750082	A	T	39,440	0.32	0.0012 (0.0009)	0.19	15,461	0.67	0.0070 (0.0039)	0.07
<i>RNF32</i>	rs6459680	T	G	40,415	0.62	-0.0005 (0.0009)	0.56	16,435	0.54	0.0049 (0.0034)	0.15
<i>A1CF</i>	rs10994860	T	C	41,963	0.10	0.0014 (0.0017)	0.42	15,461	0.22	0.0060 (0.0045)	0.18
<i>KCNQ1</i>	rs163160	A	G	40,415	0.83	0.0022 (0.0011)	0.04	16,467	0.93	0.0053 (0.0067)	0.43
<i>AP5B1</i>	rs4014195	G	C	40,415	0.20	-0.0040 (0.0010)	0.00012	15,297	0.22	-0.0014 (0.0042)	0.74
<i>TSPAN9</i>	rs10491967	A	G	40,415	0.45	-0.0013 (0.0008)	0.13	15,460	0.40	0.0072 (0.0043)	0.09
<i>PTPRO</i>	rs7956634	T	C	40,415	0.68	-0.0020 (0.0009)	0.03	16,403	0.50	-0.0082 (0.0033)	0.01
<i>INHBC</i>	rs1106766	T	C	42,296	0.11	0.0003 (0.0015)	0.85	16,415	0.10	-0.0008 (0.0057)	0.89
<i>DPEP1</i>	rs164748	G	C	39,164	0.08	0.0016 (0.0024)	0.49	16,471	0.11	0.0025 (0.0055)	0.65
<i>NFATC1</i>	rs8091180	A	G	40,415	0.83	-0.0004 (0.0011)	0.69	3,838	0.20	-0.0021 (0.0101)	0.84
<i>SIPA1L3</i>	rs11666497	T	C	40,415	0.12	-0.0027 (0.0013)	0.03	16,422	0.07	0.0023 (0.0064)	0.72
<i>TP53INP2</i>	rs6088580	G	C	36,275	0.64	0.0007 (0.0009)	0.45	NA	NA	NA	NA
<i>BCAS1</i>	rs17216707	T	C	35,492	0.86	-0.0012 (0.0018)	0.52	15,399	0.94	-0.0049 (0.0073)	0.50

Supplementary Table 10. Transethnic meta-analysis of CKDGen and the African Ancestry Renal Meta-Analysis.*

SNPID	Locus name	Chr.	Effect Allele	Non-Effect Allele	log10BF#	Posterior Probability
rs17216707	<i>BCAS1</i>	20	T	C	11.63	0.038
rs4014195	<i>AP5B1</i>	11	C	G	9.66	0.025
rs10994860	<i>A1CF</i>	10	T	C	9.52	0.025
rs7956634	<i>PTPRO</i>	12	T	C	9.07	0.015
rs10277115	<i>UNCX</i>	7	A	T	8.84	0.023
rs228611	<i>NFKB1</i>	4	A	G	7.65	0.07
rs6088580	<i>TP53INP2</i>	20	C	G	7.44	0.021
rs163160	<i>KCNQ1</i>	11	A	G	7.38	0.026
rs3850625	<i>CACNA1S</i>	1	A	G	7.34	0.046
rs6795744	<i>WNT7A</i>	3	A	G	7.04	0.046
rs10491967	<i>TSPAN9</i>	12	A	G	6.93	0.725
rs2712184	<i>IGFBP5</i>	2	A	C	6.72	0.021
rs3750082	<i>KBTBD2</i>	7	A	T	6.65	0.01
rs6459680	<i>RNF32</i>	7	T	G	6.42	0.701
rs12136063	<i>SYPL2</i>	1	A	G	6.18	0.012
rs2802729	<i>SDCCAG8</i>	1	A	C	6.00	0.021
rs10513801	<i>ETV5</i>	3	T	G	5.88	0.03
rs164748	<i>DPEP1</i>	16	C	G	5.83	0.045
rs4667594	<i>LRP2</i>	2	A	T	5.72	0.027
rs11666497	<i>SIPA1L3</i>	19	T	C	5.72	0.031
rs1106766	<i>INHBC</i>	12	T	C	5.56	0.053
rs7759001	<i>ZNF204</i>	6	A	G	5.23	0.051
rs9682041	<i>SKIL</i>	3	T	C	5.04	0.032
rs8091180	<i>NFATC1</i>	18	A	G	4.87	0.044

*The analysis was performed using the MANTRA (Meta-Analysis of Trans-ethnic Association studies) software.³⁴ The posterior probability is a measure of heterogeneity of allelic effects across the individual studies.

#Log₁₀ Bayes Factor

Supplementary Table 11. SNP associations with transcript expression.*

Locus name; index SNP	eSNP									Best eSNP			
	eSNP rsID	Dist. index SNP	r ²	eQTL tissue ^{REF}	P-value	Chr	position	Probe*	Transcript**	Best eSNP rsID	P-value	r ² to eSNP	r ² to index SNP
SYPL2 rs12136063	rs12136063	0	1.00	Prefrontal cortex - all samples ³⁵	1.3E-05	1	109,815,693	10031920561	SYPL2	rs12136063	1.3E-05	Same	Same
	rs10494040	1,562	1.00	SubCutAdipose ³⁶	6.5E-50	1	109,817,255	10031920561	SYPL2	rs10494040	6.5E-50	Same	0.96
	rs10494040	1,562	1.00	Liver ³⁶	5.4E-47	1	109,817,255	10031920561	SYPL2	rs10494040	5.4E-47	Same	0.96
	rs10857787	3,881	1.00	Liver ³⁷	2.9E-35	1	109,811,812		SYPL2	rs10857787	2.9E-35	Same	0.96
	rs4970767	1,013	1.00	Liver (UChicago) ³⁸	8.9E-16	1	109,816,706	A_23_P317200	ATXN7L2	rs4970767	8.9E-16	Same	0.96
	rs10494040	1,562	1.00	OmentalAdipose ³⁶	9.4E-13	1	109,817,255	10031920561	SYPL2	rs10494040	9.4E-13	Same	0.96
	rs10494040	1,562	1.00	Liver ³⁹	1.1E-12	1	109,817,255		SYPL2	rs10494040	1.1E-12	Same	0.96
	rs4970767	1,013	1.00	Liver (UWash) ³⁸	3.3E-02	1	109,816,706	5360451	ATXN7L2	rs4970767	3.3E-02	Same	0.96
	rs4970729	34,968	0.92	Liver ³⁶	2.3E-24	1	109,780,725	10025910902	PSMA5	rs4970729	2.3E-24	Same	0.89
	rs12073497	39,720	0.92	Liver ³⁶	6.2E-09	1	109,775,973	10023805980	Contig42599 RC	rs12073497	6.2E-09	Same	0.89
	rs12073497	39,720	0.92	Visual cortex - all samples ³⁵	3.1E-06	1	109,775,973	10031920561	SYPL2	rs12073497	3.1E-06	Same	0.89
	rs4970729	34,968	0.92	Cerebellum - all samples ³⁵	1.1E-05	1	109,780,725	10025932473	AMIGO1	rs4970729	1.1E-05	Same	0.89
SDCCAG8 rs2802729	rs2781553	12,819	0.85	Liver ³⁶	4.8E-05	1	109,828,512	10025909878	PRPF38B	rs2781553	4.8E-05	Same	0.85
	rs2802723	3,451	1.00	Periph artery plaque [‡]	4.1E-07	1	241,564,935	100142973_TGI_at	SDCCAG8	rs2802723	4.1E-07	Same	1.00
	rs2490395	42,841	0.83	Blood ⁴⁰	9.2E-06	1	241,525,545	460458	SDCCAG8	rs2490395	9.2E-06	Same	0.83
NFKB1 rs228611	rs2484639	39,396	0.81	Visual cortex - all samples ³⁵	3.2E-07	1	241,528,990	10025912019	SDCCAG8	rs2484639	3.2E-07	Same	0.80
	rs228611	0	1.00	LCL MuTHER ⁴¹	1.6E-19	4	103,780,757	ILMN_1800733	MANBA	rs228611	1.6E-19	Same	Same
	rs228611	0	1.00	Prefrontal cortex (Huntington's) ³⁵	1.4E-08	4	103,780,757	10025907439	MANBA	rs228611	1.4E-08	Same	Same
	rs228611	0	1.00	Lymphocytes ⁴²	2.0E-08	4	103,780,757		MANBA	rs228611	2.0E-08	Same	Same
	rs228611	0	1.00	Visual cortex (Alzheimer's) ³⁵	5.6E-05	4	103,780,757	10025907439	MANBA	rs228611	5.6E-05	Same	Same
	rs228611	0	1.00	Cerebellum - all samples ³⁵	6.7E-02	4	103,780,757	10025907439	MANBA	rs228611	6.7E-02	Same	Same
	rs909349	5,393	0.97	Monocytes ⁴³	2.5E-121	4	103,775,364		MANBA	rs909349	2.5E-121	Same	0.97
	rs7665090	10,106	0.97	OmentalAdipose ³⁶	6.5E-42	4	103,770,651	10025907439	MANBA	rs7665090	6.5E-42	Same	0.97
	rs228611	0	1.00	Blood ⁴⁰	1.3E-30	4	103,780,757	4230168	MANBA	rs7665090	3.9E-37	0.97	0.97
	rs7665090	10,106	0.97	Prefrontal cortex - all samples ³⁵	2.4E-26	4	103,770,651	10025907439	MANBA	rs7665090	2.4E-26	Same	0.97

Supplementary Table 11 (continued).

Locus name; index SNP	eSNP									Best eSNP			
	eSNP rsID	Dist. index SNP	r ²	eQTL tissue ^{REF}	P-value	Chr	position	Probe*	Transcript**	Best eSNP rsID	P-value	r ² to eSNP	r ² to index SNP
NFKB1 rs228611	rs228614	16,928	0.91	Blood ⁴⁰	2.0E-22	4	103,797,685	4230168	MANBA	rs7665090	3.9E-37	0.87	0.97
	rs2866413	4,632	0.97	Prefrontal cortex - all samples ³⁵	5.8E-17	4	103,776,125	10025907439	MANBA	rs7665090	2.4E-26	1	0.97
	rs2866413	4,632	0.97	Visual cortex - all samples ³⁵	5.8E-08	4	103,776,125	10025907439	MANBA	rs2866413	5.8E-08	Same	0.97
	rs228614	16,928	0.91	Blood ⁴⁴	2.1E-05	4	103,797,685	HSG00228144	MANBA	rs228614	2.1E-05	Same	0.91
	rs404574	21,499	0.90	SubCutAdipose MuTHER ⁴¹	1.7E-14	4	103,802,256	ILMN_1800733	MANBA	rs404574	1.7E-14	Same	0.84
	rs7674640	20,929	0.81	Lung ⁴⁵	<2E-16	4	103,759,828	100150393_TGI_at	CISD2	rs7674640	<2E-16	Same	0.81
ZNF204 rs7759001	rs7759001	0	1.00	Intestine normal ileum ⁴⁶	1.3E-10	6	27,341,409		ZNF391	rs7759001	1.3E-10	Same	Same
	rs7759001	0	1.00	Cerebellum - normal samples ³⁵	5.5E-09	6	27,449,388	10025913649	BC035154	rs7759001	5.5E-09	Same	Same
	rs2143062	12,443	1.00	Cerebellum (Alzheimer's) ³⁵	1.6E-11	6	27,461,831	10025913649	BC035154	rs2143062	1.6E-11	Same	1.00
	rs2143062	12,443	1.00	Prefrontal cortex (Huntington's) ³⁵	1.4E-05	6	27,461,831	10025913649	BC035154	rs2143062	1.4E-05	Same	1.00
	rs9368508	19,561	1.00	Cerebellum (Huntington's) ³⁵	3.4E-06	6	27,429,827	10025913649	BC035154	rs9368508	3.4E-06	Same	0.95
	rs10755644	4,696	1.00	SubCutAdipose ³⁶	7.0E-05	6	27,444,692	10025933640	ZNF391	rs10755644	7.0E-05	Same	0.90
	rs10807021	15,516	0.95	OmentalAdipose ³⁶	1.5E-08	6	27,464,904	10025913649	BC035154	rs10807021	1.5E-08	Same	0.86
	rs4713086	9,037	0.95	Periph artery plaque†	1.9E-07	6	27,458,425	100139132_TGI_at	BC035154	rs4713086	1.9E-07	Same	0.86
	rs980963	19,528	0.86	Cerebellum - all samples ³⁵	2.3E-05	6	27,468,916	10025933640	ZNF391	rs980963	2.3E-05	Same	0.86
	rs10807020	15,493	0.95	Cerebellum - normal samples ³⁵	5.3E-05	6	27,464,881	10025907341	NM_178534	rs10807020	5.3E-05	Same	0.86
KBTBD2 rs3750082	rs10807021	15,516	0.95	Visual cortex - all samples ³⁵	1.3E-03	6	27,464,904	10025908866	ZNF184	rs10807021	1.3E-03	Same	0.86
	rs6462431	15,512	0.93	Blood ⁴⁰	2.3E-38	7	32,901,964	4920372	KBTBD2	rs6462431	2.3E-38	Same	0.89
	rs3750082	0	1.00	LCL asthmatics ⁴⁷	8.1E-17	7	32,886,452	212447_at	KBTBD2	rs7785065	8.1E-17	0.89	0.89
	rs3750082	0	1.00	LCL MuTHER ⁴¹	1.1E-06	7	32,886,452	ILMN_1784540	KBTBD2	rs13230763	8.3E-08	0.80	0.89
	rs2392152	19,378	0.89	Liver ³⁷	2.0E-10	7	32,867,074		HSS00226368	rs2392152	2.0E-10	Same	0.85
	rs4723221	34,280	0.82	Lung ⁴⁵	6.4E-08	7	32,920,732	100132413_TGI_at	KBTBD2	rs4723221	6.4E-08	Same	0.82

Supplementary Table 11 (continued).

Locus name; index SNP	eSNP									Best eSNP			
	eSNP rsID	Dist. index SNP	r ²	eQTL tissue ^{REF}	P-value	Chr	position	Probe*	Transcript**	Best eSNP rsID	P-value	r ² to eSNP	r ² to index SNP
AP5B1 rs4014195	rs11227281	10,264	1.00	Intestine -normal ileum ⁴⁶	3.8E-05	11	65,496,558		EIF1AD	rs11227281	3.8E-05	Same	1.00
	rs11604451	44,888	1.00	LCL (Degner - DNase QTLs) ⁴⁸	2.9E-04	11	65,308,286		DNase QTL 65308500-65308600	rs11604451	2.9E-04	Same	1.00
PTPRO rs7956634	rs2193172	11,148	1.00	Liver (UChicago) ³⁸	6.6E-03	12	15,223,609	A_23_P204304	PTPRO	rs2193172	6.6E-03	Same	1.00
INHBC rs1106766	rs1106766	0	1.00	ER+ breast tumor cells ⁴⁹	7.7E-06	12	56,095,723		GLS2	rs11614506	3.4E-07	0.80	0.80
DPEP1 rs164748	rs164749	68	1.00	Liver ³⁷	2.5E-09	16	88,235,725		C16orf55 (SPATA33)	rs164749	2.5E-09	Same	1.00
	rs460879	4,597	1.00	Prefrontal cortex (Huntington's) ³⁵	3.0E-08	16	88,240,390	10025907286	C16orf55 (SPATA33)	rs460879	3.0E-08	Same	1.00
	rs460879	4,597	1.00	OmentalAdipose ³⁶	2.7E-06	16	88,240,390	10025902450	CHMP1A	rs460879	2.7E-06	Same	1.00
	rs154657	196	1.00	Periph artery plaque [‡]	1.7E-05	16	88,235,597	100143418_TGI_at	SPATA2L	rs154657	1.7E-05	Same	1.00
	rs460879	4,597	1.00	Cerebellum (Huntington's) ³⁵	3.1E-05	16	88,240,390	10025902450	CHMP1A	rs460879	3.1E-05	Same	1.00
	rs459920	22,535	0.97	OmentalAdipose ³⁶	3.7E-44	16	88,258,328	10025907286	C16orf55 (SPATA33)	rs459920	3.7E-44	Same	0.97
	rs459920	22,535	0.97	SubCutAdipose ³⁶	1.2E-30	16	88,258,328	10025907286	C16orf55 (SPATA33)	rs459920	1.2E-30	Same	0.97
	rs459920	22,535	0.97	Cerebellum -all samples ³⁵	1.8E-30	16	88,258,328	10025907286	C16orf55 (SPATA33)	rs459920	1.8E-30	Same	0.97
	rs459920	22,535	0.97	Prefrontal cortex - all samples ³⁵	4.8E-23	16	88,258,328	10025907286	C16orf55 (SPATA33)	rs459920	4.8E-23	Same	0.97
	rs459920	22,535	0.97	Liver ³⁶	2.1E-21	16	88,258,328	10025907286	C16orf55 (SPATA33)	rs459920	2.1E-21	Same	0.97
	rs459920	22,535	0.97	Visual cortex -all samples ³⁵	2.7E-17	16	88,258,328	10025907286	C16orf55 (SPATA33)	rs459920	2.7E-17	Same	0.97
	rs2115401	32,317	0.97	Peripheral artery plaque [‡]	3.6E-07	16	88,268,110	100138862_TGI_at	C16orf55 (SPATA33)	rs2115401	3.6E-07	Same	0.97
	rs467035	31,993	0.97	Peripheral artery plaque [‡]	6.0E-06	16	88,267,786	100131451_TGI_at	SNAI3	rs467035	6.0E-06	Same	0.97
	rs258319	23,732	0.97	Temporal cortex ⁵⁰	4.8E-05	16	88,259,525	ILMN_1759261	C16orf55 (SPATA33)	rs258319	4.8E-05	Same	0.97
	rs2115401	32,317	0.97	Liver (UWash) ³⁸	1.7E-04	16	88,268,110	1300411	C16orf55 (SPATA33)	rs2115401	1.7E-04	Same	0.97

Supplementary Table 11 (continued).

<i>Locus name; index SNP</i>	<i>eSNP</i>									<i>Best eSNP</i>			
	<i>eSNP rsID</i>	<i>Dist. index SNP</i>	<i>r²</i>	<i>eQTL tissue^{REF}</i>	<i>P-value</i>	<i>Chr</i>	<i>position</i>	<i>Probe*</i>	<i>Transcript**</i>	<i>Best eSNP rsID</i>	<i>P-value</i>	<i>r² to eSNP</i>	<i>r² to index SNP</i>
<i>DPEP1</i> rs164748	rs2115401	32,317	0.97	Liver (UChicago) ³⁸	<1e-16	16	88,268,110	A_24_P159335	C16orf55 (SPATA33)	rs2115401	<1e-16	Same	0.97
	rs459920	22,535	0.97	Liver (UChicago) ³⁸	<1e-16	16	88,258,328	A_23_P106694	CHMP1A	rs459920	<1e-16	Same	0.97
<i>TP53INP2</i> rs6088580	rs2273684	244,713	0.94	Blood ⁴⁰	7.6E-08	20	32,993,427	7150537	ACSS2	rs2273684	7.6E-08	Same	0.94
	rs2273684	244,713	0.94	Blood ⁴⁰	1.5E-06	20	32,993,427	2680161	GSS	rs2273684	1.5E-06	Same	0.94
	rs6059909	145,362	0.84	Lymphocytes ⁵¹	2.2E-03	20	32,603,352	GI_31563517-A	MAP1LC3A	rs6059909	2.2E-03	Same	0.81

* Expression QTL results were identified for our index SNPs or their proxies within the following dataset sources: whole blood samples,^{40, 44} Epstein-Barr transformed B-lymphoblastoid cell lines (LCL) from population samples,^{41, 42, 47} DNase-I QTLs in LCLs,⁴⁸ fresh lymphocytes,⁵¹ peripheral blood monocytes,⁴³ ER+ breast cancer tumor cells,⁴⁹ omental and/or subcutaneous adipose,^{36, 41} peripheral artery plaque intestine,⁴⁶ lung,⁴⁵ brain,^{35, 50} and liver.³⁶⁻³⁹

‡ Unpublished (Emilsson)

Supplementary Table 12. Background information on novel replicated loci.

SNPID (Locus name)	Evidence from functional databases		Gene prioritized for functional work	Description of Genes in each Region*
	Source	Gene		
rs228611 (<i>NFKB1</i>)	eSNP	<i>MANBA</i>	<i>NFKB1</i> closest gene	<i>MANBA</i> encodes beta-mannosidase, a lysosomal enzyme that catalyzes the final exoglycosidase step in the degradation pathway for N-linked oligosaccharide moieties of glycoproteins (RefSeq). Rare mutations have been identified as the cause for beta-mannosidosis (OMIM #248510). A mouse model has linked <i>MANBA</i> to lysosomal storage disease in multiple organs, including the kidney. ⁵² <i>NFKB1</i> encodes a pleiotropic transcription factor that is present in almost all cell types and is the endpoint of a series of signal transduction events that are initiated by a vast array of stimuli related to many biological processes such as inflammation, immunity, differentiation, cell growth, tumorigenesis and apoptosis (RefSeq).
	GRAIL	NO		
	DEPICT	NO		
rs7759001 (<i>ZNF204</i>)	eSNP	NA	NO	ZNF204 is known as a transcribed pseudogene (RefSeq). ZNF391 and ZNF184 are two neighboring zinc finger protein-coding genes (RefSeq). Little is known about other nearby genes in the region.
	GRAIL	NO		
	DEPICT	NO		
rs1106766 (<i>INHBC</i>)	eSNP	NO	NO	<i>R3HDM2</i> encodes a protein with R3H domain. <i>INHBC</i> encodes the beta C chain of inhibin, which forms heterodimers with beta A and beta B subunits. Inhibins are involved in hormonal secretion and growth and differentiation of various cell types (RefSeq). <i>GLI1</i> encodes a transcription factor that is activated by the sonic hedgehog signal transduction cascade that regulates stem cell proliferation (RefSeq). The hedgehog-Gli pathway has been implicated in kidney fibrosis in mouse studies. ⁵³ <i>ARHGAP9</i> encodes a member of the Rho-GAP family of GTPase activating proteins, converting them to an inactive GDP-bound state (RefSeq).
	GRAIL	NO		
	DEPICT	NO		
rs10513801 (<i>ETV5</i>)	eSNP	NA	NO	The <i>ETV5</i> gene is a ubiquitously expressed transcription factor. A role in renal development has been shown in a mouse model ⁵⁴ . In humans, it has not yet been connected to kidney disease. <i>DGKG</i> encodes diacylglycerol Kinase, Gamma, a glycerol kinase that metabolizes 1,2,diacylglycerol to produce the second messenger phosphatidic acid. The transcript is expressed in kidney.
	GRAIL	NO		
	DEPICT	NO		
rs3850625 (<i>CACNA1S</i>)	eSNP	NA	NO	<i>CACNA1S</i> encodes one of five subunits of the slowly inactivating L-type voltage-dependent calcium channel, which plays a role in skeletal muscle contraction. Known mutations in <i>CACNA1S</i> have been associated with susceptibility to malignant hyperthermia, hypokalemic periodic paralysis, and thyrotoxic periodic paralysis. The identified variant is a non-synonymous coding SNP at a highly conserved position (R1539C on protein level) that is predicted as damaging or pathogenic by 3 out of 4 prediction software.
	GRAIL	NO		
	DEPICT	NO		
rs10491967 (<i>TSPAN9</i>)	eSNP	NA	<i>TSPAN9</i>	<i>TSPAN9</i> encodes for a member of the tetraspanin family, which assemble in complexes with additional proteins such as integrins to transduct signals. Its association to renal function is unclear; mutations in some integrins cause monogenic kidney disease, as do mutations in the gene paralog CD151. Other genes in the region have not specifically been connected to kidney disease.
	GRAIL	NO		
	DEPICT	<i>TSPAN9</i> FDR<0.05		

Supplementary Table 12 (continued).

SNPID (Locus name)	Evidence from functional databases		Gene prioritized for functional work	Description of Genes in each Region*
	Source	Gene		
rs164748 (<i>DPEP1</i>)	eSNP	<i>C16orf55</i> (also known as <i>SPATA33</i>)	<i>DPEP1</i>	<p><i>DPEP1</i> encodes dipeptidase 1, an enzyme responsible for hydrolysis of glutathione and certain types of antibiotics in the kidney membrane^{55,56} and is highly expressed in the kidney and pancreas (GeneCards), but there are no studies linking this gene to kidney function. <i>RPL13</i> encodes ribosomal protein L13, a protein within the large 60S ribosomal subunit complex⁵⁷. There are no publications linking RPL13 to the kidney or kidney function. <i>SNORD68</i> encodes small nucleolar RNA C/D box 68 a class molecules that modify other RNAs. This particular class of snoRNA is responsible for RNA methylation. There are no known connections to kidney function or disease. <i>SPG7</i> encodes paraplegin, an enzyme component of m-AAA protease (a mitochondrial enzyme responsible for degradation of malformed or misfolded proteins).⁵⁸ Mutations in the <i>SPG7</i> gene lead to monogenic forms of spastic paraplegia.^{59, 60, 60, 61, 61} There are no reported studies linking <i>SPG7</i> and kidney function. <i>ANKRD11</i> encodes ankyrin repeat domain protein 11, a protein inhibiting ligand-dependent activation of transcription. Variation in <i>ANKRD11</i> can lead to KBG syndrome (a disorder with abnormal skeletal development and delay in neurological development).^{62, 63, 63} No associations of <i>ANKRD11</i> and kidney function have been published. <i>C16orf55</i> (also known as <i>SPATA33</i>) encodes spermatogenesis-associated protein 33 and is thought to be involved in spermatogenesis. No associations with kidney disease have been identified.</p>
	GRAIL	NO		
	DEPICT	<i>DPEP1</i> FDR<0.05		
rs8091180 (<i>NFATC1</i>)	eSNP	NA	<i>NFATC1</i>	<p><i>NFATC1</i> encodes for nuclear factor of activated T-cells (cytoplasmic, calcineurin dependent 1) and is a part of a complex involved in the activation of immune response, specifically activation of the T-cell antigen receptor.⁶⁴ In rodents, <i>NFAT1C</i> is potentially involved in proximal tubules after injury⁶⁵ and activation of this nuclear factor may lead to glomerulosclerosis by mutant forms of <i>TRPC6</i> - a podocyte protein involved in maintaining the filtration barrier.⁶⁶ <i>NFAT1C</i> has not been linked with kidney function in humans. <i>ATP9B</i> encodes an ATPase, class II, type 9b - a transmembrane transporter. No association with kidney disease has been previously reported. <i>CTDP1</i> encodes the c-terminal domain of RNA polymerase II subunit A phosphatase. Variations in <i>CTDP1</i> have been associated with congenital cataracts, facial dimorphism and neuropathy in Bulgarian gypsy populations.⁶⁷ There have been no previous reports of <i>CTDP1</i> and kidney disease.</p>
	GRAIL	<i>NFATC1</i> p=0.03		
	DEPICT	NO		
rs11666497 (<i>SIPA1L3</i>)	eSNP	NA	NO	<p><i>SIPA1L3</i> encodes the signal-induced proliferation-associated 1 like 3 protein. There are no published reports in the literature describing this gene or its protein product. <i>WDR87</i> encodes the WD repeat-containing protein 87, a protein 2.8K amino acids in length. There are no published reports of this gene or gene product in the literature.</p>
	GRAIL	NO		
	DEPICT	NO		

Supplementary Table 12 (continued).

SNPID (Locus name)	Evidence from functional databases		Gene prioritized for functional work	Description of Genes in each Region*
	Source	Gene		
rs4667594 (<i>LRP2</i>)	eSNP	NA	NO	<i>LRP2</i> encodes the megalin receptor, which plays an important role in the reabsorption of albumin and other low-molecular-weight proteins ⁶⁸ along with cubilin (<i>CUBN</i>) and amnionless (<i>AMN</i>). <i>LRP2</i> mutations are associated with Donnai-Barrow and facio-oculo-acoustico-renal syndromes. ⁶⁹ <i>DHRS9</i> is a protein-coding gene that encodes dehydrogenase/reductase member 9 (GeneCards). It is a 3-alpha-hydroxysteroid dehydrogenase that is responsible for the synthesis of dihydroxyprogesterone; low levels of activity with retinoids have also been identified (UCSC). There are no published studies linking this gene to kidney function. <i>ABCB11</i> belongs to the ATP-binding cassette superfamily. The protein is involved in bile salt export (www.genecards.org), and mutations are involved in familial intrahepatic cholestasis (OMIM). In addition, this gene may be involved with bile acid transport in the kidney. ⁷⁰
	GRAIL	<i>LRP2</i> p=0.0004		
	DEPICT	<i>LRP2</i> FDR<0.05		
rs6795744 (<i>WNT7A</i>)	eSNP	NA	<i>WNT7A</i> closest gene	<i>WNT7A</i> is a member of set of genes that are signaling proteins, specifically those that are involved in embryogenesis (www.genecards.org). Mutations in this gene have been associated with Fuhrmann syndrome (OMIM #228930) and the Al-Awadi/Raas-Rothschild/Schinzler phocomelia syndrome (AARRS; OMIM #276820), disorders characterized by limb malformation. Wnt-7a signalling may allow for the development of sexual dimorphism via development of the mullerian ducts. ⁷¹ <i>TPRXL</i> is a homeobox gene thought to be involved in embryonic development (GeneCards). There are no published papers linking this gene to kidney function. Of the remaining genes in the region, only <i>XPC</i> has been previously linked to kidney function in the published literature. <i>XPC</i> is a DNA repair gene. Mutations result in Xeroderma pigmentosum, a disease characterized by sunlight sensitivity and early carcinomas. Mutations in <i>XPC</i> may be associated with renal cell carcinoma. ⁷²
	GRAIL	NO		
	DEPICT	NO		
rs7956634 (<i>PTPRO</i>)	eSNP	<i>PTPRO</i>	<i>PTPRO</i> closest gene	<i>RERG</i> belongs to the RAS superfamily and is involved with cell proliferation and tumor pathogenesis (GeneCards). <i>RERG</i> expression may be lost in kidney cancer. ⁷³ <i>PTPRO</i> is expressed in the podocyte foot processes of the kidney; mutations in <i>PTPRO</i> are associated with autosomal-recessive nephrotic syndrome. ⁷⁴ Knock-out mice display reduced eGFR but no proteinuria ⁷⁵ . <i>EPS8</i> is an epidermal growth factor receptor pathway substrate gene (GeneCards). Eps8 proteins are involved in the organization of actin filaments. ⁷⁶
	GRAIL	NO		
	DEPICT	NO		
rs10277115 (<i>UNCX</i>)	eSNP	NA	<i>UNCX</i> (no other genes in the LD block)	<i>UNCX</i> is a transcription factor that is involved in neurogenesis and somitogenesis (GeneCards). In addition, it may be involved in differentiation of the axial skeleton (GeneCards). There is no published literature linking it to kidney function. <i>ZFAND2A</i> is a protein coding gene involved in zinc ion binding (GeneCards). It has been shown to be part of a network of genes expressed in human renal epithelial cells in response to cadmium exposure, a known nephrotoxin. ⁷⁷ <i>GPER</i> is a member of the G-protein coupled receptor family with a primary role of binding estrogen (GeneCards). <i>GPER</i> may mediate the effects of estrogen (but not aldosterone) ⁷⁸ on the vasculature ⁷⁹ including the rat kidney. ⁸⁰ <i>GPER</i> binds estrogen in addition to other substances including endocrine disruptors. ⁸¹
	GRAIL	NO		
	DEPICT	NO		

Supplementary Table 12 (continued).

SNPID (Locus name)	Evidence from functional databases		Gene prioritized for functional work	Description of Genes in each Region*
	Source	Gene		
rs12136063 (SYPL2)	eSNP	SYPL2	SYPL2	<p><i>SYPL2</i> encodes synaptophysin-like 2 protein (also known as mitsugumin 29), a membrane protein involved in the communication between the transverse tubular and junctional sarcoplasmic reticulum membranes (Entrez Gene). It is expressed in the kidney (www.proteinatlas.org). There are no publications linking this gene to kidney function. <i>PSMA5</i> encodes proteasome (prosome, macropain) subunit alpha type 5, which is a proteasome involved in the processing of MHC class I peptides (Entrez Gene) and is expressed in many tissues including the kidney (ProteinAtlas). There are no publications linking this gene to kidney function. There is no disease linked to this gene in OMIM. <i>ATXN7L2</i> encodes for ataxin 7-like 2 protein (GeneCards), which is expressed in several tissues including the kidney (ProteinAtlas). There is no publication linking this gene to kidney function. <i>AMIGO1</i> encodes adhesion molecule with Ig-like domain 1, which is part of a family of transmembrane proteins involved in axon tract development.⁸² It is expressed in the kidney. <i>CELSR2</i> encodes cadherin EGF LAG seven-pass G-type receptor 2, which is a member of the flamingo subfamily cadherins which does not interact with catenins (Entrez Gene). It is a plasma membrane protein postulated to be involved in contact-mediated communication and expressed in many tissues including the kidney.</p>
	GRAIL	NO		
	DEPICT	SYPL2 FDR<0.05		
rs2802729 (SDCCAG8)	eSNP	SDCCAG8	NO	<p><i>SDCCAG8</i> encodes serologically defined colon cancers antigen 8, a centrosome associated protein involved in interphase and mitosis (Entrez Gene). Truncating mutations cause Senior-Loken syndrome 7 (OMIM #613615), an autosomal recessive ciliopathy with nephronophthisis and Leber congenital amaurosis. <i>CEP170</i> encodes centrosomal protein 170kDa, which is the major microtubule-organizing center in animals (Entrez Gene). It is expressed in several tissues including the kidney, but has not been linked to kidney function in any publication. <i>AKT3</i> encodes v-akt murine thymoma viral oncogene homolog 3, which is a kinase regulating cell signaling in response to insulin and growth factors; it is stimulated by PDGF, insulin and IGF-1 (Entrez Gene). It shows low-level expression in the kidney (ProteinAtlas). Rare mutations cause megalencephaly-polymicrogyria-polydactyly-hydrocephalus syndrome (MPPH, OMIM #603387), which does not show organ involvement beyond central nervous defects. There are no publications linking this gene to kidney function.</p>
	GRAIL	NO		
	DEPICT	NO		
rs17216707 (BCAS1)	eSNP	NA	BCAS1 but no fish orthologs	<p><i>BCAS1</i> encodes breast carcinoma amplified sequence 1, which has been implicated as a breast cancer oncogene (Entrez Gene). There is no expression in the kidney (ProteinAtlas). <i>CYP24A1</i> encodes cytochrome P450, family 24, subfamily A, polypeptide 1, which initiates the degradation of 1,25 hydrox-Vitamin D3, thus regulating calcium homeostasis (Entrez Gene). Rare mutations in this gene cause infantile hypercalcemia (OMIM #143880). <i>CYP24A1</i> is expressed in the kidney. <i>PFDN4</i> encodes prefoldin subunit 4, which is part of a molecular chaperone complex needed for correct folding of newly synthesised polypeptides (Entrez Gene). There are no publications linking <i>PFDN4</i> to kidney function. <i>SUMO1P1</i> encodes <i>SUMO1</i> (small ubiquitin-like modifier 1) pseudogene 1 (Entrez Gene); <i>SUMO1</i> is part of a post-translational modification system regulating NFkB under high glucose conditions in kidney mesangial cells;⁸³ sumoylation protects from oxidative stress.⁸⁴ There is no publication linking <i>SUMO1P1</i> to kidney function.</p>
	GRAIL	NO		
	DEPICT	NO		

Supplementary Table 12 (continued).

SNPID (Locus name)	Evidence from functional databases		Gene prioritized for functional work	Description of Genes in each Region*
	Source	Gene		
rs3750082 (<i>KBTD2</i>)	eSNP	<i>KBTD2</i>	<i>KBTD2</i>	AVL9 plays a role in late exocytic transport, ⁸⁵ at the level of the Golgi. ⁸⁶ This gene has not formally been associated with kidney function in the published literature. <i>LSM5</i> is a Sm-like protein that plays a role in pre-mRNA splicing (GeneCards). It has not previously been linked to kidney function in the published literature. There is little known about <i>KBTD2</i> , and no published literature linking it to kidney function. There are several other genes in the region.
	GRAIL	NO		
	DEPICT	<i>KBTD2</i> FDR<0.05		
rs6088580 (<i>TP53INP2</i>)	eSNP	NO	NO	<i>TP53INP2</i> regulates both transcription and autophagy (UCSC). It is a scaffold protein that interacts with <i>VMP1</i> . ⁸⁷ It has not been formally linked to kidney function in the literature. There are dozens of additional genes in the region.
	GRAIL	NO		
	DEPICT	<i>ACSS2</i> and <i>NCOA6</i> FDR<0.05		
rs6459680 (<i>RNF32</i>)	eSNP	NA	NO	<i>RNF32</i> plays a role in spermatogenesis. The gene has not previously been identified in association with kidney function. <i>LMBR1</i> plays a role in limb malformation, which may occur via altered Sonic hedgehog signaling. The gene has not been connected to kidney function or disease.
	GRAIL	NO		
	DEPICT	NO		
rs4014195 (<i>AP5B1</i>)	eSNP	NA	<i>AP5B1</i> but no existing Morpholino	<i>KAT5</i> encodes for K(Lysine) acetyltransferase 5, which is important in transcriptional regulation. It has not been previously identified in association with kidney function. <i>OVOL1</i> encodes for a transcription factor. <i>OVOL1</i> deficient mice of C57BL/6 background show increased perinatal lethality and other abnormalities, including cystic kidneys. ⁸⁸
	GRAIL	NO		
	DEPICT	<i>AP5B1</i> FDR<0.05		
rs2712184 (<i>IGFBP5</i>)	eSNP	NA	NO	<i>IGFBP5</i> (insulin-like growth factor binding protein 5) is a protein-coding gene that either induces or suppresses cell proliferation. <i>IGFBP5</i> participates in cellular pathways of adaptation to hypertonicity in renal medulla under TonEBP control. ⁸⁹ <i>IGFBP2</i> (insulin-like growth factor binding protein 2, 36kDA) regulates cell growth by enhancing or suppressing IGF bioavailability. RNA and protein are ubiquitously expressed. <i>IGFBP2</i> is the most expressed among IGF binding proteins, particularly in glomerular mesangial cells where it is controlled by angiotensin and glucose concentrations. ⁹⁰ <i>TNP1</i> (transition protein 1) is a protein-coding gene. In the course of spermiogenesis, <i>TNP1</i> is a spermatid-specific product that replaces histone and is itself replaced in the mature sperm by protamines. It is expressed in testis. There is no published literature relating this gene or its protein product to kidney function.
	GRAIL	NO		
	DEPICT	NO		
rs10994860 (<i>A1CF</i>)	eSNP	NA	<i>A1CF</i>	<i>A1CF</i> is a protein coding gene that mediates the deamination of apolipoprotein B mRNA through a multi-component enzyme complex including <i>APOBEC-1</i> and a complementation factor coded by <i>A1CF</i> gene. It is expressed in the gastrointestinal tract, liver, pancreas, brain and kidneys. One publication ²⁰ links this gene to kidney reabsorption of urate (elevated serum urate concentrations). <i>ASAH2B</i> (N-acylsphingosine amidohydrolase (non-lysosomal ceramidase) 2B) is a protein-coding gene. Ceramidases are a group of enzymes which catalyze the hydrolysis of ceramides to produce sphingosine but <i>ASAH2B</i> lacks the active site and therefore could be catalytically inactive. No publications link this gene to kidney function. <i>PRKG1</i> encodes the soluble alpha and I beta1 isoforms of the cGMP dependent protein kinase, involved in the regulation of cardiovascular and neuronal functions, smooth muscle relaxation, platelet aggregation, and in cell growth by modulating cellular calcium.
	GRAIL	NO		
	DEPICT	<i>A1CF</i> FDR<0.05		

Supplementary Table 12 (continued).

SNPID (Locus name)	Evidence from functional databases		Gene prioritized for functional work	Description of Genes in each Region*
	Source	Gene		
rs9682041 (SKIL)	eSNP	NA	SKIL	The protein encoded by SKI-like oncogene is a component of the SMAD pathway, that has a regulatory role on the cell division and differentiation through TGF β . RNA and protein are expressed in most human tissues except pancreas. <i>SKIL</i> is implicated in ubiquitin dependent tubulointerstitial fibrosis along with TGF β 1. ⁹¹ Claudin 11 is a member of the claudin family of tight junction associated proteins. The protein encoded by this gene is a major component of the central nervous system (CNS) myelin and plays an important role in regulating proliferation and migration of oligodendrocytes. RNA and protein are expressed in Sertoli cells in testis and oligodendrocytes in the CNS. CLDN11 Claudin family is differentially expressed in tight junctions of human cortical nephron and influences pathologies involving abnormalities of absorption. ⁹² Its expression is significantly unregulated in genetic model of polycystic kidney in early stages. ⁹³ RNA is expressed in cerebral cortex, while the protein is strongly expressed in hippocampus and prostate. There are no publications linking this gene to kidney function.
	GRAIL	SKIL p=0.02		
	DEPICT	SKIL FDR<0.05		
rs163160 (KCNQ1)	eSNP	NA	KCNQ1 work already completed	KCNQ1 encodes a voltage-gated potassium channel required for cardiac repolarization. RNA is ubiquitously expressed but the protein product is mostly expressed in glandular cells. In addition to be recognized as a gene conferring risk of T2D in Caucasians and more recently in African Americans, ⁹⁴ it is also reported as a gene conferring susceptibility to diabetic nephropathy in Asians ⁹⁵ and CKD in African Americans. ⁹⁶ TRPM5 encodes the transient receptor potential cation channel, subfamily M, member 5 gene, a member of the transient receptor potential (TRP) protein family with structural features typical of ion channels (non-selective cations except Ca ²⁺) and mediates a transient membrane depolarization in the presence of low concentrations of intracellular calcium. mRNA is found mainly in the prostate, testis, ovary, colon and leukocytes but the protein is expressed in the majority of human tissues. There are no publications linking this gene or its protein to kidney function. The tumor suppressing subtransferable candidate 4 (TSSC4) gene is located in the p15.5 region of the chromosome 11, a region involved in Wilm's tumor and a known important tumor-suppressor gene region. TSSC4 RNA is widely expressed in human tissues while protein is strongly expressed in renal tubules and moderately in digestive system as well as male and female reproductive systems.
	GRAIL	NO		
	DEPICT	IGF2 (FDR<0.05) but very far away from lead SNP		

*Cited web resources: Entrez Gene: <http://www.ncbi.nlm.nih.gov/gene>; GenCards: www.genecards.org;

OMIM: <http://www.ncbi.nlm.nih.gov/omim>; ProteinAtlas: www.proteinatlas.org; RefSeq:

<http://www.ncbi.nlm.nih.gov/refseq/>; UCSC: www.genome.ucsc.edu.

Supplementary Table 13. DEPICT tissue and cell-type enrichment.*

MeSH ID	Name	MeSH# first level term	MeSH second level term	P-value	FDR	Genes at associated loci§
A05.810	Urinary Tract	Urogenital System	Urinary Tract	0.000226	<0.025	<i>NAT8, SLC17A3, SLC22A2, SLC17A1, LRP2, UMOD, SLC7A9, SLC6A13, SLC34A1, ENSG00000204872, PCK1, SLC47A1, WDR72</i>
A05.810.453	Kidney	Urogenital System	Urinary Tract	0.000292	<0.025	<i>NAT8, SLC17A3, SLC22A2, SLC17A1, LRP2, UMOD, SLC7A9, SLC6A13, SLC34A1, ENSG00000204872, PCK1, SLC47A1, WDR72</i>
A03.620	Liver	Digestive System	Liver	0.000663	<0.025	<i>ITIH1, LPA, SLC17A2, SLC22A1, ITIH3, KNG1, LEAP2, GCKR, SLC22A7, CPS1, A1CF</i>
A11.436.348	Hepatocytes	Cells	Epithelial Cells	0.001338	<0.025	<i>ITIH3, A1CF, LEAP2, SLC17A2, GCKR, AGMAT, ITIH1, SCGN, LCAT, SLC22A3, SLC7A9, SLC22A1</i>
A06.407.071.140	Adrenal Cortex	Endocrine System	Endocrine Glands	0.001574	<0.025	<i>ENSG00000256731, SLC47A1, FNDC4, KCNQ1, RERG, CASP9</i>
A06.407.071	Adrenal Glands	Endocrine System	Endocrine Glands	0.002988	0.025	<i>SLC47A1, FNDC4, ENSG00000256731, KCNQ1, RERG, MAMSTR, CASP9</i>

*DEPICT was used to assess whether genes at associated loci were highly expressed in any of 209 tissue and cell type annotations. In total we found 6 significantly enriched tissues (FDR < 0.05). Only significant tissues are shown in the table.

MeSH: Medical Subject Headings.

§ Lists of genes that are within an associated region and highly expressed in the given tissue or cell type.

Supplementary Table 14. DEPICT pathway analysis. Gene sets with P-value < 1e-05 are shown.

Reconstituted gene set ID	Reconstituted gene set name	Part of meta gene set	P-value	FDR	Reconstituted gene set genes at associated loci
ENSG00000186350	RXRA protein complex	NCOA1 protein complex	5.94E-08	<0.002	<i>NCOA6, SKIL, TRIB1, MED1, CDK12, EDC4, RELA, ARNT, ERBB2, PGAP3, NFATC1, PTPN12, NFKB1, A1CF, NRBP1, TPRKB, AFF4, NRIP1, NFATC3, INO80</i>
MP:0011423	Kidney Cortex Atrophy	Dilated Renal Tubules	3.40E-07	<0.002	<i>SLC7A9, SLC22A2, SLC34A1, VEGFA, NAT8, UMOD, DPEP1, AGMAT, PCK1, LRP2, PTPRO, IGF2, ENSG00000204872, ADAMTS5, DAB2, BMP4, SLC12A4, CA12, SLC6A13, TRIB1</i>
ENSG00000125124	BBS2 protein complex	BBS4 protein complex	4.76E-07	<0.002	<i>KNG1, UMOD, CPS1, ITIH3, ESRP2, RASIP1, PCK1, SLC7A9, AGMAT, SLC22A1, NAT8, LRP2, NUTF2, DACH1, ITIH4, EDC4, SLC22A2, ENSG00000204872, PSMD12, SGMS1</i>
ENSG00000124151	NCOA3 protein complex	NCOA1 protein complex	7.21E-07	<0.002	<i>TRIB1, MED1, NCOA6, FBXL20, ETV5, CDK12, VEGFA, GCKR, GLI2, RAI1, PSKH1, GTF3C2, MOV10, PIK3R1, PHLDA1, PGAP3, ITIH3, NFATC1, SETBP1, RMND5A</i>
MP:0003918	Decreased Kidney Weight	Dilated Renal Tubules	8.75E-07	0.002	<i>RAPSN, IGF2-AS, ACVR2A, NFATC3, PCK1, NAT8, SLC7A9, SLC6A13, IGF2, SLC34A1, PTPRO, ADAMTS5, PAPP, IGF2R, UMOD, GSS, LCAT, TCEA3, SVEP1, AGMAT</i>
GO:0015145	Monosaccharide Transmembrane Transporter Activity	Monosaccharide Trans-membrane Transporter Activity	1.21E-06	0.002	<i>SLC7A9, ENSG00000204872, SLC34A1, AGMAT, LEAP2, NAT8, SLC22A2, TSPAN9, TRIM58, SLC2A9, DPEP1, C12orf68, ENSG00000230288, VEGFA, SLC28A2, DAB2, UMOD, SLC6A13, XYLB, OR2W3</i>
GO:0008514	Organic Anion Transmembrane Transporter Activity	Organic Cationanionzwitterion Transport	2.08E-06	0.006	<i>SLC34A1, SLC22A2, NAT8, SLC6A13, KNG1, SLC22A7, UMOD, SLC22A1, ITIH1, SLC7A9, AGMAT, WDR72, ENSG00000204872, INHBC, SLC47A1, A1CF, LRP2, XYLB, PCK1, DPEP1</i>
MP:0000521	Abnormal Kidney Cortex Morphology	Dilated Renal Tubules	2.60E-06	0.005	<i>UMOD, SLC7A9, SLC22A2, SLC34A1, AGMAT, VEGFA, NAT8, CA12, IGF2-AS, STC1, KNG1, GLI2, RASIP1, LRP2, DPEP1, EYA4, DACH1, SETBP1, PCK1, TSPAN9</i>
GO:2000117	Negative Regulation Of Cysteine-Type Endopeptidase Activity	Negative Regulation Of Cysteine-Type Endopeptidase Activity	3.26E-06	0.004	<i>OR2W3, UMOD, SLC34A1, NRIP1, RMND5A, PSMA5, PRELID1, PTPN12, ENSG00000187446, SNX17, SLC7A6, ENSG00000232656, PSKH1, IZUMO1, SLC22A2, TRIB1, IDI1, NFE2L2, DCDC5, LRP2</i>

Supplementary Table 14 (continued).

Reconstituted gene set ID	Reconstituted gene set name	Part of meta gene set	P-value	FDR	Reconstituted gene set genes at associated loci
MP:0001698	Decreased Embryo Size	Complete Embryonic Lethality During Organogenesis	4.30E-06	0.006	<i>PTPN12, HNRNPR, NCOA6, HSPA4, IGF2, AFF4, IGF2R, ACVR2B, NSD1, VEGFA, MED1, CELF1, NRF1, BMP4, SETDB1, LAMA5, JARID2, RIF1, RMND5A, INO80</i>
MP:0011108	Partial Embryonic Lethality During Organogenesis	Complete Embryonic Lethality During Organogenesis	4.61E-06	0.005	<i>IGF2, LAMA5, VEGFA, BMP4, TRIB1, RARB, PHLDA1, R3HDM2, RHOC, TSPAN9, JARID2, ACVR2B, AFF4, NSD1, PTPN12, SIPA1L3, RASIP1, A1CF, GLI2, GRB10</i>
MP:0001711	Abnormal Placenta Morphology	Abnormal Placenta Labyrinth Morphology	4.80E-06	0.005	<i>VEGFA, ITIH1, GRHL2, TRIB1, KNG1, IGF2, SPATA5L1, NSD1, ITIH3, ESRP2, INO80, RHOC, LEAP2, LAMA5, CELF1, ARNT, ITCH, ERBB2, NCOA6, PAPP</i>
GO:0005355	Glucose Transmembrane Transporter Activity	Monosaccharide Transmembrane Transporter Activity	5.07E-06	0.005	<i>SLC7A9, ENSG00000204872, SLC34A1, AGMAT, SLC2A9, DPEP1, NAT8, SLC22A2, C12orf68, LEAP2, TRIM58, ENSG00000230288, VEGFA, SLC28A2, TSPAN9, GGT7, UMOD, SLC22A7, TFDP2, SLC6A13</i>
GO:0001655	Urogenital System Development	Kidney Development	5.43E-06	0.006	<i>UMOD, EYA4, DACH1, SLC34A1, SLC22A2, RARB, DPEP1, IGF2-AS, ADAMTS5, IGF2, SETBP1, GRHL2, ERBB2, GLI2, LRP2, SHH, PAPP, ACVR2B, BMP4, STC1</i>
GO:0015149	Hexose Transmembrane Transporter Activity	Monosaccharide Transmembrane Transporter Activity	6.45E-06	0.006	<i>SLC7A9, ENSG00000204872, SLC34A1, AGMAT, SLC2A9, NAT8, SLC22A2, DPEP1, TSPAN9, LEAP2, C12orf68, TRIM58, ENSG00000230288, VEGFA, SLC28A2, GGT7, UMOD, PRELID1, SLC22A7, OR2W3</i>
MP:0002981	Increased Liver Weight	Decreased Circulating Cholesterol Level	7.33E-06	0.006	<i>ITIH1, PCK1, KNG1, CPS1, LEAP2, SLC22A1, SLC22A7, ACSS2, SLC7A9, ITIH3, GCKR, IDI1, LPA, INHBC, FASN, A1CF, DPEP1, ITIH4, GSS, IGF2R</i>
GO:0072001	Renal System Development	Kidney Development	7.35E-06	0.005	<i>UMOD, DPEP1, DACH1, SLC22A2, RARB, SLC34A1, ADAMTS5, EYA4, IGF2, IGF2-AS, SETBP1, BMP4, LRP2, NAT8, GLI2, SHH, CLDN11, ERBB2, LAMA5, CA12</i>
MP:0011346	Renal Tubule Atrophy	Dilated Renal Tubules	8.41E-06	0.005	<i>UMOD, SLC34A1, SLC7A9, DPEP1, VEGFA, PTPRO, SLC47A1, AGMAT, SLC22A2, SLC22A7, KNG1, PCK1, MANBA, CPS1, ENSG00000204872, NAT8, WDR72, SLC2A9, LEAP2, SLC22A3</i>
MP:0005459	Decreased Percent Body Fat	Abnormal Glucose Homeostasis	8.78E-06	0.005	<i>CTRL, CELA2B, TRIB1, PCK1, RAPSN, SORT1, ACSS2, SLC34A1, SLC7A9, ENSG00000187446, RSBN1L, FASN, STC1, THADA, LPA, PIK3R1, LMAN2, CASP9, IGF2R, PIGU</i>

Supplementary Table 15. Zebrafish knock-down results.*

	MO	Dose	<i>Pax2a</i>	<i>Nephrin</i>	<i>Slc20a1a</i>
Clutch 1	Control	N/A	0/33 (0%)	0/36 (0%)	0/51 (0%)
	<i>Wnt7aa</i>	250 uM	3/38 (8%) p= 0.24	0/29 (0%)	0/29 (0%)
	<i>Nfkb1</i>	250 uM	0/33 (0%)	0/32 (0%)	0/30 (0%)
	<i>Ptpro</i>	250 uM	0/22 (0%)	0/27 (0%)	0/14 (0%)
	<i>A1cf</i>	250 uM	0/31 (0%)	0/36 (0%)	0/23 (0%)
	<i>Tspan9a+b</i>	125 uM each	1/25 (4%) p=0.43	0/22 (0%)	1/19 (5%) p=0.27
Clutch 2	Control	N/A	3/18 (17%)	0/21 (0%)	0/20 (0%)
	<i>Kbtbd2</i>	250 uM	0/20 (0%) p=0.10	0/22 (0%)	0/17 (0%)
	<i>Uncx</i>	250 uM	3/30 (10%) p=0.66	1/26 (4%) p=1.00	0/19 (0%)
	<i>Dpep</i>	250 uM	1/26 (4%) p=0.29	1/23 (4%) p=1.00	0/24 (0%)
	<i>Sypl2a+b</i>	250 uM each	7/35 (20%) p=1.00	1/25 (4%) p=1.00	0/29 (0%)
Clutch 3	Control	N/A	0/37 (0%)	0/55 (0%)	0/41 (0%)
	<i>Nfatc1</i>	250 uM	2/26 (8%) p=0.17	0/29 (0%)	0/31 (0%)
	<i>Skila+b</i>	62.5 uM each	0/41 (0%)	0/46 (0%)	0/32 (0%)

Zebrafish embryos were injected with morpholinos (MOs) targeting orthologous loci, and renal gene expression was assessed at 48 hours post-fertilization (hpf) by whole mount *in situ* hybridization. Renal markers examined include *pax2a* (global kidney), *nephrin* (podocyte), and *slc20a1a* (proximal tubule). Data are presented as the number of observed abnormalities per total number of embryos scored.

*P-values provided for non-0 results only.

Supplementary Table 16. Study information: discovery. Extensive study names are reported in the Acknowledgements section.

Study ^{REF}	Study Design	N [§]	Study characteristics	Creatinine Measurement	Cystatin measurement
3C ^{97, 98}	Prospective population-based	6440	Study exclusions or disease enrichment. None. Exclusions. Of the 6440, those with genotypes and creatinine were 6431 and those with genotypes and cystatin C were 1243.	Modified kinetic Jaffe reaction.	Particle-enhanced immuno-nephelometric method (BNII, Dade-Behring/ Siemens)
Advance ⁹⁹	Randomised controlled trial	11,140	Study exclusions or disease enrichment. Multicenter trial done by 215 collaborating centres in 20 countries. All patients had T2D and were of Caucasian origin. Those with genotypes and creatinine were 2301. Exclusions. NA	Serum creatinine was measured in local laboratories at individual study sites.	NA
AGES ¹⁰⁰	Population based	3664	Study exclusions or disease enrichment. None. Exclusions. Sample exclusion criteria included sample failure, genotype mismatch with reference panel, and sex mismatch, resulting in clean genotype data on 3219 individuals.	Jaffé reaction.	NA
Amish ^{101, 102}	Population based "founder" cohort	1264	Study exclusions or disease enrichment. None. Exclusions. Age <20, severe chronic disease, call rate <95%, pHWE<10E-6.	Modified kinetic Jaffe reaction.	Particle-enhanced immunonephelometric method (BNII, Dade-Behring).
ARIC ¹⁰³	Prospective, population-based	9713	Study exclusions or disease enrichment. None. Exclusions. Of the 9713 genotyped individuals of European ancestry, we excluded 658 individuals based on discrepancies with previous genotypes, disagreement between reported and genotypic sex, one randomly selected member of a pair of first-degree relatives, or outlier based on measures of average DST or >8 SD away on any of the first 10 principal components.	Modified kinetic Jaffé reaction.	Particle enhanced immunonephelometric assay (N Latex Cystatin C, Dade Behring).
ASPS ^{104, 105}	Prospective study	922	Study exclusions or disease enrichment. Excluded were subjects with history of neuropsychiatric disease, previous stroke and/or TIA, and dementia. Exclusions. Of the 922 participants who underwent genotyping, we made the following exclusions: sample call rate <98% (74). This resulted in 848 genotyped individuals.	Modified kinetic Jaffé reaction.	NA
AUSTWIN ¹⁰⁶	Families, population-based	9592	Study exclusions or disease enrichment. NA. Exclusions. NA.	Jaffé method, Hitachi 917 analyser.	NA
BLSA ¹⁰⁷	Population based study	1200	Study exclusions or disease enrichment. None. Exclusions. Those of non-European descent or with missing phenotype information.	Modified kinetic Jaffé reaction.	NA
BMES ¹⁰⁸⁻¹¹⁰	Prospective cohort study	2761	Study exclusions or disease enrichment. None. Exclusions. Sample call rate <95% (n=9), outlying autosomal heterozygosity (n=28), sex discrepancies or ambiguous sample identification (n=69), cryptic relatedness (average IBD sharing proportion > 0.1875: n=121), non-European ancestry (n=13). This resulted in 2534 genotyped individuals.	Measured within 4 hours of collection using a Hitachi 747 Biochemistry analyser (Roche reagents, modified kinetic Jaffé).	NA

Abbreviations: T2D = type 2 diabetes. §Total genotyped sample size.

Supplementary Table 16 (continued).

Study ^{REF}	Study Design	N [§]	Study characteristics	Creatinine Measurement	Cystatin measurement
CHS ^{111, 112}	Prospective population-based	3397	Study exclusions or disease enrichment. A total of 1908 persons excluded from the GWAS study sample due to the presence at study baseline of coronary heart disease, congestive heart failure, peripheral vascular disease, valvular heart disease, stroke or transient ischemic attack or lack of available DNA. Exclusions. The present report is based upon genotyping results from 3329 Caucasian participants, who were free of clinical cardiovascular disease at baseline, consented to genetic testing, and had DNA available for genotyping. Genotyping was successful in 3291 persons.	Colorimetric method (Ektachem 700, Eastman Kodak).	Particle-enhanced immunonephelometric assay [N Latex Cystatin C, Dade Behring (now Siemens), Deerfield, Ill, USA] with a nephelometer [BNII, Dade Behring (now Siemens)].
CROATIA-KORCULA ¹¹³	Cross-sectional, family-based	888	Study exclusions or disease enrichment. None. Exclusions. Subjects were excluded if they fulfilled any of the following criteria: genotypic call rate <97%; reported vs. genotypic sex mismatch; unexpectedly low genomic sharing with 1 st degree relatives; excess of autosomal heterozygosity; outliers identified by IBS clustering analysis; pregnant women.	Jaffé rate method in plasma.	NA
CROATIA-SPLIT ¹¹⁴	Population-based	499	Study exclusions or disease enrichment. Fasting urine and blood samples were collected from 1012 healthy volunteers aged 18+ from Split on the Dalmatian coast in Croatia in 2008/2011. Exclusions. Missing creatinine levels	Jaffé protein compensated method in the serum.	NA
CROATIA-VIS ^{115, 116}	Cross-sectional, family-based	768	Study exclusions or disease enrichment. None. Exclusions. Subjects were excluded if they fulfilled any of the following criteria: genotypic call rate <97%, mismatch between reported and genotypic sex, unexpectedly low genomic sharing with first degree relatives, excess autosomal heterozygosity, or outliers identified by IBS clustering analysis. Pregnant women were excluded from the study.	Enzymatic photometric assay using an ADVIA1650 clinical chemistry analyzer (Siemens Healthcare Diagnostics GmbH, Eschborn, Germany) ¹¹⁷ at the Institute for Clinical Chemistry and Laboratory Medicine, Regensburg University Medical Center, Germany.	NA
DESIR ¹¹⁸	Population-based	731	Study exclusions or disease enrichment. Excluded 15 ethnicity outliers. Exclusions. Individuals with genotype call rate <0.90, outlying heterozygosity, gender discrepancies, missing clinical data, cryptic relationships, non-European.	Kinetic colorimetry using Jaffé's Method. The assay utilized a Technicon DAX24 automated analyser from Bayer Diagnostics, Puteaux, France or a Specific or a Delta from Konelab, Evry, France.	NA
EGCUT 370K ^{119, 120} EGCUT Omni ^{119, 120}	Population-based	2700 9500	Study exclusions or disease enrichment. None. Exclusions. Missing creatinine levels; genetic outliers; cryptic relatedness (one random member up to 2nd cousins was only included)	Modified Jaffé protein compensated method in the serum.	NA

Abbreviations: T2D = type 2 diabetes. §Total genotyped sample size.

Supplementary Table 16 (continued).

Study^{REF}	Study Design	N[§]	Study characteristics	Creatinine Measurement	Cystatin measurement
ERF ¹²¹	Population based family study	2834	Study exclusions or disease enrichment. None. Exclusions. NA.	Jaffé rate method using a Synchron LX20.	NA
FamHS ¹²²	Family based	3838	Study exclusions or disease enrichment. None. Exclusions. Age <18, call rate <98%, pHWE <10E-6, sex mismatch, non-European ancestry.	Thin film adaptation of the amidohydrolase enzymatic method using the Vitros analyzer (Johnson & Johnson Clinical Diagnostics, Inc. Rochester NY 14650).	Immune particle-enhanced turbidimetric (PET) kit (DAKO A/S, Produktionsvej 42, DK-2600 Glostrup, Denmark. Code no. K0071)
FHS ¹²³⁻¹²⁵	Prospective family-based	9300	Study exclusions or disease enrichment. None. Exclusions. Of the 9,274 participants who underwent genotyping, we made the following exclusions: sample call rate <97% (n=666), genotype heterozygosity > 5 SDs, and ambiguous family data (n=127). This resulted in 8481 genotyped individuals.	Modified kinetic Jaffé reaction.	Particle-enhanced immunonephelometric method, Cystatin C was measured (BNII, Dade-Behring).
GENOA ¹²⁶⁻¹²⁸	Family-based	1553	Study exclusions or disease enrichment. 5 Non-white, 1 Missing Exam. Exclusions. For the Affymetrix 6.0 data, we excluded 25 subjects who failed pre-processing, 123 with Contrast QC<0.4, 2 for inconsistent relatedness, 11 identical twins. We re-ran the samples that failed pre-processing or the Contrast QC filter on the Affymetrix 6.0 data along with 50 that had passed. Of these samples, 19 failed genotyping completely, 9 had call rate <0.95, 2 had inconsistent relatedness, and 2 were identical twins. Of the 1509 remaining samples, 346 had no serum creatinine information. Our final data had 1163 samples with genotype and phenotype data.	Compensated rate Jaffé reaction.	NA
HABC	prospective cohort study	1663	Study exclusions or disease enrichment. None. Exclusions. Sample failure, genotypic sex mismatch, and first-degree relative of an included individual based on genotype data.	Colorimetric technique on a Johnson & Johnson VITROS 950 Chemistry Analyzer (Johnson & Johnson, New Brunswick, N.J., USA) using the enzymatic method.	BNII nephelometer (Dade Behring Inc., Deerfield, Ill., USA) that utilized a particleenhanced immunonephelometric assay (N Latex Cystatin C).
HCS ¹²⁹	Population-based	2204	Study exclusions or disease enrichment. None. Exclusions. Individuals with genotype call rate <0.95, outlying heterozygosity, gender discrepancies, missing clinical data, cryptic relationships, non-European ancestry or missing creatinine measurement.	Siemens Dimension Vista 1500 Intelligent Lab System using a modified Jaffé assay in a NATA accredited lab.	NA
HPFS ¹³⁰⁻¹³⁴	Nested case-control study of T2D	2487	Study exclusions or disease enrichment. All subjects had T2D. Exclusions. Of the 2487 subjects with genome-wide scans, 818 (all T2D cases) had creatinine measured.	Modified kinetic Jaffé reaction in plasma.	NA

Abbreviations: T2D = type 2 diabetes. §Total genotyped sample size.

Supplementary Table 16 (continued).

Study^{REF}	Study Design#	N*	Study characteristics	Creatinine Measurement	Cystatin measurement
HYPERGENES HTN cases ¹³⁵	Cases from a nested case-control study of Hypertension	2125	Study exclusions or disease enrichment. We excluded 81 samples with call rate <0.95, 33 with genotypic sex mis-match, 39 duplicated, 94 related subjects, and 13 outliers. Exclusions. Of the 1865 subjects with GWA data after QC, 1591 had creatine measured	Kinetic Jaffé reaction.	NA
HYPERGENES HTN controls ¹³⁵	Controls from a nested case-control study of Hypertension	1934	Study exclusions or disease enrichment. We excluded 62 samples with call rate <0.95, 23 with genotypic sex mismatch, 25 duplicated, 62 related subjects, and 12 outliers. Exclusions. Of the 1750 subjects with GWA data after QC, 1662 had creatine measured	Kinetic Jaffé reaction.	NA
INCIPE ¹³⁶	Cross-sectional, population based	940	Study exclusions or disease enrichment. Excluded were individuals <40 year old. Exclusions. Pregnant women	Kinetic rates using the Jaffé method, recalibrated to standardized creatinine determinations obtained at the Cleveland Clinic Research Laboratory.	Immunonephelometric method that used monospecific antisera on an Immage 800 (Beckman Coulter).
INGI-CARLANTINO ¹³⁷⁻¹³⁹	Isolated population	679	Study exclusions or disease enrichment. None. Exclusions. We obtained the levels of creatinine of 447 participants; excluded were those with call rate <97%.	Jaffé reaction.	NA
INGI-CILENTO ¹⁴⁰⁻¹⁴⁷	Cross-sectional population-based study of isolated populations with pedigree information	859	Study exclusions or disease enrichment. None. Exclusions. 38 individuals aged <20 years	Modified kinetic Jaffé reaction.	NA
INGI-FVG ¹³⁷⁻¹³⁹	Isolated population	1376	Study exclusions or disease enrichment. None. Exclusions. We obtained the levels of creatinine of 874 participants; excluded were those with call rate <97%.	Jaffé reaction.	NA
INGI-VAL BORBERA ¹⁴⁸	Family Population-based	1665	Study exclusions or disease enrichment. None. Exclusions. Of the 1665 participants who underwent genotyping, one was excluded because of a sample call rate <95%.	Jaffé reaction.	NA

Abbreviations: T2D = type 2 diabetes. §Total genotyped sample size.

Supplementary Table 16 (continued).

Study ^{REF}	Study Design	N[§]	Study characteristics	Creatinine Measurement	Cystatin measurement
JUPITER ^{149, 150}	Randomized, placebo-controlled trial	8782	Study exclusions or disease enrichment. Excluded were subjects with LDL-C \geq 130 mg/dl or CRP $<$ 2 mg/l. Exclusions. None	Performed in JUPITER core central laboratories using the Roche Modular Analytics Chemistry System with Roche creatinine reagents (modified Jaffe reaction with rate blanking) (Roche Diagnostics, Township of Branchburg, New Jersey).	NA
KORA-F3 ^{151, 152}	Prospective population based	1644	Study exclusions or disease enrichment. None. Exclusions. None.	Modified kinetic Jaffe reaction.	Particle-enhanced immunonephelometric method (BNII, Dade-Behring).
KORA-F4 ^{151, 152}		1814			
MESA ¹⁵³	Community-based cohort study	2520	Study exclusions or disease enrichment. None. Exclusions. NA.	Serum creatinine was measured by rate reflectance spectrophotometry using thin film adaptation of the creatine amidinohydrolase method on the Vitros analyzer (Johnson & Johnson Clinical Diagnostics, Inc., Rochester, NY 14650). The laboratory analytical CV is 2.2%. All creatinine measurements for the MDRD Study were performed at Cleveland Clinic Labs using a CX3 assay. The Vitros analyzer used here was previously calibrated to a CX3 machine with the Cleveland Clinic lab and found the results were nearly identical.	BNII nephelometer (Dade Behring Inc., Deerfield, IL) that utilizes a particle enhanced immunonephelometric assay (N Latex Cystatin-C) 7 on fasting plasma specimens stored at -70°C. The assay is stable over 5 cycles of freeze / thaw. Among 61 healthy individuals with 3 cystatin-C measurements over a 6-month period, the intra-individual coefficient of variation was 7.7%.
MICROS ¹⁵⁴⁻¹⁵⁶	Cross-sectional, population-based study on extended pedigrees	1391	Study exclusions or disease enrichment. None. Exclusions. Excluded were samples with overall SNP call rate $<$ 95%, showing excess of heterozygosity, or being classified as outliers by IBS clustering analysis.	Enzymatic photometric assay using an ADVIA1650 clinical chemistry analyzer (Siemens Healthcare Diagnostics GmbH, Eschborn, Germany).	BN-ProSpec analyzer (Dade Behring, Marburg, Germany) at the Institute for Clinical Chemistry and Laboratory Medicine, Regensburg University Medical Center, Germany.

Abbreviations: T2D = type 2 diabetes. §Total genotyped sample size.

Supplementary Table 16 (continued).

Study ^{REF}	Study Design	N§	Study characteristics	Creatinine Measurement	Cystatin measurement
NESDA ^{157, 158}	Longitudinal cohort study	1862	Study exclusions or disease enrichment. Individuals were almost all cases with major depression or anxiety disorder (N=1705) and of Western-European ancestry. Exclusions. Excluded were ethnic outliers, XO and XXY samples, and samples with a call rate <95%, high genome-wide homo- or heterozygosity, or excess of IBS.	Enzymatic assay from Roche Modular P unit (CREAplus; Roche Diagnostics, Ltd., Lewes, UK). Outliers (>4SD from mean) were excluded.	BNII nephelometer on plasma specimens (N Latex Cystatin C; Dade Behring Inc., Deerfield, IL). Excluded were values >4SD from the mean.
NHS ^{130, 132, 133, 159, 160}	Nested case-control study of T2D	3286	Study exclusions or disease enrichment. All individuals had T2D. Exclusions. Of the 3286 subjects with GWA data, 784 had creatinine measured.	Modified kinetic Jaffé reaction in plasma. Creatinine values were not normalized to the Cleveland Clinic standard.	NA
NSPHS ^{161, 162}	cross-sectional, family-based	565	Study exclusions or disease enrichment. None. Exclusions. Subjects were excluded if they fulfilled any of the following criteria: genotypic call rate <97%, mismatch between reported and genotypic sex, unexpectedly low genomic sharing with first degree relatives, excess of autosomal heterozygosity, or outliers identified by IBS clustering analysis.	Enzymatic photometric assay in plasma using an ADVIA1650 clinical chemistry analyzer (Siemens Healthcare Diagnostics GmbH, Eschborn, Germany).	NA
OGP-TALANA ¹⁶³⁻¹⁶⁵	Isolated population	1376	Study exclusions or disease enrichment. None. Exclusions. We obtained the levels of creatinine of 874 participants; exclusions were samples with call rate <97%.	Jaffé reaction.	NA
ORCADES ¹⁶⁶	cross-sectional, family-based	704	Study exclusions or disease enrichment. None. Exclusions. Subjects were excluded if they fulfilled any of the following criteria: genotypic call rate <97%, mismatch between reported and genotypic sex, unexpectedly low genomic sharing with first degree relatives, excess autosomal heterozygosity, or outliers identified by IBS clustering analysis.	Enzymatic photometric assay in plasma using an ADVIA1650 clinical chemistry analyzer (Siemens Healthcare Diagnostics GmbH, Eschborn, Germany) ¹¹⁷ at the Institute for Clinical Chemistry and Laboratory Medicine, Regensburg University Medical Center, Germany.	NA
POPGEN ¹⁶⁷	Prospective population-based	1163	Study exclusions or disease enrichment. None. Exclusions. Samples with > 5% missing genotypes, showing excess genetic dissimilarity to the remaining subjects, or with evidence for a cryptic relatedness to other study participants were removed. These quality control measures left 1241 control samples for inclusion in the study. Of them, 1163 had serum creatinine available. All sex assignments could be verified by reference to the proportion of heterozygous SNPs on the X chromosome.	Enzymatic in vitro assay (CREAplus, Cobas®, Roche Diagnostics, Indianapolis, IN)	NA
PREVEND ¹⁶⁸	population based	3634	Study exclusions or disease enrichment. Subjects were aged 28 to 75 years and were enriched for microalbuminuria. Exclusions. NA	NA	BNII nephelometer (Dade Behring Inc.).

Abbreviations: T2D = type 2 diabetes. §Total genotyped sample size.

Supplementary Table 16 (continued).

Study ^{REF}	Study Design	N [§]	Study characteristics	Creatinine Measurement	Cystatin measurement
PROSPER-PHASE ¹⁶⁹⁻¹⁷¹	randomized placebo controlled trial	5763	Study exclusions or disease enrichment. Subjects were included when they had had a vascular event or were at increased risk of vascular disease; age: 70-82. Exclusions. After QC (sample rate <97.5%, genotype heterozygosity, ambiguous family data), 5244 samples were available for analysis.	Measured at central laboratories, one in each of the three participating countries.	NA
RS-I ¹⁷²⁻¹⁷⁴	Prospective population based study	5974	Study exclusions or disease enrichment. NA. Exclusions. Excluded were samples with call rate < 97.5%, excess autosomal heterozygosity >0.336 (~FDR <0.1%), mismatch between called and phenotypic gender, or outliers identified by the IBS clustering analysis with >3 SDs from population mean or IBS probabilities >97%.	Modified kinetic Jaffé reaction.	NA
RS-II ¹⁷²⁻¹⁷⁴	Prospective population based study	1895	Study exclusions or disease enrichment. NA. Exclusions. Excluded were samples with call-rate <97.5%, excess of autosomal heterozygosity >0.336 (~FDR<0.1%), mismatch between called and phenotypic gender, and outliers identified by the IBS clustering analysis with >3 SDs from population mean or IBS probabilities >97%.	Modified kinetic Jaffé reaction.	NA
SAPALDIA	population based	1444	Study exclusions or disease enrichment. NA. Exclusions. Excluded were subjects with cryptic relatedness and call rate <95%.	Jaffé reaction (Roche) and calibrated to the Roche enzymatic gold standard reference yielding slightly lower serum creatinine measurements than the Cleveland Clinic Jaffé reaction.	NA
SHIP ^{175, 176}	Prospective population-based	4105	Study exclusions or disease enrichment. None. Exclusions. Excluded were samples with call rate <92%, duplicate samples (by IBS estimation), individuals with reported/genotyped gender mismatch.	Jaffé method. (A blood sample was drawn from the cubital vein in the supine position - the participants were non-fasting due to the duration of the cumulative examinations, 4-6 hours in total).	Siemens N Latex Cystatin C assay, a particle-enhanced nephelometric immunoassay, on the BN ProSpec® System.
SHIP-TREND	Prospective population-based	988	Study exclusions or disease enrichment. Excluded were individuals with no genotype and those with known T2D. Exclusions. Excluded were samples with call rate <94%, duplicate samples (by IBS estimation), individuals with reported/genotyped gender mismatch.	Jaffé method.	Dimension Vista® System, CYSC Flex® reagent cartridge, SIEMENS, Eschborn, Germany
SORBS ¹⁷⁷⁻¹⁷⁹	Population-based	1097	Study exclusions or disease enrichment. None. Exclusions. Sample call rate<0.94; Ethnic outliers; duplicates; gender mismatch; IBS>0.2	Kinetic enzymatic method.	NA

Abbreviations: T2D = type 2 diabetes. §Total genotyped sample size.

Supplementary Table 16 (continued).

Study^{REF}	Study Design	N[§]	Study characteristics	Creatinine Measurement	Cystatin measurement
WGHS ¹⁸⁰	Prospective population based	21,940	Study exclusions or disease enrichment. Excluded were individuals of non-European ancestry. Exclusions. Samples with <98% successful SNP calls were excluded. Data had been collected on 21940 individuals had successful genotype information and verified European ancestry at the time of the analysis.	Rate-blanked method based on the Jaffé reaction using Roche Diagnostics reagents with reproducibility of 3.67% and 1.60% at concentrations of 1.17 and 6.40 mg/dL, respectively.	NA
YFS	Population based	2023	Study exclusions or disease enrichment. NA. Exclusions. None.	Jaffé method (picric acid; Olympus Diagnostica GmbH) from frozen plasma samples in 2007.	NA

Abbreviations: T2D = type 2 diabetes. §Total genotyped sample size.

Supplementary Table 17. Study information, replication. Extensive study names are reported in the Acknowledgements section.

Study Name ^{REF}	Study Design	N§	Study characteristics	Creatinine Measurement	Cystatin measurement
Bus Santé ^{181, 182}	Cross-sectional population-based	5589	Study exclusions or disease enrichment. None. Exclusions. Unsuccessful genotyping (N=106). Analyses were restricted to Caucasians, defined as self-reported citizenship corresponding to South / North America, Europe, and Australia regions (N=4671). Creatinine was available in 4408 participants.	Modified kinetic Jaffé reaction (Abbott Architect).	NA
EGCUT replic ¹²⁰	Population-based	8576	Study exclusions or disease enrichment. None. Exclusions. Sample call rate <95% (N=279), genotype heterozygosity >5 SDs and ambiguous family data (N=264), gender mismatch (N=241).	Modified kinetic Jaffé reaction.	Immune turbidimetric method.
ESTHER ¹⁸³⁻¹⁸⁵	Prospective cohort study	3490	Study exclusions or disease enrichment. Inclusion criteria: age ≥50 and good knowledge of German language. Exclusions. Samples with insufficient amount of DNA were not genotyped.	Modified kinetic Jaffé reaction.	NA
GENDIAN ^{186, 187}	Cohort study of T2D complications	1026	Study exclusions or disease enrichment. After exclusion of N=53 subjects due to subject QC, excluded were patients with ESRD (N=438) or advanced, histologically proven diabetic nephropathy (N=84) or missing phenotype (N=1). Exclusions: call-rate<95% (N=22); relatedness and duplicates (N=11); gender mismatch (N=16); ethnicity check (N=4).	Enzymatic assay.	Dade Behring assay (BNII)
GHS-I ^{188, 189} GHS-II ^{188, 189}	Population-based	3422 1438	Study exclusions or disease enrichment. Excluded if age <35 and >74. Exclusions: N=426 based on a call-rate <97%, a rate of heterozygosity of 3 SDs away from the mean, disagreement between reported and genotypic sex, estimated IBD >0.25, IBS based principal components. This resulted in 2996 genotyped individuals.	Modified kinetic Jaffé reaction (Abbott).	NA
GSK ¹⁹⁰⁻¹⁹²	Case-control study	1776	Study exclusions or disease enrichment. Cases were patients with unipolar recurrent depression; exclusion criteria: presence of manic or hypomanic episodes, mood incongruent psychotic symptoms, lifetime diagnosis of drug abuse and depressive symptoms secondary to alcohol or substance abuse or dependence or to a medical illness or medication. Controls: exclusion criteria: anxiety and affective disorders. Exclusions: call-rate<98% (N=2).	Standard clinical chemistry assays from Roche Hitachi.	NA
HRS	Population-based Longitudinal	12,507	Study exclusions or disease enrichment. Non-European participants were excluded. Exclusions. Sample call-rate<98%, gender discrepancy, unexpected duplicates and/or relatives with KC>1/32, missing cystatin C data.	NA	See note (1)
KORAF3 non-GWAS ^{151, 152} KORAF4 non-GWAS ^{151, 152}	Prospective Population based	1498 1202	Study exclusions or disease enrichment. None. Exclusions. None.	Modified kinetic Jaffé reaction.	Particle-enhanced immunonephelometric method (BNII, Dade-Behring).

§ Total genotyped sample size.

(1) Measured from dried blood spots (DBS) obtained during a face-to-face interview with the respondent at the same time as saliva collection. Blood was taken by pricking the participant's finger with a sterile lancet after cleansing the finger with an alcohol swab. Droplets of blood were extracted from the finger and directly placed on specially treated filter paper. Cystatin C for was assayed at the University of Vermont in 2006 and 2008. Serum equivalent values were calculated using a formula derived from a validation study done in conjunction with the USC/UCLA Center on Biodemography and Population Health.

Supplementary Table 17 (continued).

Study Name ^{REF}	Study Design	N§	Study characteristics	Creatinine Measurement	Cystatin measurement
IPM ¹⁹³	Hospital-based	1747	Study exclusions or disease enrichment. None. Exclusions. None.	Colorimetric method in (CPT82565) performed by the New York State / CLIA Clinical Chemistry Laboratory at Mount Sinai Medical Center.	NA
LURIC ¹⁹⁴	Case control	3061	Study exclusions or disease enrichment. Inclusion criteria: Caucasian, availability of coronary angiogram, stable condition except acute coronary syndrome. Exclusion criteria: any chronic disease other than cardiovascular, any acute illness other than ACS, any malignancy in the previous five years. Exclusions. Sample call-rate<95%, ambiguous sex, duplicates, relatedness. This resulted in 3061 genotyped individuals.	Jaffé method (CREA/Hitachi 717).	N LATEX Cystatin C/ Behring nephelometer II.
OGP ¹⁶³⁻¹⁶⁵	Population-based study with pedigree information	9554	Study exclusions or disease enrichment. None. Exclusions. NA.	Modified kinetic Jaffé reaction or enzymatic reaction.	NA
SAPHIR ^{195, 196}	Healthy working population	1726	Study exclusions or disease enrichment. None. Exclusions. None.	Modified kinetic Jaffé reaction (CREA®, Roche Diagnostics GmbH, Mannheim, Germany).	NA
SKIPOGH ¹⁹⁷	cross-sectional family-based population-based	941	Study exclusions or disease enrichment. None. Exclusions. Excluded were 71 participants with call-rate <90%, resulting in 870 genotyped individuals.	IDMS-traceable Jaffé kinetic compensated method.	NA
Vanderbilt Omni1 Vanderbilt Omni5	Practice-based cohort	5184 2005	Study exclusions or disease enrichment. Samples chosen based on being a case or control for one of 31 pharmacogenetic analyses. Exclusions. Excluded individuals of non-white ancestry in the electronic medical record. Also excluded any lab measurements of individuals after initiation of dialysis or a kidney transplant. The median outpatient creatinine value was chosen.	Jaffé reaction.	NA
Vanderbilt 660W	Practice-based cohort	3021	Study exclusions or disease enrichment. Samples chosen for normal cardiac conduction, meaning that at some point in time they had a normal electrocardiogram without the presence of heart disease, arrhythmias, or electro-cardiographically-active medications. Exclusions. Children (age<18) and individuals of non-white ancestry in the electronic medical record. Also excluded any lab measurements from individuals after initiation of dialysis or a kidney transplant. The median outpatient creatinine value was chosen. At some point in their electronic medical record, the patients were absent of heart disease, but could later develop it.	Jaffé reaction.	NA

§ Total genotyped sample size.

Supplementary Table 18. Genotyping information, discovery studies.

Study name	Array type	Genotype calling	QC filters genotyped SNPs used for imputation‡	No. of SNPs used for imputation	Imputation software	Imputation Backbone (NCBI build)	Filtering of imputed genotypes†	Data management and statistical analysis
3C	Illumina Human610-Quad	Illumina BeadStudio	call rate < 98%, pHWE<10e-6, MAF<1%	492,897	MACH	1000 Genomes - CEU - Dec 2010 (build 37)	none	R, ProbABEL
Advance	Affymetrix 5.0 Affymetrix 6.0	Affymetrix	Affymetrix 5.0: call rate < 96% (<99% if MAF < 5%); Affymetrix 6.0: call rate < 97% (<99% if MAF<5%)	876,688	IMPUTE 2 v2.1.2	1000 Genomes - CEU Pilot - Jun 2010 plus HapMap3 rel. 2 (all available haplotypes) – Feb 2009 (build 36)	imputation info<0.5	SNPTEST
AGES	Illumina Hu370CNV	Illumina	pHWE<1e-6, call rate<97%, mishap p<1e-9, MAF<1%, SNPs not in Hapmap or with strand issues when merging with Hapmap	329,804	MACH v1.0.16	HapMap rel. 22 (build 36)	none	R, ProbABEL, Linear and Logistic Regression
Amish	Affymetrix 500K	BRLMM	MAF<1%, non-HapMap, call rate < 95%, pHWE<10e-6	338,598	MACH v1.0.15	HapMap rel. 22 - phased CEU haplotypes (build 36)	none	Measured genotype accounting for polygenic component
ARIC	Affy 6.0	Birdseed	call rate <95%, MAF<1%, pHWE <10e-5	669,450	MACH v1.0.16	HapMap rel. 22 (build 36)	none	R, ProbABEL, PLINK
ASPS	Illumina Human610-Quad BeadChip	Illumina	pHWE<1e-6, call rate < 98%, mishap p<1e-9, MAF<1%, Mendelian errors>100, SNPs not in Hapmap or with strand issues when merging with Hapmap	550,635	MACH v1.0.15	HapMap rel. 22 (build 36)	none	R, linear and logistic fixed effects model
AUSTWIN	Illumina370, iLLumina610	BeadStudio-gencall v3.0	call rate < 95%, pHWE<1e-6	271,069	MACH v1.0.15	HapMap rel. 22 - phased CEU haplotypes (build 36)	Rsqr<0.3	Stata, SPSS
BLSA	Illumina Infinium HumanHap 550K	BeadStudio	call rate <99%, MAF <1%, pHWE<10e-4	501,764	MACH v1.0.15	HapMap rel. 21 - phased CEU haplotypes (build 35)	MAF<0.01, r2hat < 0.3	SAS, MERLIN, R

Supplementary Table 18 (continued).

Study name	Array type	Genotype calling	QC filters genotyped SNPs used for imputation‡	No. of SNPs used for imputation	Imputation software	Imputation Backbone (NCBI build)	Filtering of imputed genotypes†	Data management and statistical analysis
BMES	Illumina 670K-Quad	CHIAMO	call rate <95%, pHWE<10e-6, MAF<1%, genotype discrepancies in > 2.5% of 1356 samples independently genotyped for the Illumina 610K array	501,910	MACH v1.0.16	HapMap rel. 22 - phased CEU haplotypes (build 36.1)	none	Plink, mach2dat, SAS
CHS	Illumina 370CNV	Illumina BeadStudio	call rate<97%, heterozygotes=0, pHWE<10e-5, SNP not in HapMap	306,655	BimBam v0.99	HapMap CEU rel. 22 (build 36)	dosage variance < 0.01	R, linear and logistic regression, robust SE estimates
CROATIA-KORCULA	Illumina Infinium HumanCNV370 v1 SNP bead microarrays	BeadStudio	MAF<1%, pHWE<1e-6, call rate < 98%	317,896	MACH v1.0.16	HapMap rel. 22 (build 36)	none	R, GenABEL, ProbABEL;
CROATIA-SPLIT	Illumina HAP370CNV	Illumina	call rate <98%, pHWE<10e-10	330,997	MACH v1.0.15	HapMap rel. 22 - phased CEU haplotypes (build 36)	none	R (GenABEL, ProABEL)
CROATIA-VIS	Illumina HumanHap300 beadchip	BeadStudio	MAF≤1% , pHWE ≤1e-6 , call rate ≤ 98%	305,068	MACH v1.0.15	HapMap rel. 22 (build 36)	none	R, GenABEL, ProbABEL;
DESIR	Illumina Human CNV370-Duo Array and Illumina HAP300 array	Illumina Beadstation Genotyping Solution	call rate < 95% p HWE < 1e-4	291,609	IMPUTE v0.3.2 (genotyped SNPs used where available)	HapMap rel. 22 - phased CEU haplotypes (build 36)	none	SNPTEST v2.2.0
EGCUT 370K	Illumina 370CNV; Illumina OmniExpress	GenCall GenomeStudio	call rate <95%, pHWE<10e-6, MAF<1%	320784; 622800	IMPUTE v1.0	HapMap 2 CEU rel. 22	none	PLINK
EGCUT Omni	Illumina 370CNV; Illumina OmniExpress	GenCall GenomeStudio	call rate <95%, pHWE<10e-6, MAF<1%	320784; 622800	IMPUTE v1.0	HapMap 2 CEU rel. 22	none	PLINK
ERF	Illumina 6K/318K/380K, Affy 250K	BeadStudio (Affymetrix)	pHWE<1e-6, call rate<98%, MAF<1%, gender mismatch, excess heterozygosity	487,573	MACH v1.0.15	HapMap rel. 22 (build 36)	none	R, GenABEL, ProbABEL (linear mixed effect models)
FamHS	Illumina 550K, Illumina 610K, and Illumina 1M	BeadStudio-GenCall v3.0	call rate <98%, pHWE<10e-6, MAF<1%	499,979	MACH v1.0.16	HapMap rel. 22 - phased CEU haplotypes (build 36)	none	R (quantitative traits), SAS (qualitative trait)

Supplementary Table 18 (continued).

Study name	Array type	Genotype calling	QC filters genotyped SNPs used for imputation‡	No. of SNPs used for imputation	Imputation software	Imputation Backbone (NCBI build)	Filtering of imputed genotypes†	Data management and statistical analysis
FHS	Affymetrix 500K Affymetrix 50K Supplementary	Affymetrix	call rate <95%, pHWE<10e-6	503,526	MACH v1.0.15	HapMap rel. 22 - phased CEU haplotypes (build 36)	none	R
GENOA	Affymetrix 6.0 (primary), Illumina 610- Quad, Illumina 660-Quad, Illumina 1M- Duo	Birdseed (Affymetrix data), Genome Studio (Illumina data)	call rate < 95%, pHWE < 0.001, MAF<1%	1,233,495 (because of the different platforms, some SNPs may have had many missings)	MACH v1.0.16	HapMap CEU rel. 22 (build 36)	none	R, multic and GEE
HABC	Illumina 1M	BeadStudio v3.3.7	MAF<1%, call rate < 97%, pHWE<1e-6	914,263	MACH v1.0.16	HapMap CEU rel. 22 (build 36)	None	R, linear regression and logistic regression
HCS	Illumina 610K- Quad	Illumina	call rate <95%, pHWE<10e-6, MAF<1%	513,977	MACH v1.0.16	HapMap rel. 24 - phased CEU haplotypes (build 36.1)	MAF<0.01, Rsqr<0.3	plink, SAS
HPFS	Affymetrix Genome-Wide Human 6.0 array	Birdseed	call rate <97%, pHWE<10e-4, MAF <0.02, more than 1 discordance over 29 replicates, significant plate associations	607,569 (autosomal)	MACH v1.0.15	HapMap rel. 22 - phased CEU haplotypes (build 36)	none	ProbABEL (R), SAS 9.0, PLINK
HYPERGEN ES HTN cases	Illumina 1M Duo	Illumina	call rate <99%, pHWE<10e-8	882,935	MACH v1.0	HapMap rel. 22 - phased CEU haplotypes (build 36)	Rsqr<0.3	PLINK, R
HYPERGEN ES HTN ctrls	Illumina 1M Duo	Illumina	call rate <99%, pHWE<10e-8	882,935	MACH v1.0	HapMap rel. 22 - phased CEU haplotypes (build 36)	Rsqr<0.3	PLINK, R
INCIPE	Illumina	Illumina	call rate <95%, pHWE<10e-6	635,646	IMPUTE 2	HapMap rel. 22 - phased CEU haplotypes (build 36)	none	R
INGI- CARLANTINO	Illumina 370K	BeadStudio	MAF<5%, call rate < 90%, pHWE < 0.05	374,498	MACH	HapMap rel. 22 (build 36)	Rsqr<0.3 and <5 copies of the rare allele.	R, GenABEL, mmscore
INGI- CILENTO	370 K Illumina	Illumina	call rate < 95%, SNPs not in Hapmap	285,674	MACH v1.0.15	HapMap rel. 22 (build 36)	none	R, linear model, GenABEL, ProbABEL (mmscore)
INGI-FVG	Illumina 370K	BeadStudio	MAF<5%, call rate < 90%, pHWE < 0.05	374,498	MACH	HapMap rel. 22 (build 36)	Rsqr<0.3 and <5 copies of the rare allele.	R, GenABEL, mmscore

Supplementary Table 18 (continued).

Study name	Array type	Genotype calling	QC filters genotyped SNPs used for imputation‡	No. of SNPs used for imputation	Imputation software	Imputation Backbone (NCBI build)	Filtering of imputed genotypes†	Data management and statistical analysis
INGI-VAL BORBERA	Illumina SNP array 370K - HumanCNV370 -Quadv3	BeadStudio	pHWE<1e-4, call rate < 90%, MAF<1%	324,319	MACH	HapMap rel. 22 (build 36)	none	R
JUPITER	Omni 1M Quad	Illumina	sample call rate <98%, call rate < 90%, pHWE<1e-6	740,416	MACH v1.0	1000 Genomes - CEU - Pilot - Jun 2010	none	R, probABEL
KORA-F3	Affymetrix 500K	BRLMM	per-chip call rate <93%, MAF <5%,discrepancy for one of the 50 SNPs common on both chips, gender mismatch	380,407	MACH	HapMap rel. 22 (build 35)	none	MACH2QTL, PROBABEL, R, VISUAL BASIC
KORA-F4	Affymetrix 6.0	BRLMM	per-chip call rate <93%, per SNP call rate <93%, MAF<1%, gender mismatch	629,893	MACH	HapMap rel. 22 (build 36)	none	MACH2QTL, PROBABEL, R, VISUAL BASIC
MESA	Affymetrix Genome-Wide Human SNP Array 6.0	Birdseed v2	call rate < 95%, MAF≤1%	897,979	IMPUTE v2.1.0	HapMap rel. 22 - phased CEU haplotypes (build 36)	none	PLINK
MICROS	Illumina Infinium HumanHap300 v2 SNP bead microarrays	BeadStudio	call rate <98%, MAF<1%, pHWE<10e-6	292,917	MACH v1.0.16	HapMap rel. 22 (build 36)	none	R, GenABEL, ProbABEL;
NESDA	Perlegen 600K	Perlegen	call rate <95%, MAF<1%, >5% genotype mismatches, >5% Mendelian errors, unknown SNP location	435,291	IMPUTE v0.4.2	HapMap rel. 24 - phased CEU haplotypes (build 36)	none	SNPTEST v2.2.0, R
NHS	Affymetrix Genome-Wide Human 6.0 array	Birdseed	call rate <97%, pHWE<10e-4, MAF <2%, >1 discordance/12 replicates, significant plate associations	606,626 (autosomal)	MACH v1.0.15	HapMap rel. 22 - phased CEU haplotypes (build 36)	none	R, ProbABEL, SAS 9.0, PLINK
NSPHS	Illumina 300K	BeadStudio	MAF≤1%, pHWE≤1e-5 , call rate ≤ 97%	318,049	MACH v1.0.15	HapMap rel. 22 (build 36)	none	R, GenABEL, ProbABEL
OGP-TALANA	Affymetrix500k	Affymetrix	call rate <95%, pHWE<10e-6	329,122	MACH v1.0.16	HapMap rel. 22 - phased CEU haplotypes (build 36)	Rs<0.3	R, GenABEL, ProbABEL (mmscore)
ORCADES	Illumina 300K	BeadStudio	MAF≤1%, pHWE ≤ 1e-6, call-rate ≤ 98%	306,207	MACH v1.0.15	HapMap rel. 22 (build 36)	none	R, GenABEL, ProbABEL

Supplementary Table 18 (continued).

Study name	Array type	Genotype calling	QC filters genotyped SNPs used for imputation‡	No. of SNPs used for imputation	Imputation software	Imputation Backbone (NCBI build)	Filtering of imputed genotypes†	Data management and statistical analysis
POPGEN	Affymetrix 6.0	Birdseed v2	sample call rate < 0.90, call rate < 0.95, pHWE<1e-4, MAF<1%	709,003	MACH v1.0.16	HapMap rel. 22 - phased CEU haplotypes (build 36)	none	PLINK, R
PREVEND	Illumina CytoSNP12 v2	GenomeStudio	call rate < 95%, pHWE<1e-5	232,571		HapMap rel. 22 - phased CEU haplotypes (build 36)	none	STATA
PROSPER-PHASE	Illumina 660K	Illumina	call rate < 97,5%	557	MACH v1.0.15	HapMap rel. 22 (build 36)	none	R
RS-I	Version 3 Illumina Infinium II HumanHap550	BeadStudio	pHWE<1e-5, call rate<90%, MAF<1%, Mendelian errors > 100, SNPs not in Hapmap or strandedness issues merging with Hapmap	491,875	MACH	HapMap rel. 22 (build 36)	none	ProbABEL
RS-II	Version 3 Illumina Infinium II HumanHap550	BeadStudio	pHWE<1e-5, call rate<90%, MAF<1%, Mendelian errors>100, SNPs not in Hapmap or strandedness issues merging with Hapmap	495,478	MACH	HapMap rel. 22 (build 36)	none	ProbABEL
SAPALDIA	Illumina Human610-Quad BeadChip	Gencall	none	567,589	MACH v1.0.16	HapMap rel. 22 - phased CEU haplotypes (build 36)	none	ProbABEL, STATA v11
SHIP	Affymetrix 6.0	Affymetrix Birdseed2	none	869,224	IMPUTE v0.5.0	HapMap rel. 22 (build 36)	none	SNPTEST v1.1.5, QUICKTEST v0.94, R, InforSense, InterSystems Caché,
SHIP-TREND	Illumina Human Omni 2.5	Illumina	pHWE ≤ 1e-4, call rate ≤ 0.9, monomorphic SNPs	1,782,967	IMPUTE v2.1.2.3	HapMap rel. 22 - phased CEU haplotypes (build 36)	duplicated rsID with different positions	QUICKTEST v0.95, R, InforSense, InterSystems Caché

Supplementary Table 18 (continued).

Study name	Array type	Genotype calling	QC filters genotyped SNPs used for imputation‡	No. of SNPs used for imputation	Imputation software	Imputation Backbone (NCBI build)	Filtering of imputed genotypes†	Data management and statistical analysis
SORBS	500K Affymetrix GeneChip (250K Sty and 250K Nsp arrays, Affymetrix, Inc) and Affymetrix Genome-Wide Human SNP Array 6.0	Microarray Core Facility of the Interdisciplinary Centre for Clinical Research, University of Leipzig, Germany and ATLAS Biolabs GmbH, Berlin, Germany	MAF<1%, pHWE<1e-4, call rate < 95%,	378,513	IMPUTE	HapMap CEU rel. 21 (build 35)	proper_info ≤ 0.4	ProbABEL with robust variance option to account for remaining relatedness
WGHS	Illumina HumanHap300 Duo "plus"	BeadStudio v3.1	pHWE<1e-6, call rate < 98%, MAF<1%	331,993	MACH v1.0.15	HapMap rel. 22 (build 36)	none	PLINK, R, ProbABEL
YFS	Illumina 670k custom	Illuminus	call rate <95%, pHWE<10e-6	546,674	IMPUTE 2.1.2	1000 Genomes - Jun 2011 phased haplotypes	none	SAS

‡QC filters for genotyped SNPs used for imputation (SNPs satisfying the filter were excluded).

†Filtering of imputed genotypes (SNPs satisfying the filter were excluded).

Supplementary Table 19. Genotyping platform – *in silico* replication studies.

Study Name	Array type	Genotype calling	QC filters for genotyped SNPs used for imputation	No of SNPs used for imputation	Imputation	Imputation Backbone (NCBI build)	Filtering of imputed genotypes	Data management and statistical analysis
EGCUT-replic	Illumina OmniExpress	Genome Studio (GenCall)	call rate<95%, pHWE<10e-6, MAF<1%	616,063	IMPUTE v2.2	HapMap rel. 22 - phased CEU haplotypes (build 36)	none	Plink, SNPTEST 2, R
GENDIAN	Affymetrix Genome-Wide Human SNP Array 6.0	Birdseed (BRLMM)	N=126,259 SNPs (chr 1-chr22, chr X) were excluded from imputation by SNP quality control.	776,075	Mach 1.0.18.c MiniMac 2012-10-09	GIANT ALL 1000 Genomes v3 ref panel GRCh build 37	none	R
GSK cases/controls	Illumina 550 K	Illumina	call rate <98%, pHWE<10e-5	522,008	IMPUTE v2	CEU in HapMap3 and 1000 Genomes Pilot 1	none	R, Plink
Gutenberg Health Study	Affymetrix 6.0	Birdseed	call rate<95%, MAF<1%, pHWE<10e-4	662,405	IMPUTE v2.1.0	HapMap rel. 24 (build 36)	none	R
HRS	Illumina Omni2.5 Beadchip	Genome Studio v2011.2	call rate <98%, pHWE<10e-4	551,936	MACH v1.0.16	HapMap rel. 22 - phased CEU haplotypes (build 36)	none	SAS, R
IPM_EA_Affy	Affymetrix 6.0	Birdseed	sample call rate <95%, SNP call rate <95%, pHWE<1e-4, MAF<1%	711,270	IMPUTE 2	Phase I integrated variant set release (v3) (NCBI build 37)	none	SNPTEST 2, R
IPM_EA_Illu	Illumina Human OmniExpress Exome-8v1	Genome Studio	sample call rate <99%, SNP call rate <95%, pHWE<5e-5, MAF<1%	865,711	IMPUTE 2	Phase I integrated variant set release (v3) (NCBI build 37)	none	SNPTEST 2, R
LURIC	Affymetrix 6.0	Birdseed v2	call rate <98%, pHWE<10e-4	686,195	MACH	HapMap rel. 22 - phased CEU haplotypes (build 36)	none	QUICKTEST
Vanderbilt Omni1	Illumina HumanOmni1-Quad	BeadStudio	call-rate<98%, IBD (ZO<0.8), Mendel errors >0, Duplicate concordance <100%	946,523	IMPUTE v2.3.0	1000 Genomes Phase 1 integrated v3	Genotype Likelihood <0.9	Plink, R
Vanderbilt Omni5	Illumina HumanOmni5-Quad	BeadStudio	call rate <98%, IBD (ZO<0.8), Mendel errors >0, Duplicate concordance <100%	3,819,154	IMPUTE v2.3.0	1000 Genomes Phase 1 integrated v3	Genotype Likelihood <0.9	Plink, R
Vanderbilt 660W	Illumina Human660W-Quad	BeadStudio	call rate <98%, IBD (ZO<0.8), Mendel errors >0, Duplicate concordance <100%	530,014	IMPUTE v2.3.0	1000 Genomes Phase 1 integrated v3	Genotype Likelihood <0.9	Plink, R

Supplementary Table 20. Genotyping information: *de novo* replication studies.

Study Name	Bus Santé	ESTHER	KORA F3 and F4	OGP	SAPHIR	SKIPOGH
Genotyping platform	LGC Genomics SNP-line, using KASP™ Chemistry and 1536-well plates	LGC Genomics SNP-line, using KASP™ Chemistry	MALDITOF MS, Bruker Daltonik GmbH, Leipzig, Germany	LGC Genomics SNP-line, using KASP™ Chemistry	Sequenom platform	LGC Genomics SNP-line, using KASP™ Chemistry and 1536-well plates
Amount of DNA used per SNP (in ng)	DNA amplified, but typically 5 -7.5 ng of gDNA per genotype	3.75 ng	1 ng	5 ng	15 ng	5 -7.5 ng of gDNA per genotype
Genotyping method	See note ¹	See note ¹	iPlex Gold	See note ¹	Mass ARRAY Analyzer 4 system	See note ¹
No. of duplicates and concordance per SNP (provided per individual SNP)	Not duplicated	LGC Genomics does not add duplicates. The data for each SNP represents one reaction per sample.	At least 15% duplicate genotyping per SNP. Concordance ≥ 95%, median = 100%	LGC Genomics does not add duplicates. The data for each SNP represents one reaction per sample.	70 duplicates; of 46 SNPs genotyped, 44 had a concordance of 100%, 2 had 1 discordant sample each	29 samples genotyped in duplicate. SNP concordance varied from 86% to 100%.
Number attempted / number genotyped samples per individual SNP	The genotyping call rate (%) was typically between 0.95-0.97	Call-rate range 0.98-1	NA	Allele call rate 0.99	46 SNPs were genotyped: mean call-rate = 99.3% (min = 98.15%, max = 99.65%)	Median SNP call-rate = 97.2% (min = 94.5%, max = 99.5%)
Other QC laboratory-specific indices	See note ²	none indicated by the lab	NA	none indicated by the lab	automatic calculation of the HWE, comparison of the obtained genotypes with HapMap Data	See note ²

¹ Genotyped using KASPar (Kompetitive Allele Specific PCR) v4.0 after whole genome amplification by primer extension preamplification (PEP) using thermostable DNA polymerases.

² All assays have been validated on an in-house DNA panel (44 random Caucasian DNA samples). All sample plates genotyped include at least two negative controls. ie. blank/water controls. All genotyping data is initially generated by an automated algorithm (genotype calling based upon recorded fluorescence values). All genotyping data is manually checked and verified by no less than two experienced scientists at LGC genomics.

Supplementary Table 21. Morpholino (MO) sequences.

Target gene	Splice or ATG	MO sequence
<i>A1cf</i>	ATG	5' CCCCACATTTTTGATTGGTTTCCAT 3'
<i>Dpep</i>	ATG	5' AATCTTGACCCATTCCATCATCACC 3'
<i>Kbtbd2</i>	ATG	5' TGTTCGTATCCCATGAGTTTTCAAC 3'
<i>Nfatc1</i>	Splice	5' CGCATCTGTAAGGTACAATCACATT 3'
<i>Nfkb1</i>	ATG	5' TCCTCGCCAGCCATGATTCCTTTGC 3'
<i>Ptpro</i>	ATG	5' GATCGCACTCTTTGATTCTCGGCGT 3'
<i>Skila</i>	Splice	5' TTGCCCTGCAAACACACATACACAC 3'
<i>Skilb</i>	Splice	5' CCCGGATGACTGAAACAAGTCAAAA 3'
<i>Sypl2a</i>	Splice	5' TTGAAATGTGGTTGTTTATACCTGA 3'
<i>Sypl2b</i>	Splice	5' AGACTCTTTAATGAGGTTTACCTAA 3'
<i>Tspan9a</i>	Splice	5' GTAGGAGTGGCAAACTTACGCTCA 3'
<i>Tspan9b</i>	ATG	5' GCACAGGCATCCACGAGCCATCTTC 3'
<i>Uncx</i>	Splice	5' ATCCCCCGAATCTATGTAAGAAACA 3'
<i>Wnt7aa</i>	ATG	5' TCCAGCGGCGCGTTTTCTGCTCAT 3'

Supplementary Note.

Study specific acknowledgements and funding sources for participating studies, alphabetical order.

3C. Three-City Study. The work was made possible by the participation of the control subjects, the patients, and their families. We thank Dr. Anne Boland (CNG) for her technical help in preparing the DNA samples for analyses. This work was supported by the National Foundation for Alzheimer's disease and related disorders, the Institut Pasteur de Lille and the Centre National de Génotypage. The 3C Study was performed as part of a collaboration between the Institut National de la Santé et de la Recherche Médicale (Inserm), the Victor Segalen Bordeaux II University and Sanofi-Synthélabo. The Fondation pour la Recherche Médicale funded the preparation and initiation of the study. The 3C Study was also funded by the Caisse Nationale Maladie des Travailleurs Salariés, Direction Générale de la Santé, MGEN, Institut de la Longévité, Agence Française de Sécurité Sanitaire des Produits de Santé, the Aquitaine and Bourgogne Regional Councils, Fondation de France and the joint French Ministry of Research/INSERM "Cohortes et collections de données biologiques" programme. Lille Génopôle received an unconditional grant from Eisai.

Advance. Action in Diabetes and Vascular disease: preteraX and diamicron mr Controlled Evaluation study. The genetic epidemiological work was funded by Prognomix Inc. and by grants from Genome Quebec and Canadian Institutes for Health Research (CIHR). The clinical study was managed by the George Institute for International Health (Sydney, Australia) with grants received from Les Laboratoires Servier, France and from Medical Research Council of Australia. The genotyping was performed at the genomic platform of CRCHUM. The authors acknowledge the technical help of Carole Long and Mounif Haloui and the bioinformatic analyses performed by Gilles Godefroid, François-Christophe Blanchet-Marois and François Harvey. The members of the genetic sub-study of ADVANCE, Stephen Harrap and Michel Marre are also acknowledged.

AGES. Age, Gene/Environment Susceptibility-Reykjavik Study. This study has been funded by NIH contract N01-AG-1-2100, the NIA Intramural Research Program, Hjartavernd (the Icelandic Heart Association), and the Althingi (the Icelandic Parliament). The study is approved by the Icelandic National Bioethics Committee, VSN: 00-063. The researchers are indebted to the participants for their willingness to participate in the study.

Amish. Amish studies. We thank our Amish research volunteers for their long-standing partnership in research, and the research staff at the Amish Research Clinic for their work. We are supported by grants and contracts from the NIH including R01 AG18728 (Amish Longevity Study), R01 HL088119 (Amish Calcification Study), U01 GM074518-04 (PAPI Study), U01 HL072515-06 (HAPI Study), U01 HL084756 and NIH K12RR023250 (University of Maryland MCRDP), the University of Maryland General Clinical Research Center, grant M01 RR 16500, the Baltimore Veterans Administration Medical Center Geriatrics Research and Education Clinical Center and the Paul Beeson Physician Faculty Scholars in Aging Program.

ARIC. Atherosclerosis Risk in Communities study. The ARIC study is carried out as a collaborative study supported by National Heart, Lung, and Blood Institute contracts (HHSN268201100005C, HHSN268201100006C, HHSN268201100007C, HHSN268201100008C, HHSN268201100009C, HHSN268201100010C, HHSN268201100011C, and HHSN268201100012C), R01HL087641, R01HL59367 and R01HL086694; National Human Genome Research Institute contract U01HG004402; and National Institutes of Health contract HHSN268200625226C. The authors thank the staff and participants of the ARIC study for their important contributions. Infrastructure was partly supported by Grant Number UL1RR025005, a component of the National Institutes of Health and NIH Roadmap for Medical Research. This work as well as YL and AK were supported by the Emmy Noether Program of the German Research Foundation (KO 3598/2-1 to AK).

ASPS. Austrian Stroke Prevention Study. The research reported in this article was funded by the Austrian Science Fond (FWF) grant number P20545-P05 and P13180. The Medical University of Graz supports the databank of the ASPS. The authors thank the staff and the participants of the ASPS for their valuable contributions. We thank Birgit Reinhart for her long-term administrative commitment and Ing Johann Semmler for the technical assistance at creating the DNA-bank.

AUSTWIN. The *Australian Twin-Family Studies* were supported by NIH grants AA07535, AA07728, AA13320, AA13321, AA13326, AA14041, AA11998, AA17688, DA012854, DA019951; by grants from the Australian National Health and Medical Research Council (241944, 339462, 389927, 389875, 389891, 389892, 389938, 442915, 442981, 496739, 552485, 552498); from the Australian Research Council (A7960034, A79906588, A79801419, DP0770096, DP0212016, DP0343921); and the FP-5 GenomEUtwin Project (QLG2-CT-2002-01254). Genotyping was partially supported by grant AA13320 to the late Richard Todd, PhD, MD. R.P.M., and G.W.M. are supported by the National Health and Medical Research Council (NHMRC) Fellowship Scheme.

BLSA. *Baltimore Longitudinal Study of Aging.* The BLSA was supported in part by the Intramural Research Program of the NIH, National Institute on Aging.

BMES. *Blue Mountains Eye Study.* The BMES has been supported by the Australian RADGAC grant (1992-94) and Australian National Health & Medical Research Council, Canberra Australia (Grant Nos: 974159, 211069, 991407, 457349). The GWAS studies of Blue Mountains Eye Study population are supported by the Australian National Health & Medical Research Council (Grant Nos: 512423, 475604, 529912) and the Wellcome Trust, UK (2008). EGH and JJW are funded by the Australian National Health & Medical Research Council Fellowship Schemes.

Bus Santé study. This research was conducted in part using data and resources from the Bus Santé study of the Geneva University Hospitals. This work was partially supported by the Geneva University Hospitals, the Canton of Geneva (Switzerland), the General Directorate of Health (Canton of Geneva), the Swiss School of Public Health Plus (SSPH+) and the Swiss Foundation for Science (Contract No.495 33CM30-124087). Data on creatinine was founded by the Loterie Romande. The investigators of the Bus Santé study thank all the collaborators of the Unit of Population Epidemiology, and Abbott Diagnostics (Baar, Switzerland). We thank Dr Olivier Golaz and Dr Pierre Lescuyer, Geneva University Hospitals.

CARDIA. *Coronary Artery Risk in Young Adults.* University of Alabama at Birmingham (N01-HC-48047), University of Minnesota (N01-HC-48048), Northwestern University (N01-HC-48049), Kaiser Foundation Research Institute (N01-HC-48050), University of Alabama at Birmingham (N01-HC-95095), Tufts-New England Medical Center (N01-HC-45204), Wake Forest University (N01-HC-45205), Harbor-UCLA Research and Education Institute (N01-HC-05187), University of California, Irvine (N01-HC-45134, N01-HC-95100).

CHS. *Cardiovascular Health Study.* The CHS research reported in this article was supported by contract numbers N01-HC-85079 through N01-HC-85086, N01-HC-35129, N01 HC-15103, N01 HC-55222, N01-HC-75150, N01-HC-45133, grant numbers U01 HL080295 and R01 HL087652 from the National Heart, Lung, and Blood Institute, with additional contribution from the National Institute of Neurological Disorders and Stroke. A full list of principal CHS investigators and institutions can be found at <http://www.chs-nhlbi.org/pi.htm>. DNA handling and genotyping was supported in part by National Center for Research Resources grant M01RR00425 to the Cedars-Sinai General Clinical Research Center Genotyping core and National Institute of Diabetes and Digestive and Kidney Diseases grant DK063491 to the Southern California Diabetes Endocrinology Research Center.

CROATIA-KORCULA. The Korcula study in the Croatian island of Vis was supported through the grants from the Medical Research Council UK to H.C., A.F.W. and I.R.; and Ministry of Science, Education and Sport of the Republic of Croatia to I.R. (number 108-1080315-0302). We would like to acknowledge the contributions of the recruitment team in Korcula, the administrative teams in Croatia and Edinburgh (Rosa Bisset) and the people of Korcula.

CROATIA-SPLIT. The CROATIA-Split study was supported through the grants from the Medical Research Council UK; and Ministry of Science, Education and Sport of the Republic of Croatia. (number 108-1080315-0302). We would like to acknowledge the invaluable contributions of the recruitment team from the Croatian Centre for Global Health, University of Split, the administrative teams in Croatia and Edinburgh and the people of Split. The SNP genotyping for the SPLIT cohort was performed by AROS Applied Biotechnology, Aarhus, Denmark.

CROATIA-VIS. The Vis study in the Croatian island of Vis was supported through the grants from the Medical Research Council UK to H.C., A.F.W. and I.R.; and Ministry of Science, Education and Sport of the Republic of Croatia to I.R. (number 108-1080315-0302) and the European Union framework program 6 EUROSPAN project (contract no. LSHG-CT-2006-018947). We would like to acknowledge the invaluable contributions of the recruitment

team (including those from the Institute of Anthropological Research in Zagreb) in Vis, the administrative teams in Croatia and Edinburgh (Rosa Bisset) and the people of Vis.

DESIR. *Data from an Epidemiological Study on the Insulin Resistance syndrome.* DESIR genetic data was supported in part by the "Conseil Regional Nord-Pas-de-Calais: Fonds européen de développement économique et régional (CPER 2011-2013); Genome Quebec-Genome Canada; and the British Medical Research Council.

EGCUT. *Estonian Genome Center University of Tartu.* EGCUT received financing from FP7 grants (278913, 306031, 313010) and targeted financing from Estonian Government (SF0180142s08). EGCUT studies were covered from Infra-structure grant no. 3.2.0304.11-0312 funded mostly by the European Regional Development Fund, Center of Excellence in Genomics (EXCEGEN) and University of Tartu (SP1GVARENG). We acknowledge EGCUT technical personnel, especially Mr V. Soo and S. Smit. Data analyses were carried out in part in the High Performance Computing Center of the University of Tartu.

ERF. *Erasmus Rucphen Family study.* The ERF study was supported by grants from the Netherlands Organization for Scientific Research (NWO; Pioneergrant), Erasmus Medical Center, the Centre for Medical Systems Biology (CMSB), and the Netherlands Kidney Foundation. We are grateful to all patients and their relatives, general practitioners and neurologists for their contributions and to P. Veraart for her help in genealogy, Jeannette Vergeer for the supervision of the laboratory work and P. Sniijders for his help in data collection.

ESTHER. *Epidemiologische Studie zu Chancen der Verhütung, Früherkennung und optimierten Therapie chronischer ERkrankungen in der älteren Bevölkerung.* We thank all the individuals who took part in this study and all the researchers, clinicians, technicians and administrative staff who have enabled this work to be carried out. This work was supported in part by the Baden-Württemberg State Ministry of Science, Research and Arts; by the German Federal Ministry of Education and Research (grant number 01ET0717); and by the CHANCES project funded in the Seventh Framework Programme of the Directorate-General for Research and Innovation in the European Commission (grant number 242244).

FamHS. *Family Heart Study.* The FHS work was supported in part by NIH grants 5R01HL08770003, 5R01HL08821502 (Michael A. Province) from the NHLBI and 5R01DK07568102, 5R01DK06833603 from the NIDDK (I.B.B.). The authors thank the staff and participants of the FamHS for their important contributions.

FHS. *Framingham Heart Study.* This research was conducted in part using data and resources from the Framingham Heart Study of the National Heart Lung and Blood Institute of the National Institutes of Health and Boston University School of Medicine. The analyses reflect intellectual input and resource development from the Framingham Heart Study investigators participating in the SNP Health Association Resource (SHARe) project. This work was partially supported by the National Heart, Lung and Blood Institute's Framingham Heart Study (Contract No. N01-HC-25195) and its contract with Affymetrix, Inc for genotyping services (Contract No. N02-HL-6-4278). A portion of this research utilized the Linux Cluster for Genetic Analysis (LinGA-II) funded by the Robert Dawson Evans Endowment of the Department of Medicine at Boston University School of Medicine and Boston Medical Center.

GENDIAN. *GENetics of DIAbetic Nephropathy study.* The support of the physicians, the patients, and the staff of the Diabetes Zentrum Mergentheim (Head: Prof. Dr. Thomas Haak), the diabetes outpatient clinic Dr Nusser - Dr Kreisel, the dialysis centers KfH Amberg, KfH Bayreuth, KfH Deggendorf, KfH Donauwörth, KfH Freising, KfH Freyung, KfH Fürth, KfH Hof, KfH Ingolstadt, KfH Kelheim, KfH München Elsenheimerstraße, KfH München-Schwabing, KfH Neumarkt, KfH Neusäß, KfH Oberschleißheim, KfH Passau, KfH Plauen, KfH Regensburg Günzstraße, KfH Regensburg Caritas-Krankenhaus, KfH Straubing, KfH Sulzbach-Rosenberg, KfH Weiden, Dialysezentrum Augsburg Dr. Kirschner, Dialysezentrum Bad Alexandersbad, KfH Bamberg, Dialysezentrum Emmering, Dialysezentrum Klinikum Landshut, Dialysezentrum Landshut, Dialysezentrum Pfarrkirchen, Dialysezentrum Schwandorf, Dr. Angela Götz, the medical doctoral student Johanna Christ and the Study Nurse Ingrid Lugauer. The expert technical assistance of Claudia Strohmeier is acknowledged. Phenotyping was funded by the Dr. Robert Pfleger-Stiftung (Dr Carsten A. Böger), the MSD Stipend Diabetes (Dr Carsten A. Böger) and the University Hospital of Regensburg (intramural grant ReForM A to Dr. A. Götz, ReForM C to Dr. Carsten Böger). Genome-wide genotyping was funded by the KfH Stiftung Präventivmedizin e.V. (Dr. Carsten A. Böger, Dr. Jens Brüning), the Else Kröner-Fresenius-Stiftung (2012_A147 to Dr Carsten A. Böger and Dr Iris M. Heid) and the University Hospital Regensburg (Dr Carsten A. Böger). Data analysis was funded by the Else Kröner-Fresenius Stiftung (Dr. Iris M. Heid and Dr. Carsten A. Böger: 2012_A147; Dr. Carsten A. Böger and Dr. Bernhard K. Krämer: P48/08//A11/08). GENDIAN Study Group: Mathias Gorski, Iris M. Heid, Bernhard K. Krämer, Myriam Rheinberger, Michael Broll, Alexander Lammert, Jens

Brüning, Matthias Olden, Klaus Stark, Claudia Strohmeier, Simone Neumeier, Sarah Hufnagel, Petra Jackermeier, Emilia Ruff, Johanna Christ, Peter Nürnberg, Thomas Haak, Carsten A. Böger.

GENOA. *Genetic Epidemiology Network Of Arteriopathy.* This research was partially supported by the National Heart Lung and Blood Institute of the National Institutes of Health R01 HL-87660.

GHS. *Gutenberg Heart Study.* The Gutenberg Health study is funded through the government of Rheinland-Pfalz (No. AZ 961-386261/733), the research program “Wissen schafft Zukunft” and the “Schwerpunkt Vaskuläre Prävention” of the Johannes Gutenberg University of Mainz and its contract with Boehringer Ingelheim and PHILIPS medical systems, including an unrestricted grant for the Gutenberg Health Study. Further, this project has been supported by the National Genome Network “NGFNplus” by the federal Ministry of Education and Research, Germany (No. 01GS0833 and 01GS0831, projects A3/D1).

GSK cases/controls. Funding Source: Max-Planck Society, German Federal Ministry of Education and Research (BMBF) in the framework of the National Genome Research Network (NGFN), Foerderkennzeichen 01GS0481.

HABC. *Health Aging and Body Composition Study.* The HABC study was funded by the National Institutes of Aging. This research was supported by NIA contracts N01AG62101, N01AG62103, and N01AG62106. The genome-wide association study was funded by NIA grant 1R01AG032098-01A1 to Wake Forest University Health Sciences and genotyping services were provided by the Center for Inherited Disease Research (CIDR). CIDR is fully funded through a federal contract from the National Institutes of Health to The Johns Hopkins University, contract number HHSN268200782096C. This research was supported in part by the Intramural Research Program of the NIH, National Institute on Aging.

HANDLS. This research was supported by the Intramural Research Program of the NIH, National Institute on Aging and the National Center on Minority Health and Health Disparities (Intramural Project # Z01-AG000513 and human subjects protocol # 2009-149). Data analyses for the HANDLS study utilized the high-performance computational capabilities of the Biowulf Linux cluster at the National Institutes of Health, Bethesda, Md. (<http://biowulf.nih.gov>).

HCS. *Hunter Community Study.* The University of Newcastle provided \$300,000 from its Strategic Initiatives Fund, and \$600,000 from the Gladys M Brawn Senior Research Fellowship scheme; Vincent Fairfax Family Foundation, a private philanthropic trust, provided \$195,000; The Hunter Medical Research Institute provided media support during the initial recruitment of participants; and Dr Anne Crotty, Prof. Rodney Scott and Associate Prof. Levi provided financial support towards freezing costs for the long-term storage of participant blood samples. The authors would like to thank the men and women participating in the HCS as well as all the staff, investigators and collaborators who have supported or been involved in the project to date. A special thank you should go to Alison Koschel and Debbie Quain who were instrumental in setting up the pilot study and initial phase of the project.

HPFS. *Health Professionals Follow-Up Study.* The NHS/HPFS type 2 diabetes GWAS (U01HG004399) is a component of a collaborative project that includes 13 other GWAS (U01HG004738, U01HG004422, U01HG004402, U01HG004729, U01HG004726, U01HG004735, U01HG004415, U01HG004436, U01HG004423, U01HG004728, RFAHG006033; National Institute of Dental & Craniofacial Research: U01DE018993, U01DE018903) funded as part of the Gene Environment-Association Studies (GENEVA) under the NIH Genes, Environment and Health Initiative (GEI). Assistance with phenotype harmonization and genotype cleaning, as well as with general study coordination, was provided by the GENEVA Coordinating Center (U01HG004446). Assistance with data cleaning was provided by the National Center for Biotechnology Information. Genotyping was performed at the Broad Institute of MIT and Harvard, with funding support from the NIH GEI (U01HG04424), and Johns Hopkins University Center for Inherited Disease Research, with support from the NIH GEI (U01HG004438) and the NIH contract “High throughput genotyping for studying the genetic contributions to human disease”(HHSN268200782096C). Additional funding for the current research was provided by the National Cancer Institute (P01CA087969, P01CA055075), and the National Institute of Diabetes and Digestive and Kidney Diseases (R01DK058845). We thank the staff and participants of the NHS and HPFS for their dedication and commitment.

HUFS. The *Howard University Family Study* was supported by National Institutes of Health grants S06GM008016-320107 and S06GM008016-380111. Enrollment was carried out at the Howard University General Clinical Research Center, supported by National Institutes of Health grant 2M01RR010284. This research was supported in part by the

Intramural Research Program of the Center for Research on Genomics and Global Health. The Center for Research on Genomics and Global Health is supported by the National Human Genome Research Institute, the National Institute of Diabetes and Digestive and Kidney Diseases, the Center for Information Technology, and the Office of the Director at the National Institutes of Health (Z01HG200362). Genotyping support was provided by the Coriell Institute for Medical Research. We thank the participants of the Howard University Family Study. The contents of this publication are solely the responsibility of the authors and do not necessarily represent the official view of the National Institutes of Health.

HYPERGENES. *European Network for Genetic-Epidemiological Studies: building a method to dissect complex genetic traits, using essential hypertension as a disease model.* HYPERGENES (FP7 - HEALTH-F4-2007-201550); INTEROMICS (MIUR - CNR Italian Flagship Project); IC15-CT98-0329-EPOGH, LSHM-CT-2006-037093; HEALTH-2011-278249-EU-MASCARA, and ERC Advanced Grant-2011-294713-EPLORE and the Fonds voor Wetenschappelijk Onderzoek Vlaanderen, Ministry of the Flemish Community, Brussels, Belgium (grants G.0575.06 and G.0734.09).

Patricia B. Munroe's work forms part of the research themes contributing to the translational research portfolio for the NIHR Barts Cardiovascular Biomedical Research Unit.

INCIPE. The INCIPE study was co-sponsored by Fondazione Cassa di Risparmio di Verona, Azienda Ospedaliera di Verona, and University of Verona, Italy.

INGI-CARLANTINO. *Italian Network on Genetic Isolates – Carlantino.* We thank Anna Morgan and Angela D'Eustacchio for technical support. We are grateful to the municipal administrators for their collaboration on the project and for logistic support. We would like to thank all participants to this study.

INGI-CILENTO. *Italian Network on Genetic Isolates – Cilento.* We thank the populations of Cilento for their participation in the study. This work was supported by grants from the EU (Vasoplus-037254), the Italian Ministry of Universities (FIRB -RBIN064YAT), the Assessorato Ricerca Regione Campania, the Ente Parco Nazionale del Cilento e Vallo di Diano and the Fondazione Banco di Napoli to MC.

INGI-FVG. *Italian Network on Genetic Isolates – Friuli Venezia-Giulia.* We thank Anna Morgan and Angela D'Eustacchio for technical support. We are grateful to the municipal administrators for their collaboration on the project and for logistic support. We would like to thank all participants to this study.

INGI-VAL BORBERA. *Italian Network on Genetic Isolates – Val Borbera.* The research was done using data obtained thanks to funds from Compagnia di San Paolo, Torino, Italy, Cariplo Foundation, Milano, Italy and Italian Ministry of Health Progetto Finalizzato 2007 and 2009.

IPM. *Mount Sinai BioMe Biobank Program.* The Mount Sinai BioMe Biobank Program is supported by The Andrea and Charles Bronfman Philanthropies.

JHS. *Jackson Heart Study.* Jackson State University (N01-HC-95170), University of Mississippi (N01-HC-95171), Tougaloo College (N01-HC-95172).

JUPITER. We thank Jean MacFadyen, Lynda Rose, the JUPITER study participants and the >1000 physicians worldwide for their personal time, effort and commitment to JUPITER. This work was supported by research funds from AstraZeneca to Drs. Ridker and Chasman.

HRS. *Health and Retirement Study.* HRS is supported by the National Institute on Aging (NIA U01AG009740). The genotyping was funded as a separate award from the National Institute on Aging (RC2 AG036495). Our genotyping was conducted by the NIH Center for Inherited Disease Research (CIDR) at Johns Hopkins University. Genotyping quality control and final preparation of the data were performed by the Genetics Coordinating Center at the University of Washington.

KORA F3 and F4. The genetic epidemiological work was funded by the NIH subcontract from the Children's Hospital, Boston, US, (H.E.W., I.M.H, prime grant 1 R01 DK075787-01A1), the German National Genome Research Net NGFN2 and NGFNplus (H.E.W. 01GS0823; WK project A3, number 01GS0834), the Munich Center of Health

Sciences (MC Health) as part of LMUinnovativ, and by the Else Kröner-Fresenius-Stiftung (P48/08//A11/08; C.A.B., B.K.K; 2012_A147 to CAB and IMH.). The kidney parameter measurements in F3 were funded by the Else Kröner-Fresenius-Stiftung (C.A.B., B.K.K.) and the Regensburg University Medical Center, Germany; in F4 by the University of Ulm, Germany (W.K.). Genome wide genotyping costs in F3 and F4 were in part funded by the Else Kröner-Fresenius-Stiftung (C.A.B., B.K.K.). Denovo genotyping in F3 and F4 were funded by the Else Kröner-Fresenius-Stiftung (C.A.B., B.K.K.). The KORA research platform and the MONICA Augsburg studies were initiated and financed by the Helmholtz Zentrum München, German Research Center for Environmental Health, by the German Federal Ministry of Education and Research and by the State of Bavaria. Genotyping was performed in the Genome Analysis Center (GAC) of the Helmholtz Zentrum München. The LINUX platform for computation were funded by the University of Regensburg for the Department of Epidemiology and Preventive Medicine at the Regensburg University Medical Center.

LURIC. *Ludwigshafen Risk and Cardiovascular Health Study.* We extend our appreciation to the participants of the LURIC study; without their collaboration, this article would not have been written. We thank the LURIC study team who were either temporarily or permanently involved in patient recruitment as well as sample and data handling, in addition to the laboratory staff at the Ludwigshafen General Hospital and the Universities of Freiburg and Ulm, Germany. LURIC has received funding from the 6th Framework Program (integrated project Bloodomics, grant LSHM-CT-2004-503485) and 7th Framework Program (Atheroremo, grant agreement number 201668 and RiskyCAD, grant agreement number 305739) of the European Union.

MESA. *Multi-Ethnic Study of Atherosclerosis.* University of Washington (N01-HC-95159), Regents of the University of California (N01-HC-95160), Columbia University (N01-HC-95161), Johns Hopkins University (N01-HC-95162, N01-HC-95168), University of Minnesota (N01-HC-95163), Northwestern University (N01-HC-95164), Wake Forest University (N01-HC-95165), University of Vermont (N01-HC-95166), New England Medical Center (N01-HC-95167), Harbor-UCLA Research and Education Institute (N01-HC-95169), Cedars-Sinai Medical Center (R01-HL-071205), University of Virginia (subcontract to R01-HL-071205)

MICROS. *Microisolates in South Tyrol study.* We owe a debt of gratitude to all participants. We thank the primary care practitioners Raffaella Stocker, Stefan Waldner, Toni Pizzecco, Josef Plangger, Ugo Marcadent and the personnel of the Hospital of Silandro (Department of Laboratory Medicine) for their participation and collaboration in the research project. We thank Dr. Peter Riegler (Hemodialysis Unit, Hospital of Merano) for the important discussions. In South Tyrol, the study was supported by the Ministry of Health and Department of Educational Assistance, University and Research of the Autonomous Province of Bolzano, the South Tyrolean Sparkasse Foundation, and the European Union framework program 6 EUROSPAN project (contract no. LSHG-CT-2006-018947).

NESDA. *The Netherlands Study of Depression and Anxiety.* The infrastructure for the NESDA is funded through the Geestkracht programme of the Dutch Scientific Organization (ZON-MW, grant number 10-000-1002) and matching funds from participating universities and mental health care organizations. Genotyping in NESDA was funded by the Genetic Association Information Network (GAIN) of the Foundation for the US National Institutes of Health. Statistical analyses were carried out on the Genetic Cluster Computer (<http://www.geneticcluster.org>), which is financially supported by the Netherlands Scientific Organization (I 480-05-003) along with a supplement from the Dutch Brain Foundation.

NHS. *Nurses' Health Study.* The NHS/HPFS type 2 diabetes GWAS (U01HG004399) is a component of a collaborative project that includes 13 other GWAS (U01HG004738, U01HG004422, U01HG004402, U01HG004729, U01HG004726, U01HG004735, U01HG004415, U01HG004436, U01HG004423, U01HG004728, RFAHG006033; National Institute of Dental & Craniofacial Research: U01DE018993, U01DE018903) funded as part of the Gene Environment-Association Studies (GENEVA) under the NIH Genes, Environment and Health Initiative (GEI). Assistance with phenotype harmonization and genotype cleaning, as well as with general study coordination, was provided by the GENEVA Coordinating Center (U01HG004446). Assistance with data cleaning was provided by the National Center for Biotechnology Information. Genotyping was performed at the Broad Institute of MIT and Harvard, with funding support from the NIH GEI (U01HG04424), and Johns Hopkins University Center for Inherited Disease Research, with support from the NIH GEI (U01HG004438) and the NIH contract "High throughput genotyping for studying the genetic contributions to human disease"(HHSN268200782096C). The NHS renal function and

albuminuria work was supported by DK66574. Additional funding for the current research was provided by the National Cancer Institute (P01CA087969, P01CA055075), and the National Institute of Diabetes and Digestive and Kidney Diseases (R01DK058845). We thank the staff and participants of the NHS and HPFS for their dedication and commitment.

NSPHS. *The Northern Swedish Population Health Study.* The NSPHS was supported by grants from the Swedish Natural Sciences Research Council, the European Union through the EUROSPAN project (contract no. LSHG-CT-2006-018947), the Foundation for Strategic Research (SSF) and the Linneaus Centre for Bioinformatics (LCB). We are also grateful for the contribution of samples from the Medical Biobank in Umeå and for the contribution of the district nurse Svea Hennix in the Karesuando study.

OGP (*Ogliastro Genetic Park*) including **OGP-TALANA**. We thank the Ogliastro population and all the individuals who participated in this study. We are grateful to the municipal administrators for their collaboration to the project and for economic and logistic support. This work was supported by grants from the Italian Ministry of Education, University and Research (MIUR) no.5571/DSPAR/2002 and (FIRB) D. M. no. 718/Ric/2005.

ORCADES. *Orkney Complex Disease Study.* ORCADES was supported by the the Chief Scientist Office of the Scottish Government , the Royal Society and the European Union framework program 6 EUROSPAN project (contract no. LSHG-CT-2006-018947). DNA extractions were performed at the Wellcome Trust Clinical Research Facility in Edinburgh. We would like to acknowledge the invaluable contributions of Lorraine Anderson, the research nurses in Orkney, the administrative team in Edinburgh and the people of Orkney.

POPGEN. The POPGen study was supported by the German Ministry of Education and Research (BMBF) through the National Genome Research Network (NGFN). It is currently funded by the Ministry of Science, Commerce and Transportation of the State of Schleswig-Holstein. The project has also received infrastructure support through the DFG excellence cluster "Inflammation at Interfaces" and the BMBF funded project P2N (01EY1103).

PREVEND. *Prevention of Renal and Vascular Endstage Disease Study.* PREVEND genetics is supported by the Dutch Kidney Foundation (Grant E033), the EU project grant GENECURE (FP-6 LSHM CT 2006 037697), the National Institutes of Health (grant 2R01LM010098), The Netherlands organisation for health research and development (NWO-Groot grant 175.010.2007.006, NWO VENI grant 916.761.70, ZonMw grant 90.700.441), and the Dutch Inter University Cardiology Institute Netherlands (ICIN).

PROSPER-PHASE. *Prospective Study of Pravastatin in the Elderly at Risk.* The PROSPER trial was supported by an investigator initiated grant from Bristol-Myers Squibb, USA. The study was conducted, analysed, and reported independently of the company.

RS-I and II. *The Rotterdam Study.* The GWA study was funded by the Netherlands Organisation of Scientific Research NWO Investments (nr. 175.010.2005.011, 911-03-012), the Research Institute for Diseases in the Elderly (014-93-015; RIDE2), the Netherlands Genomics Initiative (NGI)/Netherlands Consortium for Healthy Aging (NCHA) project nr. 050-060-810. We thank Pascal Arp, Mila Jhamai, Dr Michael Moorhouse, Marijn Verkerk, and Sander Bervoets for their help in creating the GWAS database. The Rotterdam Study is funded by Erasmus Medical Center and Erasmus University, Rotterdam, Netherlands Organization for the Health Research and Development (ZonMw), the Research Institute for Diseases in the Elderly (RIDE), the Ministry of Education, Culture and Science, the Ministry for Health, Welfare and Sports, the European Commission (DG XII), and the Municipality of Rotterdam. The authors are very grateful to the participants and staff from the Rotterdam Study, the participating general practitioners and the pharmacists. We would like to thank Dr. Tobias A. Knoch, Luc V. de Zeeuw, Anis Abuseiris, and Rob de Graaf as well as their institutions the Erasmus Computing Grid, Rotterdam, The Netherlands, and especially the national German MediGRID and Services@MediGRID part of the German D-Grid, both funded by the German Bundesministerium fuer Forschung und Technology under grants #01 AK 803 A-H and # 01 IG 07015 G, for access to their grid resources. Abbas Dehghan is supported by NWO grant (vici, 918-76-619).

SAPALDIA. *Swiss Study on Air Pollution and Lung Diseases in Adults.* The SAPALDIA Team: Study directorate: T Rochat (p), NM Probst Hensch (e/g), N Künzli (e/exp), C Schindler (s), JM Gaspoz (c) Scientific team: JC Barthélémy (c), W Berger (g), R Bettschart (p), A Bircher (a), O Brändli (p), C Brombach (n), M Brutsche (p), L Burdet (p), M Frey (p), U Frey (pd), MW Gerbase (p), D Gold (e/c/p), E de Groot (c), W Karrer (p), R Keller (p), B Martin (pa), D Miedinger (o), U Neu (exp), L Nicod (p), M Pons (p), F Roche (c), T Rothe (p), E Russi (p), P Schmid-Grendelmeyer

(a), A Schmidt-Trucksäss (pa), A Turk (p), J Schwartz (e), D. Stolz (p), P Straehl (exp), JM Tschopp (p), A von Eckardstein (cc), E Zemp Stutz (e). Scientific team at coordinating centers: M Adam (e/g), C Autenrieth (pa), PO Bridevaux (p), D Carballo (c), E Corradi (exp), I Curjuric (e), J Dratva (e), A Di Pasquale (s), E Dupuis Lozeron (s), E Fischer (e), M Germond (s), L Grize (s), D Keidel (s), S Kriemler (pa), A Kumar (g), M Imboden (g), N Maire (s), A Mehta (e), H Phuleria (exp), E Schaffner (s), GA Thun (g) A Ineichen (exp), M Ragetti (e), M Ritter (exp), T Schikowski (e), M Tarantino (s), M Tsai (exp) (a) allergology, (c) cardiology, (cc) clinical chemistry, (e) epidemiology, (exp) exposure, (g) genetic and molecular biology, (m) meteorology, (n) nutrition, (o) occupational health, (p) pneumology, (pa) physical activity, (pd) pediatrics, (s) statistics. Funding: The Swiss National Science Foundation (grants no 33CSCO-134276/1, 33CSCO-108796, 3247BO-104283, 3247BO-104288, 3247BO-104284, 3247-065896, 3100-059302, 3200-052720, 3200-042532, 4026-028099), the Federal Office for Forest, Environment and Landscape, the Federal Office of Public Health, the Federal Office of Roads and Transport, the canton's government of Aargau, Basel-Stadt, Basel-Land, Geneva, Luzern, Ticino, Valais, and Zürich, the Swiss Lung League, the canton's Lung League of Basel Stadt/ Basel Landschaft, Geneva, Ticino, Valais and Zurich, SUVA, Freiwillige Akademische Gesellschaft, UBS Wealth Foundation, Talecris Biotherapeutics GmbH, Abbott Diagnostics, European Commission 018996 (GABRIEL), Wellcome Trust WT 084703MA. The study could not have been done without the help of the study participants, technical and administrative support and the medical teams and field workers at the local study sites. Local fieldworkers : Aarau: S Brun, G Giger, M Sperisen, M Stahel, Basel: C Bürl, C Dahler, N Oertli, I Harreh, F Karrer, G Novicic, N Wyttenbacher, Davos: A Saner, P Senn, R Winzeler, Geneva: F Bonfils, B Blicharz, C Landolt, J Rochat, Lugano: S Boccia, E Gehrig, MT Mandia, G Solari, B Viscardi, Montana: AP Bieri, C Darioly, M Maire, Payerne: F Ding, P Danieli A Vonnez, Wald: D Bodmer, E Hochstrasser, R Kunz, C Meier, J Rakic, U Schafroth, A Walder. Administrative staff: C Gabriel, R Gutknecht.

SAPHIR. *Salzburg Atherosclerosis Prevention program in subjects at High Individual Risk.* The SAPHIR-study was partially supported by a grant from the Kamillo Eisner Stiftung to B. Paulweber and by grants from the "Genomics of Lipid-associated Disorders – GOLD" of the "Austrian Genome Research Programme GEN-AU" to F. Kronenberg.

SHIP and SHIP-TREND. *The Study of Health in Pomerania.* SHIP is part of the Community Medicine Research net of the University of Greifswald, Germany, which is funded by the Federal Ministry of Education and Research (grants no. 01ZZ9603, 01ZZ0103, and 01ZZ0403), the Ministry of Cultural Affairs as well as the Social Ministry of the Federal State of Mecklenburg-West Pomerania, and the network 'Greifswald Approach to Individualized Medicine (GANI_MED)' funded by the Federal Ministry of Education and Research (grant 03IS2061A). Genome-wide data have been supported by the Federal Ministry of Education and Research (grant no. 03ZIK012) and a joint grant from Siemens Healthcare, Erlangen, Germany and the Federal State of Mecklenburg- West Pomerania. The University of Greifswald is a member of the 'Center of Knowledge Interchange' program of the Siemens AG and the Caché Campus program of the InterSystems GmbH. The SHIP authors are grateful to Mario Stanke for the opportunity to use his Server Cluster for the SNP imputation as well as to Holger Prokisch and Thomas Meitinger (Helmholtz Zentrum München) for the genotyping of the SHIP-TREND cohort.

SIGNET-Sea Islands and SIGNET-REGARDS. The *Sea Islands Genetics Network* is supported by R01 DK084350 (M Sale), and consists of data from the REasons for Geographic And Racial Differences in Stroke (REGARDS) cohort (U01 NS041588; G Howard), Project SuGAR (Sea Islands Genetic African American Registry; W.M. Keck Foundation; WT Garvey), and a South Carolina Center of Biomedical Research Excellence (COBRE) for Oral Health Project P20-RR-17696 (JK Fernandes).

SKIPOGH. *The Swiss Kidney Project on Genes in Hypertension.* This research was funded by a grant from the Swiss National Science Foundation (33CM30-124087). The study also received support from Lausanne University Hospital, Geneva University Hospital and Bern University Hospital, Switzerland. Murielle Bochud received support from the Swiss School of Public Health Plus (SSPH+).

SORBS. *The Sorbs study.* This work was supported by grants from the German Research Council (SFB- 1052 "Obesity mechanisms" to Michael Stumvoll, Anke Tönjes and Peter Kovacs), from the German Diabetes Association (to Anke Tönjes and Peter Kovacs) and from the DHFD (Diabetes Hilfs- und Forschungsfonds Deutschland to Michael Stumvoll and Peter Kovacs). IFB AdiposityDiseases is supported by the Federal Ministry of Education and Research (BMBF), Germany, FKZ: 01EO1001. We thank all those who participated in the study. Sincere thanks are

given to Knut Krohn (Microarray Core Facility of the Interdisciplinary Centre for Clinical Research, University of Leipzig) for the genotyping support.

Vanderbilt. Illumina HumanOmni1-Quad and Illumina HumanOmni5-Quad genotyping were supported by NIGMS RC2-GM092318. Illumina Human660W-Quad genotyping was supported by the eMERGE Network, initiated and funded by the National Human Genome Research Institute, with additional funding from the National Institute of General Medical Sciences (NIGMS), through U01-HG04603.

WGHS. *Women's Genome Health Study.* The WGHS is supported by HL 043851 and HL69757 from the National Heart, Lung, and Blood Institute and CA 047988 from the National Cancer Institute, the Donald W. Reynolds Foundation and the Fondation Leducq, with collaborative scientific support and funding for genotyping provided by Amgen.

YFS. *Young Finns Study.* The YFS has been financially supported by the Academy of Finland: grants 134309 (Eye), 126925, 121584, 124282, 129378 (Salve), 117787 (Gendi), and 41071 (Skidi), the Social Insurance Institution of Finland, Kuopio, Tampere and Turku University Hospital Medical Funds (grant 9M048 and 9N035 for TeLeht), Juho Vainio Foundation, Paavo Nurmi Foundation, Finnish Foundation of Cardiovascular Research and Finnish Cultural Foundation, Tampere Tuberculosis Foundation and Emil Aaltonen Foundation (T.L.). The expert technical assistance in the statistical analyses by Ville Aalto and Irina Lisinen is acknowledged.

References

1. Freedman, M. L. *et al.* Assessing the impact of population stratification on genetic association studies. *Nat. Genet.* **36**, 388-393 (2004).
2. de Bakker, P. I. *et al.* Practical aspects of imputation-driven meta-analysis of genome-wide association studies. *Hum. Mol. Genet.* **17**, R122-8 (2008).
3. Pattaro, C. *et al.* Genome-wide association and functional follow-up reveals new loci for kidney function. *PLoS Genet.* **8**, e1002584 (2012).
4. Kottgen, A. *et al.* New loci associated with kidney function and chronic kidney disease. *Nat. Genet.* **42**, 376-384 (2010).
5. Kottgen, A. *et al.* Multiple loci associated with indices of renal function and chronic kidney disease. *Nat. Genet.* **41**, 712-717 (2009).
6. Chasman, D. I. *et al.* Integration of genome-wide association studies with biological knowledge identifies six novel genes related to kidney function. *Hum. Mol. Genet.* **21**, 5329-5343 (2012).
7. Johnson, A. D. *et al.* SNAP: a web-based tool for identification and annotation of proxy SNPs using HapMap. *Bioinformatics* **24**, 2938-2939 (2008).
8. Schunkert, H. *et al.* Large-scale association analysis identifies 13 new susceptibility loci for coronary artery disease. *Nat. Genet.* **43**, 333-338 (2011).
9. Vasan, R. S. *et al.* Genetic variants associated with cardiac structure and function: a meta-analysis and replication of genome-wide association data. *JAMA* **302**, 168-178 (2009).
10. Smith, N. L. *et al.* Association of genome-wide variation with the risk of incident heart failure in adults of European and African ancestry: a prospective meta-analysis from the cohorts for heart and aging research in genomic epidemiology (CHARGE) consortium. *Circ. Cardiovasc. Genet.* **3**, 256-266 (2010).
11. International Consortium for Blood Pressure Genome-Wide Association Studies *et al.* Genetic variants in novel pathways influence blood pressure and cardiovascular disease risk. *Nature* **478**, 103-109 (2011).
12. Dupuis, J. *et al.* New genetic loci implicated in fasting glucose homeostasis and their impact on type 2 diabetes risk. *Nat. Genet.* **42**, 105-116 (2010).
13. Schizophrenia Psychiatric Genome-Wide Association Study (GWAS) Consortium. Genome-wide association study identifies five new schizophrenia loci. *Nat. Genet.* **43**, 969-976 (2011).
14. Berndt, S. I. *et al.* Genome-wide meta-analysis identifies 11 new loci for anthropometric traits and provides insights into genetic architecture. *Nat. Genet.* **45**, 501-512 (2013).
15. Thorleifsson, G. *et al.* Genome-wide association yields new sequence variants at seven loci that associate with measures of obesity. *Nat. Genet.* **41**, 18-24 (2009).
16. Speliotes, E. K. *et al.* Association analyses of 249,796 individuals reveal 18 new loci associated with body mass index. *Nat. Genet.* **42**, 937-948 (2010).
17. Jostins, L. *et al.* Host-microbe interactions have shaped the genetic architecture of inflammatory bowel disease. *Nature* **491**, 119-124 (2012).
18. Mells, G. F. *et al.* Genome-wide association study identifies 12 new susceptibility loci for primary biliary cirrhosis. *Nat. Genet.* **43**, 329-332 (2011).
19. Okada, Y. *et al.* Meta-analysis identifies multiple loci associated with kidney function-related traits in east Asian populations. *Nat. Genet.* **44**, 904-909 (2012).
20. Kottgen, A. *et al.* Genome-wide association analyses identify 18 new loci associated with serum urate concentrations. *Nat. Genet.* **45**, 145-154 (2013).
21. Paternoster, L. *et al.* Meta-analysis of genome-wide association studies identifies three new risk loci for atopic dermatitis. *Nat. Genet.* **44**, 187-192 (2011).
22. Teslovich, T. M. *et al.* Biological, clinical and population relevance of 95 loci for blood lipids. *Nature* **466**, 707-713 (2010).
23. Yang, Q. *et al.* Multiple genetic loci influence serum urate levels and their relationship with gout and cardiovascular disease risk factors. *Circ. Cardiovasc. Genet.* **3**, 523-530 (2010).
24. van Meurs, J. B. *et al.* Common genetic loci influencing plasma homocysteine concentrations and their effect on risk of coronary artery disease. *Am. J. Clin. Nutr.* **98**, 668-676 (2013).
25. Barrett, J. H. *et al.* Genome-wide association study identifies three new melanoma susceptibility loci. *Nat. Genet.* **43**, 1108-1113 (2011).
26. Bishop, D. T. *et al.* Genome-wide association study identifies three loci associated with melanoma risk. *Nat. Genet.* **41**, 920-925 (2009).
27. Han, J. *et al.* A genome-wide association study identifies novel alleles associated with hair color and skin pigmentation. *PLoS Genet.* **4**, e1000074 (2008).
28. Zhang, M. *et al.* Genome-wide association studies identify several new loci associated with pigmentation traits and skin cancer risk in European Americans. *Hum. Mol. Genet.* **22**, 2948-2959 (2013).
29. Nan, H. *et al.* Genome-wide association study identifies novel alleles associated with risk of cutaneous basal cell carcinoma and squamous cell carcinoma. *Hum. Mol. Genet.* **20**, 3718-3724 (2011).

30. Sulem, P. *et al.* Genetic determinants of hair, eye and skin pigmentation in Europeans. *Nat. Genet.* **39**, 1443-1452 (2007).
31. Siddiq, A. *et al.* A meta-analysis of genome-wide association studies of breast cancer identifies two novel susceptibility loci at 6q14 and 20q11. *Hum. Mol. Genet.* **21**, 5373-5384 (2012).
32. Sulem, P. *et al.* Two newly identified genetic determinants of pigmentation in Europeans. *Nat. Genet.* **40**, 835-837 (2008).
33. Brown, K. M. *et al.* Common sequence variants on 20q11.22 confer melanoma susceptibility. *Nat. Genet.* **40**, 838-840 (2008).
34. Morris, A. P. Transethnic meta-analysis of genomewide association studies. *Genet. Epidemiol.* **35**, 809-822 (2011).
35. Zhang, B. *et al.* Integrated systems approach identifies genetic nodes and networks in late-onset Alzheimer's disease. *Cell* **153**, 707-720 (2013).
36. Greenawalt, D. M. *et al.* A survey of the genetics of stomach, liver, and adipose gene expression from a morbidly obese cohort. *Genome Res.* **21**, 1008-1016 (2011).
37. Schadt, E. E. *et al.* Mapping the genetic architecture of gene expression in human liver. *PLoS Biol.* **6**, e107 (2008).
38. Innocenti, F. *et al.* Identification, replication, and functional fine-mapping of expression quantitative trait loci in primary human liver tissue. *PLoS Genet.* **7**, e1002078 (2011).
39. Schroder, A. *et al.* Genomics of ADME gene expression: mapping expression quantitative trait loci relevant for absorption, distribution, metabolism and excretion of drugs in human liver. *Pharmacogenomics J.* **13**, 12-20 (2013).
40. Fehrmann, R. S. *et al.* Trans-eQTLs reveal that independent genetic variants associated with a complex phenotype converge on intermediate genes, with a major role for the HLA. *PLoS Genet.* **7**, e1002197 (2011).
41. Grundberg, E. *et al.* Mapping cis- and trans-regulatory effects across multiple tissues in twins. *Nat. Genet.* **44**, 1084-1089 (2012).
42. Dimas, A. S. *et al.* Common regulatory variation impacts gene expression in a cell type-dependent manner. *Science* **325**, 1246-1250 (2009).
43. Zeller, T. *et al.* Genetics and beyond--the transcriptome of human monocytes and disease susceptibility. *PLoS One* **5**, e10693 (2010).
44. Emilsson, V. *et al.* Genetics of gene expression and its effect on disease. *Nature* **452**, 423-428 (2008).
45. Hao, K. *et al.* Lung eQTLs to help reveal the molecular underpinnings of asthma. *PLoS Genet.* **8**, e1003029 (2012).
46. Kabachiev, B. & Silverberg, M. S. Expression quantitative trait loci analysis identifies associations between genotype and gene expression in human intestine. *Gastroenterology* **144**, 1488-96, 1496.e1-3 (2013).
47. Liang, L. *et al.* A cross-platform analysis of 14,177 expression quantitative trait loci derived from lymphoblastoid cell lines. *Genome Res.* **23**, 716-726 (2013).
48. Degner, J. F. *et al.* DNase I sensitivity QTLs are a major determinant of human expression variation. *Nature* **482**, 390-394 (2012).
49. Li, Q. *et al.* Integrative eQTL-based analyses reveal the biology of breast cancer risk loci. *Cell* **152**, 633-641 (2013).
50. Zou, F. *et al.* Brain expression genome-wide association study (eGWAS) identifies human disease-associated variants. *PLoS Genet.* **8**, e1002707 (2012).
51. Goring, H. H. *et al.* Discovery of expression QTLs using large-scale transcriptional profiling in human lymphocytes. *Nat. Genet.* **39**, 1208-1216 (2007).
52. Zhu, M. *et al.* Beta-mannosidosis mice: a model for the human lysosomal storage disease. *Hum. Mol. Genet.* **15**, 493-500 (2006).
53. Gill, P. S. & Rosenblum, N. D. Control of murine kidney development by sonic hedgehog and its GLI effectors. *Cell. Cycle* **5**, 1426-1430 (2006).
54. Lu, B. C. *et al.* Etv4 and Etv5 are required downstream of GDNF and Ret for kidney branching morphogenesis. *Nat. Genet.* **41**, 1295-1302 (2009).
55. Kozak, E. M. & Tate, S. S. Glutathione-degrading enzymes of microvillus membranes. *J. Biol. Chem.* **257**, 6322-6327 (1982).
56. Campbell, B. J., Forrester, L. J., Zahler, W. L. & Burks, M. Beta-lactamase activity of purified and partially characterized human renal dipeptidase. *J. Biol. Chem.* **259**, 14586-14590 (1984).
57. Mazumder, B. *et al.* Regulated release of L13a from the 60S ribosomal subunit as a mechanism of transcript-specific translational control. *Cell* **115**, 187-198 (2003).
58. Koppen, M., Metodiev, M. D., Casari, G., Rugarli, E. I. & Langer, T. Variable and tissue-specific subunit composition of mitochondrial m-AAA protease complexes linked to hereditary spastic paraplegia. *Mol. Cell. Biol.* **27**, 758-767 (2007).
59. Elleuch, N. *et al.* Mutation analysis of the paraplegin gene (SPG7) in patients with hereditary spastic paraplegia. *Neurology* **66**, 654-659 (2006).
60. Arnoldi, A. *et al.* A clinical, genetic, and biochemical characterization of SPG7 mutations in a large cohort of patients with hereditary spastic paraplegia. *Hum. Mutat.* **29**, 522-531 (2008).

61. Sanchez-Ferrero, E. *et al.* SPG7 mutational screening in spastic paraplegia patients supports a dominant effect for some mutations and a pathogenic role for p.A510V. *Clin. Genet.* **83**, 257-262 (2013).
62. Tekin, M. *et al.* The KBG syndrome: confirmation of autosomal dominant inheritance and further delineation of the phenotype. *Am. J. Med. Genet. A.* **130A**, 284-287 (2004).
63. Sirmaci, A. *et al.* Mutations in ANKRD11 cause KBG syndrome, characterized by intellectual disability, skeletal malformations, and macrodontia. *Am. J. Hum. Genet.* **89**, 289-294 (2011).
64. Li, X. *et al.* Cloning and chromosomal localization of the human and murine genes for the T-cell transcription factors NFATc and NFATp. *Cytogenet. Cell Genet.* **68**, 185-191 (1995).
65. Langworthy, M., Zhou, B., de Caestecker, M., Moeckel, G. & Baldwin, H. S. NFATc1 identifies a population of proximal tubule cell progenitors. *J. Am. Soc. Nephrol.* **20**, 311-321 (2009).
66. Wang, Y. *et al.* Activation of NFAT signaling in podocytes causes glomerulosclerosis. *J. Am. Soc. Nephrol.* **21**, 1657-1666 (2010).
67. Varon, R. *et al.* Partial deficiency of the C-terminal-domain phosphatase of RNA polymerase II is associated with congenital cataracts facial dysmorphism neuropathy syndrome. *Nat. Genet.* **35**, 185-189 (2003).
68. Ahuja, R. *et al.* Interactions of cubilin with megalin and the product of the amnionless gene (AMN): effect on its stability. *Biochem. J.* **410**, 301-308 (2008).
69. Kantarci, S. *et al.* Mutations in LRP2, which encodes the multiligand receptor megalin, cause Donnai-Barrow and facio-oculo-acoustico-renal syndromes. *Nat. Genet.* **39**, 957-959 (2007).
70. Attakpa, E. S., Djibril, N. M., Baba-Moussa, F., Yessoufou, G. & Sezan, A. Expression and role of the genes involved in the transport of bile acids in the liver and kidneys in mice. *J. Basic Clin. Physiol. Pharmacol.* **24**, 97-103 (2013).
71. Parr, B. A. & McMahon, A. P. Sexually dimorphic development of the mammalian reproductive tract requires Wnt-7a. *Nature* **395**, 707-710 (1998).
72. Hirata, H. *et al.* Polymorphisms of DNA repair genes are associated with renal cell carcinoma. *Biochem. Biophys. Res. Commun.* **342**, 1058-1062 (2006).
73. Key, M. D., Andres, D. A., Der, C. J. & Repasky, G. A. Characterization of RERG: an estrogen-regulated tumor suppressor gene. *Methods Enzymol.* **407**, 513-527 (2006).
74. Ozaltin, F. *et al.* Disruption of PTPRO causes childhood-onset nephrotic syndrome. *Am. J. Hum. Genet.* **89**, 139-147 (2011).
75. Wharram, B. L. *et al.* Altered podocyte structure in GLEPP1 (Ptpro)-deficient mice associated with hypertension and low glomerular filtration rate. *J. Clin. Invest.* **106**, 1281-1290 (2000).
76. Zwaenepoel, I. *et al.* Ezrin regulates microvillus morphogenesis by promoting distinct activities of Eps8 proteins. *Mol. Biol. Cell* **23**, 1080-1094 (2012).
77. Garrett, S. H. *et al.* Short and long term gene expression variation and networking in human proximal tubule cells when exposed to cadmium. *BMC Med. Genomics* **6 Suppl 1**, S2-S794-6-S1-S2. Epub 2013 Jan 23 (2013).
78. Cheng, S. B. *et al.* Anatomical location and redistribution of G protein-coupled estrogen receptor-1 during the estrus cycle in mouse kidney and specific binding to estrogens but not aldosterone. *Mol. Cell. Endocrinol.* **382**, 950-959 (2014).
79. Prabhushankar, R., Krueger, C. & Manrique, C. Membrane estrogen receptors: their role in blood pressure regulation and cardiovascular disease. *Curr. Hypertens. Rep.* **16**, 408-013-0408-6 (2014).
80. Kurt, A. H. & Buyukafsar, K. Vasoconstriction induced by G1, a G-protein-coupled oestrogen receptor1 (GPER-1) agonist, in the isolated perfused rat kidney. *Eur. J. Pharmacol.* **702**, 71-78 (2013).
81. Barton, M. Position paper: The membrane estrogen receptor GPER--Clues and questions. *Steroids* **77**, 935-942 (2012).
82. Osborne, M. R. *et al.* The reaction of 7,8-dihydro-7,8-dihydroxybenzo (a) pyrene-9,10-oxide with DNA in relation to the benzo (a) pyrene-DNA products isolated from cells. *Chem. Biol. Interact.* **13**, 343-348 (1976).
83. Huang, W. *et al.* High glucose induces activation of NF-kappaB inflammatory signaling through IkappaBalpha sumoylation in rat mesangial cells. *Biochem. Biophys. Res. Commun.* **438**, 568-574 (2013).
84. Pandey, D. *et al.* SUMO1 negatively regulates reactive oxygen species production from NADPH oxidases. *Arterioscler. Thromb. Vasc. Biol.* **31**, 1634-1642 (2011).
85. Zhang, L. *et al.* A high-throughput screen for chemical inhibitors of exocytic transport in yeast. *Chembiochem* **11**, 1291-1301 (2010).
86. Zhang, L., Huang, M. & Harsay, E. A chemical genetic screen for modulators of exocytic transport identifies inhibitors of a transport mechanism linked to GTR2 function. *Eukaryot. Cell.* **9**, 116-126 (2010).
87. Nowak, J. *et al.* The TP53INP2 protein is required for autophagy in mammalian cells. *Mol. Biol. Cell* **20**, 870-881 (2009).
88. Teng, A., Nair, M., Wells, J., Segre, J. A. & Dai, X. Strain-dependent perinatal lethality of Ovol1-deficient mice and identification of Ovol2 as a downstream target of Ovol1 in skin epidermis. *Biochim. Biophys. Acta* **1772**, 89-95 (2007).
89. Lee, S. D. *et al.* TonEBP stimulates multiple cellular pathways for adaptation to hypertonic stress: organic osmolyte-dependent and -independent pathways. *Am. J. Physiol. Renal Physiol.* **300**, F707-15 (2011).
90. Davis, L. K., Rodgers, B. D. & Kelley, K. M. Angiotensin II- and glucose-stimulated extracellular matrix production: mediation by the insulin-like growth factor (IGF) axis in a murine mesangial cell line. *Endocrine* **33**, 32-39 (2008).

91. Fukasawa, H. *et al.* Ubiquitin-dependent degradation of SnoN and Ski is increased in renal fibrosis induced by obstructive injury. *Kidney Int.* **69**, 1733-1740 (2006).
92. Kirk, A., Campbell, S., Bass, P., Mason, J. & Collins, J. Differential expression of claudin tight junction proteins in the human cortical nephron. *Nephrol. Dial. Transplant.* **25**, 2107-2119 (2010).
93. Kugita, M. *et al.* Global gene expression profiling in early-stage polycystic kidney disease in the Han:SPRD Cy rat identifies a role for RXR signaling. *Am. J. Physiol. Renal Physiol.* **300**, F177-88 (2011).
94. Ng, M. C. *et al.* Transferability and fine mapping of type 2 diabetes loci in African Americans: the Candidate Gene Association Resource Plus Study. *Diabetes* **62**, 965-976 (2013).
95. Lim, X. L. *et al.* KCNQ1 SNPS and susceptibility to diabetic nephropathy in East Asians with type 2 diabetes. *Diabetologia* **55**, 2402-2406 (2012).
96. Liu, C. T. *et al.* Genetic association for renal traits among participants of African ancestry reveals new loci for renal function. *PLoS Genet.* **7**, e1002264 (2011).
97. 3C Study Group. Vascular factors and risk of dementia: design of the Three-City Study and baseline characteristics of the study population. *Neuroepidemiology* **22**, 316-325 (2003).
98. Lambert, J. C. *et al.* Genome-wide association study identifies variants at CLU and CR1 associated with Alzheimer's disease. *Nat. Genet.* **41**, 1094-1099 (2009).
99. Patel, A. *et al.* Effects of a fixed combination of perindopril and indapamide on macrovascular and microvascular outcomes in patients with type 2 diabetes mellitus (the ADVANCE trial): a randomised controlled trial. *Lancet* **370**, 829-840 (2007).
100. Harris, T. B. *et al.* Age, Gene/Environment Susceptibility-Reykjavik Study: multidisciplinary applied phenomics. *Am. J. Epidemiol.* **165**, 1076-1087 (2007).
101. Mitchell, B. D. *et al.* The genetic response to short-term interventions affecting cardiovascular function: rationale and design of the Heredity and Phenotype Intervention (HAPI) Heart Study. *Am. Heart J.* **155**, 823-828 (2008).
102. Rampersaud, E. *et al.* The association of coronary artery calcification and carotid artery intima-media thickness with distinct, traditional coronary artery disease risk factors in asymptomatic adults. *Am. J. Epidemiol.* **168**, 1016-1023 (2008).
103. The ARIC Investigators. The Atherosclerosis Risk in Communities (ARIC) Study: design and objectives. The ARIC investigators. *Am. J. Epidemiol.* **129**, 687-702 (1989).
104. Schmidt, R. *et al.* Assessment of cerebrovascular risk profiles in healthy persons: definition of research goals and the Austrian Stroke Prevention Study (ASPS). *Neuroepidemiology* **13**, 308-313 (1994).
105. Schmidt, R., Fazekas, F., Kapeller, P., Schmidt, H. & Hartung, H. P. MRI white matter hyperintensities: three-year follow-up of the Austrian Stroke Prevention Study. *Neurology* **53**, 132-139 (1999).
106. Middelberg, R. P. *et al.* Loci affecting gamma-glutamyl transferase in adults and adolescents show age x SNP interaction and cardiometabolic disease associations. *Hum. Mol. Genet.* **21**, 446-455 (2012).
107. Shock, N. W., *et al.* Normal Human Aging: The Baltimore Longitudinal Study of Aging. *Superintendent of Documents, U.S. Government Printing Office, Washington, DC 20402.* (1984).
108. Mitchell, P., Smith, W., Attebo, K. & Wang, J. J. Prevalence of age-related maculopathy in Australia. The Blue Mountains Eye Study. *Ophthalmology* **102**, 1450-1460 (1995).
109. Attebo, K., Mitchell, P. & Smith, W. Visual acuity and the causes of visual loss in Australia. The Blue Mountains Eye Study. *Ophthalmology* **103**, 357-364 (1996).
110. Leeder, S. R. *et al.* Low hemoglobin, chronic kidney disease, and risk for coronary heart disease-related death: the Blue Mountains Eye Study. *J. Am. Soc. Nephrol.* **17**, 279-284 (2006).
111. Fried, L. P. *et al.* The Cardiovascular Health Study: design and rationale. *Ann. Epidemiol.* **1**, 263-276 (1991).
112. Heard-Costa, N. L. *et al.* NRXN3 is a novel locus for waist circumference: a genome-wide association study from the CHARGE Consortium. *PLoS Genet.* **5**, e1000539 (2009).
113. Polasek, O. *et al.* Genome-wide association study of anthropometric traits in Korcula Island, Croatia. *Croat. Med. J.* **50**, 7-16 (2009).
114. Rudan, I. *et al.* "10001 Dalmatians:" Croatia launches its national biobank. *Croat. Med. J.* **50**, 4-6 (2009).
115. Rudan, I., Campbell, H. & Rudan, P. Genetic epidemiological studies of eastern Adriatic Island isolates, Croatia: objective and strategies. *Coll. Antropol.* **23**, 531-546 (1999).
116. Rudan, I. *et al.* Effects of inbreeding, endogamy, genetic admixture, and outbreeding on human health: a (1001 Dalmatians) study. *Croat. Med. J.* **47**, 601-610 (2006).
117. Guder, W. G. *et al.* Multicentre evaluation of an enzymatic method for creatinine determination using a sensitive colour reagent. *J. Clin. Chem. Clin. Biochem.* **24**, 889-902 (1986).
118. Balkau, B. An epidemiologic survey from a network of French Health Examination Centres, (D.E.S.I.R.): epidemiologic data on the insulin resistance syndrome. *Rev. Epidemiol. Sante Publique* **44**, 373-375 (1996).
119. Metspalu, A. The Estonian Genome Project. *Drug Dev. Res.* **62**, 97-101 (2004).
120. Nelis, M. *et al.* Genetic structure of Europeans: a view from the North-East. *PLoS One* **4**, e5472 (2009).
121. Aulchenko, Y. S. *et al.* Linkage disequilibrium in young genetically isolated Dutch population. *Eur. J. Hum. Genet.* **12**, 527-534 (2004).
122. Higgins, M. *et al.* NHLBI Family Heart Study: objectives and design. *Am. J. Epidemiol.* **143**, 1219-1228 (1996).

123. Kannel, W. B., Feinleib, M., McNamara, P. M., Garrison, R. J. & Castelli, W. P. An investigation of coronary heart disease in families. The Framingham offspring study. *Am. J. Epidemiol.* **110**, 281-290 (1979).
124. Feinleib, M., Kannel, W. B., Garrison, R. J., McNamara, P. M. & Castelli, W. P. The Framingham Offspring Study. Design and preliminary data. *Prev. Med.* **4**, 518-525 (1975).
125. Splansky, G. L. *et al.* The Third Generation Cohort of the National Heart, Lung, and Blood Institute's Framingham Heart Study: design, recruitment, and initial examination. *Am. J. Epidemiol.* **165**, 1328-1335 (2007).
126. Daniels, P. R. *et al.* Familial aggregation of hypertension treatment and control in the Genetic Epidemiology Network of Arteriopathy (GENOA) study. *Am. J. Med.* **116**, 676-681 (2004).
127. Turner, S. T. *et al.* Influence of genomic loci on measures of chronic kidney disease in hypertensive sibships. *J. Am. Soc. Nephrol.* **17**, 2048-2055 (2006).
128. Rule, A. D. *et al.* A comparison of serum creatinine-based methods for identifying chronic kidney disease in hypertensive individuals and their siblings. *Am. J. Hypertens.* **19**, 608-614 (2006).
129. McEvoy, M. *et al.* Cohort profile: The Hunter Community Study. *Int. J. Epidemiol.* **39**, 1452-1463 (2010).
130. Qi, L. *et al.* Genetic variants at 2q24 are associated with susceptibility to type 2 diabetes. *Hum. Mol. Genet.* **19**, 2706-2715 (2010).
131. Rimm, E. B. *et al.* Prospective study of alcohol consumption and risk of coronary disease in men. *Lancet.* **338**, 464-468 (1991).
132. Hu, F. B. *et al.* Physical activity and television watching in relation to risk for type 2 diabetes mellitus in men. *Arch. Intern. Med.* **161**, 1542-1548 (2001).
133. Manson, J. E. *et al.* Physical activity and incidence of non-insulin-dependent diabetes mellitus in women. *Lancet* **338**, 774-778 (1991).
134. Lin, J., Hu, F. B. & Curhan, G. Serum adiponectin and renal dysfunction in men with type 2 diabetes. *Diabetes Care* **30**, 239-244 (2007).
135. Salvi, E. *et al.* Genomewide association study using a high-density single nucleotide polymorphism array and case-control design identifies a novel essential hypertension susceptibility locus in the promoter region of endothelial NO synthase. *Hypertension* **59**, 248-255 (2012).
136. Gambaro, G. *et al.* Prevalence of CKD in northeastern Italy: results of the INCIPE study and comparison with NHANES. *Clin. J. Am. Soc. Nephrol.* **5**, 1946-1953 (2010).
137. D'Adamo, P. *et al.* Metabonomics and population studies: age-related amino acids excretion and inferring networks through the study of urine samples in two Italian isolated populations. *Amino Acids* **38**, 65-73 (2010).
138. Tepper, B. J. *et al.* Variation in the bitter-taste receptor gene TAS2R38, and adiposity in a genetically isolated population in Southern Italy. *Obesity (Silver Spring)* **16**, 2289-2295 (2008).
139. Sala, C. *et al.* Variation of hemoglobin levels in normal Italian populations from genetic isolates. *Haematologica* **93**, 1372-1375 (2008).
140. Ciullo, M. *et al.* New susceptibility locus for hypertension on chromosome 8q by efficient pedigree-breaking in an Italian isolate. *Hum. Mol. Genet.* **15**, 1735-1743 (2006).
141. Colonna, V. *et al.* Campora: a young genetic isolate in South Italy. *Hum. Hered.* **64**, 123-135 (2007).
142. Ciullo, M. *et al.* Identification and replication of a novel obesity locus on chromosome 1q24 in isolated populations of Cilento. *Diabetes* **57**, 783-790 (2008).
143. Colonna, V. *et al.* Comparing population structure as inferred from genealogical versus genetic information. *Eur. J. Hum. Genet.* **17**, 1635-1641 (2009).
144. Siervo, M. *et al.* Angiogenesis and biomarkers of cardiovascular risk in adults with metabolic syndrome. *J. Intern. Med.* **268**, 338-347 (2010).
145. Sorice, R. *et al.* Association of a variant in the CHRNA5-A3-B4 gene cluster region to heavy smoking in the Italian population. *Eur. J. Hum. Genet.* **19**, 593-596 (2011).
146. Ruggiero, D. *et al.* Genetics of VEGF serum variation in human isolated populations of Cilento: importance of VEGF polymorphisms. *PLoS One* **6**, e16982 (2011).
147. Siervo, M. *et al.* Body mass index is directly associated with biomarkers of angiogenesis and inflammation in children and adolescents. *Nutrition* **28**, 262-266 (2012).
148. Traglia, M. *et al.* Heritability and demographic analyses in the large isolated population of Val Borbera suggest advantages in mapping complex traits genes. *PLoS One* **4**, e7554 (2009).
149. Ridker, P. M. *et al.* Rosuvastatin to prevent vascular events in men and women with elevated C-reactive protein. *N. Engl. J. Med.* **359**, 2195-2207 (2008).
150. Chasman, D. I. *et al.* Genetic determinants of statin-induced low-density lipoprotein cholesterol reduction: the Justification for the Use of Statins in Prevention: an Intervention Trial Evaluating Rosuvastatin (JUPITER) trial. *Circ. Cardiovasc. Genet.* **5**, 257-264 (2012).
151. Wichmann, H. E., Gieger, C., Illig, T. & MONICA/KORA Study Group. KORA-gen--resource for population genetics, controls and a broad spectrum of disease phenotypes. *Gesundheitswesen* **67 Suppl 1**, S26-30 (2005).
152. Baumeister, S. E. *et al.* Effect of chronic kidney disease and comorbid conditions on health care costs: A 10-year observational study in a general population. *Am. J. Nephrol.* **31**, 222-229 (2010).
153. Bild, D. E. *et al.* Multi-ethnic study of atherosclerosis: objectives and design. *Am. J. Epidemiol.* **156**, 871-881 (2002).

154. Pattaro, C. *et al.* The genetic study of three population microisolates in South Tyrol (MICROS): study design and epidemiological perspectives. *BMC Med. Genet.* **8**, 29 (2007).
155. Marroni, F., Grazio, D., Pattaro, C., Devoto, M. & Pramstaller, P. Estimates of genetic and environmental contribution to 43 quantitative traits support sharing of a homogeneous environment in an isolated population from South Tyrol, Italy. *Hum. Hered.* **65**, 175-182 (2008).
156. Pattaro, C. *et al.* A meta-analysis of genome-wide data from five European isolates reveals an association of COL22A1, SYT1, and GABRR2 with serum creatinine level. *BMC Med. Genet.* **11**, 41 (2010).
157. Penninx, B. W. *et al.* The Netherlands Study of Depression and Anxiety (NESDA): rationale, objectives and methods. *Int. J. Methods Psychiatr. Res.* **17**, 121-140 (2008).
158. Sullivan, P. F. *et al.* Genome-wide association for major depressive disorder: a possible role for the presynaptic protein piccolo. *Mol. Psychiatry* **14**, 359-375 (2009).
159. Colditz, G. A. & Hankinson, S. E. The Nurses' Health Study: lifestyle and health among women. *Nat. Rev. Cancer.* **5**, 388-396 (2005).
160. Lin, J., Fung, T. T., Hu, F. B. & Curhan, G. C. Association of dietary patterns with albuminuria and kidney function decline in older white women: a subgroup analysis from the Nurses' Health Study. *Am. J. Kidney Dis.* **57**, 245-254 (2011).
161. Johansson, A., Vavrouch-Nilsson, V., Edin-Liljegren, A., Sjolander, P. & Gyllensten, U. Linkage disequilibrium between microsatellite markers in the Swedish Sami relative to a worldwide selection of populations. *Hum. Genet.* **116**, 105-113 (2005).
162. Johansson, A., Vavrouch-Nilsson, V., Cox, D. R., Frazer, K. A. & Gyllensten, U. Evaluation of the SNP tagging approach in an independent population sample--array-based SNP discovery in Sami. *Hum. Genet.* **122**, 141-150 (2007).
163. Portas, L. *et al.* History, geography and population structure influence the distribution and heritability of blood and anthropometric quantitative traits in nine Sardinian genetic isolates. *Genet. Res. (Camb)* **92**, 199-208 (2010).
164. Biino, G. *et al.* Genetic architecture of hand quantitative ultrasound measures: a population-based study in a Sardinian genetic isolate. *Bone* **46**, 1197-1203 (2010).
165. Pistis, G. *et al.* High differentiation among eight villages in a secluded area of Sardinia revealed by genome-wide high density SNPs analysis. *PLoS One* **4**, e4654 (2009).
166. McQuillan, R. *et al.* Runs of homozygosity in European populations. *Am. J. Hum. Genet.* **83**, 359-372 (2008).
167. Krawczak, M. *et al.* PopGen: population-based recruitment of patients and controls for the analysis of complex genotype-phenotype relationships. *Community Genet.* **9**, 55-61 (2006).
168. Hillege, H. L. *et al.* Urinary albumin excretion predicts cardiovascular and noncardiovascular mortality in general population. *Circulation* **106**, 1777-1782 (2002).
169. Shepherd, J. *et al.* The design of a prospective study of Pravastatin in the Elderly at Risk (PROSPER). PROSPER Study Group. PROspective Study of Pravastatin in the Elderly at Risk. *Am. J. Cardiol.* **84**, 1192-1197 (1999).
170. Shepherd, J. *et al.* Pravastatin in elderly individuals at risk of vascular disease (PROSPER): a randomised controlled trial. *Lancet* **360**, 1623-1630 (2002).
171. Ford, I. *et al.* Reduced glomerular filtration rate and its association with clinical outcome in older patients at risk of vascular events: secondary analysis. *PLoS Med.* **6**, e16 (2009).
172. Hofman, A., Grobbee, D. E., de Jong, P. T. & van den Ouweland, F. A. Determinants of disease and disability in the elderly: the Rotterdam Elderly Study. *Eur. J. Epidemiol.* **7**, 403-422 (1991).
173. Hofman, A. *et al.* The Rotterdam Study: objectives and design update. *Eur. J. Epidemiol.* **22**, 819-829 (2007).
174. Hofman, A. *et al.* The Rotterdam Study: 2010 objectives and design update. *Eur. J. Epidemiol.* **24**, 553-572 (2009).
175. John, U. *et al.* Study of Health In Pomerania (SHIP): a health examination survey in an east German region: objectives and design. *Soz. Praventivmed.* **46**, 186-194 (2001).
176. Volzke, H. *et al.* Cohort profile: the study of health in Pomerania. *Int. J. Epidemiol.* **40**, 294-307 (2011).
177. Tonjes, A. *et al.* Genetic variation in GPR133 is associated with height: genome wide association study in the self-contained population of Sorbs. *Hum. Mol. Genet.* **18**, 4662-4668 (2009).
178. Tonjes, A. *et al.* Association of FTO variants with BMI and fat mass in the self-contained population of Sorbs in Germany. *Eur. J. Hum. Genet.* **18**, 104-110 (2010).
179. Veeramah, K. R. *et al.* Genetic variation in the Sorbs of eastern Germany in the context of broader European genetic diversity. *Eur. J. Hum. Genet.* **19**, 995-1001 (2011).
180. Ridker, P. M. *et al.* Rationale, design, and methodology of the Women's Genome Health Study: a genome-wide association study of more than 25,000 initially healthy american women. *Clin. Chem.* **54**, 249-255 (2008).
181. Guessous, I. *et al.* Caffeine intake and CYP1A2 variants associated with high caffeine intake protect non-smokers from hypertension. *Hum. Mol. Genet.* **21**, 3283-3292 (2012).
182. Guessous, I., Bochud, M., Theler, J. M., Gaspoz, J. M. & Pechere-Bertschi, A. 1999-2009 Trends in prevalence, unawareness, treatment and control of hypertension in Geneva, Switzerland. *PLoS One* **7**, e39877 (2012).
183. Weck, M. N., Stegmaier, C., Rothenbacher, D. & Brenner, H. Epidemiology of chronic atrophic gastritis: population-based study among 9444 older adults from Germany. *Aliment. Pharmacol. Ther.* **26**, 879-887 (2007).

184. Raum, E. *et al.* Changes of cardiovascular risk factors and their implications in subsequent birth cohorts of older adults in Germany: a life course approach. *Eur. J. Cardiovasc. Prev. Rehabil.* **14**, 809-814 (2007).
185. Schottker, B. *et al.* Strong associations of 25-hydroxyvitamin D concentrations with all-cause, cardiovascular, cancer, and respiratory disease mortality in a large cohort study. *Am. J. Clin. Nutr.* **97**, 782-793 (2013).
186. Boger, C. A. *et al.* Effect of ACE and AT-2 inhibitors on mortality and progression to microalbuminuria in a nested case-control study of diabetic nephropathy in diabetes mellitus type 2: results from the GENDIAN study. *Int. J. Clin. Pharmacol. Ther.* **44**, 364-374 (2006).
187. Boger, C. A. *et al.* Association of eGFR-Related Loci Identified by GWAS with Incident CKD and ESRD. *PLoS Genet.* **7**, e1002292 (2011).
188. Wild, P. S. *et al.* Distribution and categorization of left ventricular measurements in the general population: results from the population-based Gutenberg Heart Study. *Circ. Cardiovasc. Imaging* **3**, 604-613 (2010).
189. Wild, P. S. *et al.* A genome-wide association study identifies LIPA as a susceptibility gene for coronary artery disease. *Circ. Cardiovasc. Genet.* **4**, 403-412 (2011).
190. Lucae, S. *et al.* P2RX7, a gene coding for a purinergic ligand-gated ion channel, is associated with major depressive disorder. *Hum. Mol. Genet.* **15**, 2438-2445 (2006).
191. Kloiber, S. *et al.* Variations in tryptophan hydroxylase 2 linked to decreased serotonergic activity are associated with elevated risk for metabolic syndrome in depression. *Mol. Psychiatry* **15**, 736-747 (2010).
192. Kohli, M. A. *et al.* The neuronal transporter gene SLC6A15 confers risk to major depression. *Neuron* **70**, 252-265 (2011).
193. Tayo, B. O. *et al.* Genetic background of patients from a university medical center in Manhattan: implications for personalized medicine. *PLoS One* **6**, e19166 (2011).
194. Winkelmann, B. R. *et al.* Rationale and design of the LURIC study--a resource for functional genomics, pharmacogenomics and long-term prognosis of cardiovascular disease. *Pharmacogenomics* **2**, S1-73 (2001).
195. Heid, I. M. *et al.* Genetic architecture of the APM1 gene and its influence on adiponectin plasma levels and parameters of the metabolic syndrome in 1,727 healthy Caucasians. *Diabetes* **55**, 375-384 (2006).
196. Kollerits, B. *et al.* A common variant in the adiponutrin gene influences liver enzyme values. *J. Med. Genet.* **47**, 116-119 (2010).
197. Pruijm, M. *et al.* Heritability, determinants and reference values of renal length: a family-based population study. *Eur. Radiol.* **23**, 2899-2905 (2013).